

D

Treatment modalities



Table D.1 . 97

Percentages & counts of reported ESRD patients: by treatment modality
incident & point prevalent patients

Table D.2 . 99

Counts of incident ESRD patients, by initial mode of therapy: all dialysis
by age, gender, race, ethnicity, & primary diagnosis

Table D.3 . 101

Counts of incident ESRD patients, by initial mode of therapy: hemodialysis
by age, gender, race, ethnicity, & primary diagnosis

Table D.4 . 103

Counts of incident ESRD patients, by initial mode of therapy: peritoneal dialysis
by age, gender, race, ethnicity, & primary diagnosis

Table D.5 . 105

Counts of incident ESRD patients, by initial mode of therapy: transplant
by age, gender, race, ethnicity, & primary diagnosis

Table D.6 . 107

Counts of point prevalent ESRD patients, by mode of therapy: all dialysis
December 31 point prevalent patients,
by age, gender, race, ethnicity, & primary diagnosis

Table D.7 . 109

Counts of point prevalent ESRD patients, by mode of therapy: hemodialysis
December 31 point prevalent patients,
by age, gender, race, ethnicity, & primary diagnosis

Table D.8 . 111

Counts of point prevalent ESRD patients, by mode of therapy: peritoneal dialysis
December 31 point prevalent patients,
by age, gender, race, ethnicity, & primary diagnosis

Table D.9 . 113

Counts of point prevalent ESRD patients, by mode of therapy:
functioning transplant
December 31 point prevalent patients,
by age, gender, race, ethnicity, & primary diagnosis

Table D.10 . 115

Incident ESRD patients, 2005
by age, gender, race, ethnicity, primary diagnosis, & modality

Table D.11 . 116

Point prevalent ESRD patients, 2005
December 31 point prevalent patients,
by age, gender, race, ethnicity, primary diagnosis, & modality

Table D.12 . 117

Treatment status two years after first ESRD service, 2001–2003 combined
incident patients with Medicare coverage

Table D.13 . 118

Point prevalent ESRD patients, 2005: by patient vintage
December 31 point prevalent patients, by year-end treatment modality
& number of years patient has had ESRD (vintage)

Table D.14 . 119

Point prevalent ESRD patients, 2005: by number of years on modality
December 31 point prevalent patients, by year-end treatment modality

Table D.15 . 120

Incident ESRD patients, 2005: by modality & state/territory

Table D.16 . 121

Point prevalent ESRD patients, 2005: by modality & state/territory
December 31 point prevalent ESRD patients-

Throughout this reference section

The incident cohorts are determined at the time of ESRD initiation without applying the 60-day stable modality rule.
The age of the incident patient is determined as of the date of ESRD initiation.
The point prevalent cohorts are determined on December 31 of each calendar year without applying the 60-day stable modality rule.
The age of the point prevalent patient is determined as of December 31 of each calendar year.

The following symbols are used throughout this reference section

† CMS began collecting Hispanic ethnicity data in April 1995; cells for years prior to 1996 have therefore been left blank.
* Values for cells with ten or fewer patients are suppressed. " " Zero values in this cell.

Table D.1

Percentages & counts of reported ESRD patients: by treatment modality*incident & point prevalent patients*

INCIDENT COUNTS	1980	1981	1982	1983	1984	1985	1986	1987	1988	1989	1990	1991	1992	1993
Modality														
Dialysis	17,643	19,749	22,139	25,504	27,086	30,005	32,714	36,000	40,229	45,379	49,772	55,035	60,161	63,379
Started with a transplant	337	326	416	405	466	537	725	900	1,037	1,033	1,096	1,077	1,104	1,074
Dialysis type														
Center hemodialysis	15,455	16,783	18,206	21,004	22,245	23,952	26,341	29,078	33,109	37,562	40,932	45,335	49,892	52,666
Center self hemodialysis	60	56	41
Home hemodialysis	161	152	158	181	210	390	372	390	459	533	569	711	831	932
CAPD	982	1,884	3,139	3,732	3,914	4,031	4,094	4,411	4,677	5,130	5,910	6,132	6,308	6,731
CCPD	*	*	*	16	38	137	224	337	315	459	607	832	1,087	1,032
Other PD	675	440	264	283	271	585	727	664	718	770	664	769	734	664
Uncertain dialysis	367	484	365	288	408	910	956	1,120	951	925	1,090	1,196	1,253	1,313
All	17,980	20,075	22,555	25,909	27,552	30,542	33,439	36,900	41,266	46,412	50,868	56,112	61,265	64,453
PERCENTAGES														
Modality														
Dialysis	98.1	98.4	98.2	98.4	98.3	98.2	97.8	97.6	97.5	97.8	97.8	98.1	98.2	98.3
Started with a transplant	1.9	1.6	1.8	1.6	1.7	1.8	2.2	2.4	2.5	2.2	2.2	1.9	1.8	1.7
Dialysis type														
Center hemodialysis	87.6	85.0	82.2	82.4	82.1	79.8	80.5	80.8	82.3	82.8	82.2	82.4	82.9	83.1
Center self hemodialysis	0.1	0.1	0.1
Home hemodialysis	0.9	0.8	0.7	0.7	0.8	1.3	1.1	1.1	1.1	1.2	1.1	1.3	1.4	1.5
CAPD	5.6	9.5	14.2	14.6	14.5	13.4	12.5	12.3	11.6	11.3	11.9	11.1	10.5	10.6
CCPD	0.0	0.0	0.0	0.1	0.1	0.5	0.7	0.9	0.8	1.0	1.2	1.5	1.8	1.6
Other PD	3.8	2.2	1.2	1.1	1.0	1.9	2.2	1.8	1.8	1.7	1.3	1.4	1.2	1.0
Uncertain dialysis	2.1	2.5	1.6	1.1	1.5	3.0	2.9	3.1	2.4	2.0	2.2	2.2	2.1	2.1
All	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0
POINT PREVALENT COUNTS														
Modality														
Dialysis	49,814	56,082	62,798	72,657	79,990	86,430	92,986	101,022	110,571	122,879	135,827	150,956	166,565	180,893
Functioning graft	10,115	12,669	15,439	18,504	22,180	26,524	31,699	36,572	41,164	45,639	50,591	55,915	61,170	66,675
Dialysis type														
Center hemodialysis	44,146	50,213	55,301	62,426	66,240	67,442	73,107	80,719	90,454	101,007	111,421	122,700	135,352	147,766
Center self hemodialysis	.	*	*	*	.	*	*	*	*	.	.	677	763	130
Home hemodialysis	2,021	1,037	895	1,503	2,726	5,814	5,541	4,765	3,186	3,134	2,967	3,202	3,460	3,829
CAPD	890	2,596	4,324	6,695	8,724	10,537	11,458	12,365	13,384	14,983	16,884	18,828	20,691	22,466
CCPD	*	17	46	72	202	539	789	1,005	1,120	1,340	1,973	2,883	3,675	4,288
Other PD	1,416	1,083	1,011	862	770	866	770	691	648	596	486	436	368	287
Uncertain dialysis	1,337	1,135	1,220	1,098	1,328	1,231	1,320	1,475	1,778	1,819	2,096	2,230	2,256	2,127
All	59,929	68,751	78,237	91,161	102,170	112,954	124,685	137,594	151,735	168,518	186,418	206,871	227,735	247,568
PERCENTAGES														
Modality														
Dialysis	83.1	81.6	80.3	79.7	78.3	76.5	74.6	73.4	72.9	72.9	72.9	73.0	73.1	73.1
Functioning graft	16.9	18.4	19.7	20.3	21.7	23.5	25.4	26.6	27.1	27.1	27.1	27.0	26.9	26.9
Dialysis type														
Center hemodialysis	88.6	89.5	88.1	85.9	82.8	78.0	78.6	79.9	81.8	82.2	82.0	81.3	81.3	81.7
Center self hemodialysis	.	0.0	0.0	0.0	.	0.0	0.0	0.0	0.0	.	.	0.4	0.5	0.1
Home hemodialysis	4.1	1.8	1.4	2.1	3.4	6.7	6.0	4.7	2.9	2.6	2.2	2.1	2.1	2.1
CAPD	1.8	4.6	6.9	9.2	10.9	12.2	12.3	12.2	12.1	12.2	12.4	12.5	12.4	12.4
CCPD	0.0	0.0	0.1	0.1	0.3	0.6	0.8	1.0	1.0	1.1	1.5	1.9	2.2	2.4
Other PD	2.8	1.9	1.6	1.2	1.0	1.0	0.8	0.7	0.6	0.5	0.4	0.3	0.2	0.2
Uncertain dialysis	2.7	2.0	1.9	1.5	1.7	1.4	1.4	1.5	1.6	1.5	1.5	1.5	1.4	1.2
All	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0



Table D.1 (continued)

Percentages & counts of reported ESRD patients: by treatment modality

incident & point prevalent patients

INCIDENT COUNTS	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005
INCIDENT COUNTS												
Modality												
Dialysis	68,808	69,157	75,028	80,513	85,789	89,714	92,771	95,971	98,022	100,644	102,607	104,488
Started with a transplant	1,090	1,221	1,301	1,315	1,396	1,572	1,734	1,816	1,936	1,996	2,246	2,424
Dialysis type												
Center hemodialysis	57,606	58,317	64,506	71,210	76,740	80,687	84,643	87,873	90,458	92,914	94,750	96,684
Center self hemodialysis	24	50	69	55	50	54	28	45	53	47	28	31
Home hemodialysis	972	734	941	441	790	978	417	384	276	432	469	428
CAPD	6,751	7,287	6,834	6,210	5,563	5,355	5,005	4,974	4,677	4,738	4,706	4,673
CCPD	1,289	2,120	2,390	2,336	2,419	2,432	2,447	2,495	2,343	2,324	2,364	2,202
Other PD	622	172	134	115	97	69	86	71	101	71	63	47
Uncertain dialysis	1,544	477	154	146	130	139	145	129	114	118	227	423
All	69,898	70,378	76,329	81,828	87,185	91,286	94,505	97,787	99,958	102,640	104,853	106,912
PERCENTAGES												
Modality												
Dialysis	98.4	98.3	98.3	98.4	98.4	98.3	98.2	98.1	98.1	98.1	97.9	97.7
Started with a transplant	1.6	1.7	1.7	1.6	1.6	1.7	1.8	1.9	1.9	1.9	2.1	2.3
Dialysis type												
Center hemodialysis	83.7	84.3	86.0	88.4	89.5	89.9	91.2	91.6	92.3	92.3	92.3	92.5
Center self hemodialysis	0.0	0.1	0.1	0.1	0.1	0.1	0.0	0.0	0.1	0.0	0.0	0.0
Home hemodialysis	1.4	1.1	1.3	0.5	0.9	1.1	0.4	0.4	0.3	0.4	0.5	0.4
CAPD	9.8	10.5	9.1	7.7	6.5	6.0	5.4	5.2	4.8	4.7	4.6	4.5
CCPD	1.9	3.1	3.2	2.9	2.8	2.7	2.6	2.6	2.4	2.3	2.3	2.1
Other PD	0.9	0.2	0.2	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.0
Uncertain dialysis	2.2	0.7	0.2	0.2	0.2	0.2	0.2	0.1	0.1	0.1	0.2	0.4
All	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0
POINT PREVALENT COUNTS												
Modality												
Dialysis	197,383	209,479	223,728	239,079	254,870	269,111	283,329	296,212	308,109	319,490	330,534	341,319
Functioning graft	72,155	77,960	83,858	89,721	96,079	102,186	108,577	115,082	121,922	128,675	136,077	143,693
Dialysis type												
Center hemodialysis	161,800	174,285	189,570	206,744	224,440	239,547	254,608	267,839	279,848	290,849	301,718	311,919
Center self hemodialysis	191	506	520	456	354	334	266	317	271	193	184	138
Home hemodialysis	3,820	3,041	2,905	2,538	2,476	2,368	2,168	1,871	1,759	1,908	2,061	2,105
CAPD	23,602	22,232	19,985	17,911	15,930	14,475	13,257	12,458	11,576	11,252	10,840	10,732
CCPD	5,394	7,796	9,521	10,367	10,545	11,245	11,886	12,713	13,679	14,388	14,720	15,163
Other PD	275	215	141	114	97	74	68	66	71	50	42	37
Uncertain dialysis	2,301	1,404	1,086	949	1,028	1,068	1,076	948	905	850	969	1,225
All	269,538	287,439	307,586	328,800	350,949	371,297	391,906	411,294	430,031	448,165	466,611	485,012
PERCENTAGES												
Modality												
Dialysis	73.2	72.9	72.7	72.7	72.6	72.5	72.3	72.0	71.6	71.3	70.8	70.4
Functioning graft	26.8	27.1	27.3	27.3	27.4	27.5	27.7	28.0	28.4	28.7	29.2	29.6
Dialysis type												
Center hemodialysis	82.0	83.2	84.7	86.5	88.1	89.0	89.9	90.4	90.8	91.0	91.3	91.4
Center self hemodialysis	0.1	0.2	0.2	0.2	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.0
Home hemodialysis	1.9	1.5	1.3	1.1	1.0	0.9	0.8	0.6	0.6	0.6	0.6	0.6
CAPD	12.0	10.6	8.9	7.5	6.3	5.4	4.7	4.2	3.8	3.5	3.3	3.1
CCPD	2.7	3.7	4.3	4.3	4.1	4.2	4.2	4.3	4.4	4.5	4.5	4.4
Other PD	0.1	0.1	0.1	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
Uncertain dialysis	1.2	0.7	0.5	0.4	0.4	0.4	0.4	0.3	0.3	0.3	0.3	0.4
All	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0

Table D.2

Counts of incident ESRD patients, by initial mode of therapy: all dialysis*by age, gender, race, ethnicity, & primary diagnosis*

	1980	1981	1982	1983	1984	1985	1986	1987	1988	1989	1990	1991	1992	1993
0-4	41	58	83	93	97	124	125	121	116	132	125	160	128	154
5-9	84	80	87	80	95	80	101	91	83	94	120	86	89	107
10-14	161	174	207	193	199	186	166	169	179	158	194	199	197	199
15-19	409	450	426	414	393	405	403	416	459	464	475	436	490	470
20-29	1,450	1,620	1,810	1,766	1,919	1,961	2,025	2,029	2,205	2,361	2,460	2,487	2,638	2,692
30-39	2,133	2,384	2,654	2,733	2,946	3,172	3,454	3,595	4,017	4,383	4,598	4,849	5,205	5,327
40-49	2,375	2,693	2,804	3,166	3,301	3,595	3,867	4,342	4,820	5,474	6,014	6,719	7,265	7,766
50-59	3,720	4,333	4,637	4,810	5,067	5,447	5,730	6,155	7,061	7,594	8,219	8,881	9,811	10,412
60-64	2,131	2,455	2,786	3,210	3,524	3,943	4,123	4,589	5,038	5,498	6,106	6,662	6,984	7,341
65-69	2,181	2,268	2,598	3,329	3,512	3,920	4,448	5,099	5,657	6,532	7,234	8,005	8,626	8,960
70-79	2,527	2,745	3,395	4,685	4,926	5,781	6,597	7,448	8,303	9,777	10,869	12,687	14,185	14,994
80+	431	487	649	1,020	1,101	1,384	1,667	1,941	2,289	2,905	3,355	3,863	4,542	4,957
Unknown	.	*	*	*	*	*	*	*	*	*	*	*	*	.
0-19	695	762	803	780	784	795	795	797	837	848	914	881	904	930
20-44	4,666	5,275	5,771	5,976	6,387	6,814	7,311	7,715	8,527	9,294	10,016	10,579	11,304	11,599
45-64	7,143	8,210	8,920	9,709	10,370	11,304	11,888	12,995	14,614	16,016	17,381	19,019	20,599	21,939
65-74	3,788	4,012	4,755	6,199	6,448	7,402	8,274	9,462	10,427	12,044	13,496	15,288	16,786	17,591
75+	1,351	1,488	1,887	2,835	3,091	3,683	4,438	5,026	5,822	7,170	7,962	9,267	10,567	11,320
Unknown	.	*	*	*	*	*	*	*	*	*	*	*	*	.
Male	9,808	10,889	12,152	13,952	14,905	16,255	17,823	19,444	21,522	24,402	26,674	29,328	31,806	33,716
Female	7,835	8,860	9,987	11,551	12,181	13,750	14,891	16,556	18,706	20,977	23,098	25,707	28,355	29,663
Unknown	.	.	.	*	*
White	12,302	13,714	15,412	17,348	18,415	20,370	22,240	24,196	26,623	29,968	32,927	36,243	39,371	40,996
African American	4,815	5,449	6,123	7,412	7,829	8,668	9,327	10,532	12,025	13,803	14,877	16,609	18,248	19,491
Native American	125	149	192	278	282	295	376	432	510	547	588	626	754	761
Asian	108	175	340	366	481	591	660	757	907	974	1,259	1,431	1,616	1,927
Other/unknown	293	262	72	100	79	81	111	83	164	87	121	126	172	204
†Hispanic														
†Non-Hispanic														
Diabetes	2,641	3,982	5,207	6,143	7,407	8,585	9,914	11,299	12,856	15,502	17,912	20,462	22,741	24,369
Hypertension	3,124	4,516	5,812	6,335	7,159	8,219	8,928	10,308	11,914	13,888	15,306	17,012	19,033	19,750
Glomerulonephritis	2,753	3,594	4,217	4,433	4,748	4,980	5,233	5,688	6,100	6,435	6,860	7,093	7,248	7,550
Cystic kidney	768	936	1,072	1,091	1,133	1,199	1,275	1,297	1,339	1,351	1,484	1,530	1,645	1,711
Other urologic	456	659	850	952	962	1,154	1,160	1,280	1,331	1,335	1,259	1,418	1,508	1,307
Other cause	1,785	2,173	2,741	3,031	3,215	3,499	3,564	3,565	3,875	4,521	4,702	5,137	5,564	6,019
Unknown cause	1,586	1,649	1,360	1,401	1,431	1,504	1,746	1,918	1,991	1,779	1,873	2,052	2,161	2,265
Missing cause	4,530	2,240	880	2,118	1,031	865	894	645	823	568	376	331	261	408
All	17,643	19,749	22,139	25,504	27,086	30,005	32,714	36,000	40,229	45,379	49,772	55,035	60,161	63,379



Table D.2 (continued)

Counts of incident ESRD patients, by initial mode of therapy: all dialysis

by age, gender, race, ethnicity, & primary diagnosis

	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005
0-4	128	149	146	146	154	147	166	170	160	166	193	187
5-9	126	111	111	95	103	120	120	118	128	102	120	111
10-14	223	217	225	231	216	240	245	248	291	290	278	262
15-19	457	493	502	464	489	526	503	502	506	541	555	530
20-29	2,668	2,562	2,556	2,522	2,588	2,471	2,549	2,541	2,432	2,391	2,452	2,576
30-39	5,638	5,436	5,567	5,516	5,512	5,685	5,502	5,461	5,475	5,454	5,480	5,556
40-49	8,196	8,698	9,284	9,352	10,134	10,301	10,542	10,599	10,852	11,149	11,417	11,552
50-59	11,707	11,755	12,410	13,563	14,736	15,906	16,476	17,323	17,721	18,610	19,201	19,902
60-64	7,789	7,942	8,358	8,842	9,445	9,594	9,959	10,351	10,425	11,092	11,424	11,644
65-69	9,610	9,280	10,053	10,722	11,049	11,160	11,433	11,796	11,711	12,042	12,156	12,022
70-79	16,623	16,571	18,758	20,692	21,991	23,159	23,919	24,805	24,888	24,891	24,840	24,929
80+	5,641	5,940	7,058	8,368	9,371	10,404	11,357	12,057	13,433	13,916	14,488	15,215
Unknown	*	*	.	.	*	*	*	*
0-19	934	970	984	936	962	1,033	1,034	1,038	1,085	1,099	1,146	1,090
20-44	11,955	11,953	12,259	12,114	12,548	12,563	12,562	12,490	12,430	12,490	12,665	13,012
45-64	24,043	24,440	25,916	27,681	29,867	31,394	32,466	33,785	34,475	36,206	37,309	38,218
65-74	19,127	18,642	20,560	22,011	22,809	23,176	23,785	24,650	24,381	24,448	24,646	24,472
75+	12,747	13,149	15,309	17,771	19,602	21,547	22,924	24,008	25,651	26,401	26,838	27,694
Unknown	*	*	.	.	*	*	*	*
Male	36,411	36,444	39,937	42,872	45,553	47,919	49,592	51,549	53,212	54,454	56,619	57,972
Female	32,397	32,709	35,091	37,641	40,236	41,793	43,177	44,422	44,810	46,182	45,980	46,502
Unknown	.	*	.	.	.	*	*	.	.	*	*	14
White	44,383	43,697	47,726	51,789	54,850	57,895	60,295	62,338	63,340	64,706	66,560	67,896
African American	21,091	21,410	22,724	23,990	25,750	26,283	26,614	27,657	28,510	29,361	29,086	30,026
Native American	831	866	931	949	1,063	1,144	1,186	1,155	1,132	1,115	1,184	1,195
Asian	2,172	2,263	2,492	2,612	2,931	3,169	3,187	3,376	3,522	3,664	3,788	3,971
Other/unknown	331	921	1,155	1,173	1,195	1,223	1,489	1,445	1,518	1,798	1,989	1,400
†Hispanic			8,718	9,035	9,683	10,636	12,424	11,679	12,396	13,182	13,511	13,583
†Non-Hispanic			66,310	71,478	76,106	79,078	80,347	84,292	85,626	87,462	89,096	90,905
Diabetes	27,164	29,032	32,477	35,529	38,046	39,975	42,115	43,760	44,044	45,237	46,081	46,409
Hypertension	21,397	18,744	19,911	21,741	23,069	24,338	24,921	26,137	27,054	28,328	28,341	28,404
Glomerulonephritis	8,137	8,040	8,429	8,532	8,670	8,640	8,332	8,279	8,341	8,067	8,166	7,710
Cystic kidney	1,787	1,804	1,738	1,864	1,972	2,031	1,971	2,018	1,980	1,996	2,027	2,157
Other urologic	1,371	1,958	2,168	2,275	2,363	2,744	2,627	2,675	2,799	2,698	2,746	2,077
Other cause	6,484	6,958	7,284	7,524	8,206	8,472	8,878	9,029	9,668	9,929	10,509	11,850
Unknown cause	2,297	2,448	2,858	2,919	3,273	3,307	3,724	3,939	3,867	4,000	4,011	4,471
Missing cause	171	173	163	129	190	207	203	134	269	389	726	1,410
All	68,808	69,157	75,028	80,513	85,789	89,714	92,771	95,971	98,022	100,644	102,607	104,488



Table D.3

Counts of incident ESRD patients, by initial mode of therapy: hemodialysis*by age, gender, race, ethnicity, & primary diagnosis*

	1980	1981	1982	1983	1984	1985	1986	1987	1988	1989	1990	1991	1992	1993
0-4	22	18	23	26	19	32	37	21	24	37	35	36	22	38
5-9	57	47	50	34	35	29	30	32	28	31	50	31	37	36
10-14	133	128	120	110	97	102	85	89	87	86	92	83	105	101
15-19	377	393	343	324	298	298	295	292	338	347	338	332	340	335
20-29	1,301	1,400	1,532	1,470	1,562	1,617	1,625	1,600	1,736	1,899	1,904	1,975	2,107	2,131
30-39	1,850	2,019	2,157	2,163	2,375	2,477	2,696	2,797	3,155	3,417	3,553	3,711	4,048	4,164
40-49	2,101	2,359	2,288	2,571	2,680	2,855	3,060	3,458	3,894	4,333	4,778	5,242	5,775	6,152
50-59	3,281	3,726	3,887	3,970	4,170	4,436	4,706	5,038	5,917	6,260	6,721	7,313	8,041	8,651
60-64	1,924	2,092	2,344	2,692	2,969	3,242	3,427	3,821	4,257	4,679	5,124	5,589	5,877	6,246
65-69	1,969	1,952	2,202	2,902	2,994	3,230	3,722	4,282	4,873	5,677	6,227	6,962	7,600	7,812
70-79	2,229	2,391	2,869	4,046	4,308	4,858	5,610	6,346	7,236	8,689	9,642	11,285	12,657	13,435
80+	372	409	547	873	944	1,161	1,413	1,688	2,021	2,635	3,034	3,546	4,170	4,538
Unknown	.	*	*	*	*	*	*	*	*	*	*	*	.	.
0-19	589	586	536	494	449	461	447	434	477	501	515	482	504	510
20-44	4,113	4,524	4,744	4,822	5,170	5,428	5,757	6,033	6,745	7,323	7,785	8,166	8,890	9,100
45-64	6,344	7,072	7,464	8,044	8,586	9,199	9,757	10,681	12,214	13,265	14,295	15,664	16,958	18,244
65-74	3,387	3,457	4,022	5,374	5,534	6,162	6,967	7,972	9,002	10,562	11,732	13,403	14,830	15,486
75+	1,183	1,295	1,596	2,447	2,712	3,087	3,778	4,344	5,128	6,439	7,171	8,390	9,597	10,299
Unknown	.	*	*	*	*	*	*	*	*	*	*	*	.	.
Male	8,760	9,461	10,178	11,668	12,353	13,254	14,583	15,912	17,982	20,431	22,244	24,430	26,767	28,508
Female	6,856	7,474	8,186	9,516	10,102	11,088	12,130	13,556	15,585	17,664	19,257	21,676	24,012	25,131
Unknown	.	.	.	*	*
White	10,762	11,571	12,595	14,168	15,031	16,125	17,796	19,384	21,728	24,662	26,877	29,691	32,531	34,030
African American	4,391	4,835	5,234	6,376	6,724	7,386	7,933	8,984	10,455	12,048	12,910	14,524	16,061	17,120
Native American	110	136	168	232	228	248	312	371	438	472	501	534	635	662
Asian	100	159	303	326	409	522	588	670	819	842	1,130	1,266	1,412	1,662
Other/unknown	253	234	64	83	63	61	84	59	128	71	83	91	140	165
†Hispanic														
†Non-Hispanic														
Diabetes	2,195	3,269	4,105	4,876	5,963	6,823	7,985	9,122	10,668	12,908	14,847	16,998	19,021	20,592
Hypertension	2,758	3,946	4,908	5,380	6,164	6,904	7,557	8,812	10,289	12,094	13,244	14,871	16,781	17,341
Glomerulonephritis	2,416	3,125	3,515	3,688	3,907	4,002	4,220	4,540	4,905	5,192	5,475	5,664	5,769	6,079
Cystic kidney	696	792	918	922	930	986	1,028	1,034	1,068	1,068	1,142	1,147	1,250	1,288
Other urologic	400	571	717	805	810	942	984	1,095	1,168	1,181	1,086	1,254	1,302	1,124
Other cause	1,574	1,881	2,309	2,492	2,641	2,808	2,889	2,840	3,209	3,748	3,870	4,191	4,608	4,978
Unknown cause	1,409	1,354	1,120	1,136	1,165	1,211	1,370	1,556	1,643	1,477	1,549	1,732	1,850	1,934
Missing cause	4,168	1,997	772	1,886	875	666	680	469	618	427	288	249	198	303
All	15,616	16,935	18,364	21,185	22,455	24,342	26,713	29,468	33,568	38,095	41,501	46,106	50,779	53,639



Table D.3 (continued)

Counts of incident ESRD patients, by initial mode of therapy: hemodialysis

by age, gender, race, ethnicity, & primary diagnosis

	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005
0-4	25	38	24	37	27	34	41	44	35	34	37	45
5-9	50	48	38	33	43	45	52	53	55	42	55	50
10-14	113	104	103	114	112	125	131	147	167	160	147	142
15-19	333	356	364	341	375	421	389	393	410	436	428	428
20-29	2,155	2,068	2,078	2,116	2,223	2,152	2,220	2,248	2,186	2,146	2,178	2,312
30-39	4,446	4,321	4,538	4,641	4,685	4,896	4,825	4,803	4,833	4,844	4,865	4,942
40-49	6,628	6,990	7,694	7,975	8,850	9,039	9,364	9,464	9,857	10,077	10,342	10,549
50-59	9,707	9,800	10,571	11,724	12,995	14,228	14,863	15,599	16,154	17,005	17,565	18,175
60-64	6,605	6,759	7,323	7,902	8,575	8,725	9,107	9,536	9,614	10,284	10,625	10,803
65-69	8,462	8,129	9,021	9,742	10,150	10,314	10,579	10,976	10,903	11,222	11,372	11,293
70-79	14,909	14,978	17,139	19,142	20,577	21,745	22,572	23,475	23,610	23,687	23,624	23,720
80+	5,168	5,508	6,623	7,939	8,967	9,994	10,945	11,564	12,963	13,456	14,006	14,682
Unknown	*	*	.	.	*	*	*	*
0-19	521	546	529	525	557	625	613	637	667	672	667	665
20-44	9,534	9,585	10,025	10,194	10,799	10,897	11,024	11,036	11,144	11,187	11,270	11,672
45-64	20,007	20,353	22,179	24,164	26,529	28,143	29,355	30,614	31,500	33,169	34,305	35,109
65-74	16,924	16,539	18,541	20,121	21,067	21,506	22,180	23,083	22,889	22,955	23,165	23,065
75+	11,615	12,076	14,242	16,702	18,627	20,547	21,916	22,932	24,587	25,410	25,837	26,630
Unknown	*	*	.	.	*	*	*	*
Male	30,951	31,065	34,872	38,208	41,239	43,714	45,480	47,508	49,349	50,566	52,585	53,995
Female	27,651	28,033	30,644	33,498	36,341	38,004	39,608	40,794	41,438	42,820	42,657	43,142
Unknown	.	*	.	.	.	*	.	.	.	*	*	*
White	37,081	36,530	40,805	45,276	48,843	51,953	54,640	56,750	58,118	59,578	61,333	62,757
African American	18,678	19,100	20,718	22,222	24,065	24,689	25,104	26,103	27,020	27,780	27,588	28,541
Native American	711	751	819	856	960	1,059	1,074	1,063	1,049	1,038	1,110	1,116
Asian	1,852	1,922	2,160	2,305	2,627	2,903	2,905	3,057	3,197	3,338	3,472	3,626
Other/unknown	280	798	1,014	1,047	1,085	1,115	1,365	1,329	1,403	1,659	1,744	1,103
†Hispanic			7,732	8,173	8,828	9,774	11,469	10,879	11,527	12,328	12,686	12,811
†Non-Hispanic			57,784	63,533	68,752	71,945	73,619	77,423	79,260	81,065	82,561	84,332
Diabetes	23,055	24,717	28,245	31,632	34,463	36,443	38,751	40,473	40,909	42,196	43,170	43,589
Hypertension	18,934	16,652	18,110	20,003	21,427	22,733	23,380	24,585	25,517	26,725	26,730	26,857
Glomerulonephritis	6,517	6,461	6,895	7,138	7,370	7,480	7,183	7,167	7,346	7,035	7,056	6,715
Cystic kidney	1,351	1,365	1,333	1,465	1,610	1,640	1,619	1,631	1,649	1,630	1,671	1,736
Other urologic	1,161	1,694	1,928	2,045	2,149	2,523	2,400	2,453	2,580	2,507	2,529	1,951
Other cause	5,487	5,935	6,352	6,701	7,392	7,671	8,137	8,229	8,939	9,199	9,762	11,063
Unknown cause	1,976	2,143	2,520	2,622	3,016	3,055	3,450	3,649	3,631	3,762	3,773	4,205
Missing cause	121	134	133	100	153	174	168	115	216	339	556	1,027
All	58,602	59,101	65,516	71,706	77,580	81,719	85,088	88,302	90,787	93,393	95,247	97,143



Table D.4

Counts of incident ESRD patients, by initial mode of therapy: peritoneal dialysis*by age, gender, race, ethnicity, & primary diagnosis*

	1980	1981	1982	1983	1984	1985	1986	1987	1988	1989	1990	1991	1992	1993
0-4	*	28	50	57	68	82	79	86	83	70	72	98	91	99
5-9	15	22	34	41	54	44	61	51	46	52	51	44	45	54
10-14	13	29	71	74	91	73	73	74	81	56	85	87	80	82
15-19	15	31	63	74	76	84	83	99	82	107	106	84	127	111
20-29	73	143	220	249	301	275	330	323	378	357	443	409	431	465
30-39	138	241	416	501	487	558	590	629	688	787	851	965	981	970
40-49	152	232	431	529	543	586	640	695	760	897	1,029	1,235	1,246	1,353
50-59	240	439	628	749	790	795	811	882	891	1,060	1,193	1,270	1,444	1,489
60-64	108	240	366	455	475	549	512	572	608	624	790	829	871	864
65-69	84	212	327	372	425	431	458	526	513	608	761	761	742	867
70-79	119	224	451	530	517	571	542	659	696	773	910	986	1,083	1,128
80+	19	49	88	116	123	120	139	151	166	196	226	196	253	281
Unknown	.	.	*	*	*	.	.	*	.	*	.	.	*	.
0-19	52	110	218	246	289	283	296	310	292	285	314	313	343	346
20-44	281	501	847	1,004	1,039	1,099	1,233	1,310	1,441	1,570	1,814	2,021	2,028	2,077
45-64	430	794	1,214	1,479	1,557	1,664	1,650	1,791	1,884	2,155	2,492	2,687	2,945	3,064
65-74	162	367	611	701	754	779	784	939	943	1,053	1,331	1,359	1,399	1,562
75+	60	118	255	317	311	343	355	397	432	524	566	584	679	714
Unknown	.	.	*	*	*	.	.	*	.	*	.	.	*	.
Male	515	942	1,653	1,993	2,161	2,209	2,376	2,537	2,680	3,017	3,502	3,822	3,966	4,103
Female	470	948	1,493	1,755	1,791	1,959	1,942	2,211	2,312	2,572	3,015	3,142	3,429	3,660
Unknown
White	793	1,462	2,347	2,770	2,930	3,190	3,265	3,568	3,757	4,117	4,849	5,120	5,401	5,575
African American	158	401	752	902	917	877	935	1,065	1,089	1,308	1,479	1,615	1,708	1,868
Native American	12	*	18	43	46	43	57	52	67	61	74	77	92	83
Asian	*	*	27	27	55	49	53	58	71	99	103	133	173	222
Other/unknown	15	*	*	*	*	*	*	*	*	*	12	19	21	15
†Hispanic														
†Non-Hispanic														
Diabetes	254	529	951	1,131	1,249	1,341	1,453	1,652	1,675	2,049	2,424	2,724	2,975	3,049
Hypertension	214	403	791	849	847	916	926	1,003	1,187	1,320	1,625	1,630	1,706	1,867
Glomerulonephritis	207	342	583	655	728	771	792	868	948	1,005	1,114	1,197	1,239	1,232
Cystic kidney	46	105	131	148	179	171	197	209	217	230	281	321	334	379
Other urologic	23	62	106	125	123	140	117	121	117	104	132	120	146	142
Other cause	134	213	362	483	506	539	508	570	515	598	669	710	747	816
Unknown cause	90	184	190	204	208	183	223	206	204	191	225	222	226	244
Missing cause	17	52	32	153	112	107	102	119	129	92	47	40	22	34
All	985	1,890	3,146	3,748	3,952	4,168	4,318	4,748	4,992	5,589	6,517	6,964	7,395	7,763



Table D.4 (continued)

Counts of incident ESRD patients, by initial mode of therapy: peritoneal dialysis

by age, gender, race, ethnicity, & primary diagnosis

	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005
0-4	84	102	116	105	125	109	125	122	120	128	149	136
5-9	70	62	70	60	56	72	65	64	70	56	64	59
10-14	97	107	116	112	99	109	110	98	122	129	127	114
15-19	105	129	134	120	109	103	109	105	92	103	119	92
20-29	430	455	455	383	353	310	311	275	233	239	270	246
30-39	1,006	1,058	995	844	794	772	660	637	619	591	589	582
40-49	1,312	1,609	1,546	1,328	1,250	1,218	1,139	1,101	975	1,042	1,028	938
50-59	1,644	1,853	1,787	1,803	1,692	1,642	1,565	1,682	1,536	1,562	1,569	1,600
60-64	930	1,097	1,009	918	851	844	839	803	789	795	773	773
65-69	827	1,073	1,007	957	884	828	821	802	782	804	751	685
70-79	1,216	1,479	1,574	1,502	1,379	1,384	1,311	1,299	1,232	1,172	1,173	1,148
80+	319	382	415	414	390	396	397	481	450	441	458	502
Unknown	.	*
0-19	356	400	436	397	389	393	409	389	404	416	459	401
20-44	2,047	2,228	2,156	1,846	1,687	1,619	1,487	1,405	1,239	1,266	1,341	1,260
45-64	3,275	3,844	3,636	3,430	3,253	3,167	3,027	3,093	2,913	2,963	2,888	2,879
65-74	1,592	1,967	1,967	1,839	1,709	1,634	1,558	1,533	1,450	1,466	1,425	1,327
75+	770	967	1,029	1,034	944	974	971	1,049	1,014	951	957	1,008
Unknown	.	*
Male	4,257	5,007	4,899	4,504	4,184	4,088	3,985	3,916	3,746	3,783	3,878	3,719
Female	3,783	4,400	4,325	4,042	3,798	3,699	3,467	3,553	3,274	3,279	3,192	3,156
Unknown
White	5,716	6,734	6,719	6,335	5,850	5,799	5,494	5,445	5,070	4,988	5,082	4,941
African American	1,905	2,131	1,938	1,708	1,635	1,555	1,467	1,511	1,452	1,550	1,446	1,432
Native American	99	110	111	90	99	80	109	91	78	76	73	75
Asian	291	329	327	299	297	259	273	315	322	324	312	339
Other/unknown	29	103	129	114	101	94	109	107	98	124	157	88
†Hispanic			959	844	824	843	935	779	846	835	805	756
†Non-Hispanic			8,265	7,702	7,158	6,944	6,517	6,690	6,174	6,227	6,265	6,119
Diabetes	3,258	4,065	4,135	3,814	3,520	3,479	3,306	3,227	3,065	2,989	2,868	2,776
Hypertension	1,881	1,948	1,744	1,693	1,601	1,562	1,499	1,524	1,501	1,576	1,575	1,524
Glomerulonephritis	1,347	1,504	1,505	1,359	1,264	1,133	1,121	1,084	975	1,011	1,090	977
Cystic kidney	381	411	385	381	349	378	339	366	323	354	346	408
Other urologic	138	241	223	220	204	214	212	216	210	182	206	125
Other cause	782	948	900	789	786	769	707	775	704	705	717	755
Unknown cause	234	274	318	280	249	242	259	272	223	229	233	258
Missing cause	19	16	14	*	*	*	*	*	19	16	35	52
All	8,040	9,407	9,224	8,546	7,982	7,787	7,452	7,469	7,020	7,062	7,070	6,875

Table D.5

Counts of incident ESRD patients, by initial mode of therapy: transplant*by age, gender, race, ethnicity, & primary diagnosis*

	1980	1981	1982	1983	1984	1985	1986	1987	1988	1989	1990	1991	1992	1993
0-4	*	11	17	12	14	15	11	25	28	25	25	20	26	31
5-9	12	17	18	*	15	23	19	23	22	34	31	41	37	27
10-14	18	14	15	20	31	28	44	50	46	48	52	49	49	44
15-19	24	25	31	28	35	37	54	65	62	59	67	54	41	59
20-29	86	74	90	85	88	124	130	154	212	213	176	162	154	142
30-39	89	97	107	116	129	134	220	241	278	266	289	296	294	269
40-49	66	50	79	84	83	98	134	173	198	220	259	277	269	245
50-59	29	31	55	40	55	54	78	124	146	120	141	128	168	174
60-64	*	*	*	*	*	*	18	28	29	30	33	35	41	50
65-69	*	*	.	*	*	*	*	11	*	12	14	12	18	21
70-79	.	*	.	*	*	*	*	*	*	*	*	*	*	11
80+	*	*	*	.	*	*	*	*	.	*	*	.	*	.
Unknown	*
0-19	60	67	81	69	95	103	128	163	158	166	175	164	153	161
20-44	205	198	240	244	265	320	422	491	604	600	624	628	602	546
45-64	68	58	94	89	93	98	158	229	259	249	274	270	324	334
65-74	*	*	.	*	*	11	13	15	16	19	14	23	29	*
75+	*	*	*	.	*	*	*	*	*	*	*	*	*	*
Unknown	*
Male	219	220	269	256	297	312	436	528	619	601	638	636	616	625
Female	118	105	147	149	169	225	289	372	418	432	458	441	488	449
Unknown	.	*
White	260	258	320	298	364	430	604	718	796	873	903	942	945	945
African American	53	51	66	55	58	57	75	113	136	109	119	92	113	91
Native American	.	*	.	*	*	*	*	11	*	*	*	*	*	*
Asian	*	*	11	13	*	11	*	26	31	25	34	19	27	23
Other/unknown	22	13	19	31	30	30	31	38	63	24	30	14	*	*
†Hispanic														
†Non-Hispanic														
Diabetes	38	59	89	70	100	116	181	192	209	253	236	272	238	260
Hypertension	29	25	37	34	26	30	44	78	82	61	74	71	103	73
Glomerulonephritis	53	56	80	85	91	121	122	181	163	194	223	198	226	202
Cystic kidney	*	13	*	20	19	26	45	72	68	77	95	99	103	99
Other urologic	25	25	14	21	29	24	47	42	42	64	56	64	41	57
Other cause	46	38	55	46	75	68	95	120	113	139	149	144	164	167
Unknown cause	18	16	41	43	38	59	57	61	73	59	77	76	79	115
Missing cause	126	94	92	86	88	93	134	154	287	186	186	153	150	101
All	337	326	416	405	466	537	725	900	1,037	1,033	1,096	1,077	1,104	1,074



Table D.5 (continued)

Counts of incident ESRD patients, by initial mode of therapy: transplant

by age, gender, race, ethnicity, & primary diagnosis

	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005
0-4	16	19	38	31	20	34	28	33	20	32	41	31
5-9	35	35	33	37	31	34	33	33	52	46	30	37
10-14	35	62	46	50	55	59	64	61	48	47	44	75
15-19	70	75	77	67	53	82	71	81	77	83	92	92
20-29	152	134	135	142	149	168	159	166	148	153	178	148
30-39	273	311	279	268	313	275	316	293	325	279	311	322
40-49	272	287	362	322	367	418	441	459	472	471	512	529
50-59	160	198	226	261	287	328	397	441	493	535	607	674
60-64	47	57	73	82	66	86	117	131	159	169	208	248
65-69	20	30	26	47	45	57	70	82	94	128	140	174
70-79	*	13	*	*	*	29	37	35	48	52	81	88
80+	*	*	*	*	.	*	*	*
Unknown	*
0-19	156	191	194	185	159	209	196	208	197	208	207	235
20-44	565	585	606	577	643	647	698	676	692	633	719	694
45-64	339	402	469	498	539	628	732	814	905	974	1,097	1,227
65-74	28	41	31	52	51	85	105	110	133	172	209	239
75+	*	*	*	*	*	*	*	*	*	*	14	29
Unknown	*
Male	605	687	780	790	792	907	989	1,043	1,096	1,143	1,264	1,381
Female	484	532	521	525	604	665	745	773	840	853	982	1,042
Unknown	*	*	*
White	884	947	1,021	1,134	1,243	1,381	1,493	1,589	1,664	1,682	1,777	1,851
African American	163	217	218	125	97	133	142	139	177	192	190	202
Native American	*	*	12	*	*	*	11	*	*	14	*	*
Asian	30	30	40	29	32	30	52	51	64	67	58	65
Other/unknown	*	17	*	17	15	21	36	27	22	41	212	299
†Hispanic			78	78	96	96	118	113	131	137	139	126
†Non-Hispanic			1,223	1,237	1,300	1,476	1,616	1,703	1,805	1,859	2,107	2,298
Diabetes	250	283	316	293	284	313	291	366	389	371	419	442
Hypertension	84	72	65	87	77	86	112	138	179	167	198	218
Glomerulonephritis	200	253	264	246	270	296	329	341	380	357	401	390
Cystic kidney	87	128	147	122	190	170	218	250	270	305	297	338
Other urologic	61	102	79	105	76	74	117	107	140	125	129	81
Other cause	169	186	198	208	203	207	232	293	294	291	318	374
Unknown cause	87	62	75	104	68	129	98	114	131	96	100	119
Missing cause	152	135	157	150	228	297	337	207	153	284	384	462
All	1,090	1,221	1,301	1,315	1,396	1,572	1,734	1,816	1,936	1,996	2,246	2,424

Table D.6

Counts of point prevalent ESRD patients, by mode of therapy: all dialysis*December 31 point prevalent patients, by age, gender, race, ethnicity, & primary diagnosis*

	1980	1981	1982	1983	1984	1985	1986	1987	1988	1989	1990	1991	1992	1993
0-4	42	64	91	116	130	161	173	165	163	178	179	196	196	207
5-9	112	107	131	157	164	135	158	162	172	187	200	171	177	185
10-14	272	298	328	363	351	324	321	336	361	343	391	403	403	412
15-19	844	852	898	846	840	882	852	879	955	1,022	1,017	1,027	1,076	1,103
20-29	4,045	4,302	4,631	4,979	5,218	5,346	5,397	5,546	5,934	6,368	6,707	6,927	7,340	7,616
30-39	6,424	7,150	7,952	8,773	9,567	10,193	10,737	11,343	12,183	13,353	14,280	15,403	16,558	17,617
40-49	7,867	8,571	9,299	10,736	11,584	12,149	12,761	13,728	15,230	17,184	19,145	21,500	23,608	25,794
50-59	11,825	13,233	14,563	15,859	16,921	17,783	18,499	19,257	20,687	22,423	24,370	26,786	29,557	32,218
60-64	6,083	7,243	8,256	9,699	10,817	11,746	12,306	13,377	14,265	15,412	16,810	18,083	19,418	20,725
65-69	5,516	6,197	7,006	8,563	9,682	10,761	12,153	13,740	15,145	16,897	18,669	20,623	22,657	24,417
70-79	5,914	6,962	8,251	10,556	12,291	13,955	15,975	18,045	20,242	23,227	26,451	30,830	34,889	38,336
80+	868	1,099	1,386	2,001	2,415	2,983	3,640	4,437	5,229	6,279	7,603	9,004	10,686	12,263
Unknown	*	*	*	*	*	12	14	*	*	*	*	*	.	.
0-19	1,270	1,321	1,448	1,482	1,485	1,502	1,504	1,542	1,651	1,730	1,787	1,797	1,852	1,907
20-44	14,142	15,387	16,882	18,793	20,172	21,206	22,245	23,405	25,390	27,947	30,417	32,732	35,190	37,362
45-64	22,102	25,112	27,819	31,253	33,935	36,011	37,455	39,846	42,909	46,793	50,895	55,967	61,291	66,608
65-74	9,420	10,709	12,339	15,204	17,263	19,261	21,787	24,442	27,059	30,456	34,282	38,854	43,310	47,165
75+	2,878	3,549	4,304	5,916	7,125	8,438	9,981	11,780	13,557	15,947	18,441	21,603	24,922	27,851
Unknown	*	*	*	*	*	12	14	*	*	*	*	*	.	.
Male	27,112	30,142	33,632	38,693	42,553	45,595	48,592	52,401	57,125	63,472	70,193	78,131	86,038	93,712
Female	22,702	25,940	29,166	33,963	37,437	40,835	44,394	48,621	53,445	59,407	65,634	72,825	80,527	87,181
Unknown	.	.	.	*	*
White	33,466	37,145	41,172	47,052	51,056	54,543	58,004	62,205	67,083	73,687	80,670	88,921	97,086	103,974
African American	15,013	17,361	19,867	23,528	26,568	29,162	31,892	35,294	39,386	44,543	49,785	55,772	62,191	68,408
Native American	294	384	476	640	757	863	1,000	1,158	1,331	1,503	1,696	1,908	2,213	2,469
Asian	319	449	668	898	1,142	1,454	1,710	2,031	2,429	2,893	3,421	4,080	4,745	5,667
Other/unknown	722	743	615	539	467	408	380	334	342	253	255	275	330	375
†Hispanic														
†Non-Hispanic														
Diabetes	4,900	6,807	9,122	11,541	14,383	17,035	19,892	23,250	27,157	32,603	38,618	45,523	52,688	59,132
Hypertension	8,496	10,928	13,874	16,873	19,641	22,159	24,936	28,001	31,562	36,201	40,729	45,954	51,624	56,393
Glomerulonephritis	10,177	11,597	13,197	15,196	16,704	17,954	18,971	20,123	21,383	22,937	24,347	25,872	27,337	28,975
Cystic kidney	3,217	3,649	4,139	4,696	5,035	5,330	5,486	5,675	5,798	5,974	6,194	6,486	6,728	7,012
Other urologic	1,203	1,519	1,919	2,328	2,705	3,071	3,273	3,588	3,876	4,083	4,204	4,478	4,739	4,740
Other cause	5,002	5,811	6,991	8,248	9,212	10,063	10,661	11,162	12,004	13,230	14,148	15,123	16,211	17,144
Unknown cause	5,327	5,621	5,509	5,850	6,016	5,904	6,023	6,263	6,318	6,037	6,069	6,299	6,412	6,742
Missing cause	11,492	10,150	8,047	7,925	6,294	4,914	3,744	2,960	2,473	1,814	1,518	1,221	826	755
All	49,814	56,082	62,798	72,657	79,990	86,430	92,986	101,022	110,571	122,879	135,827	150,956	166,565	180,893



Table D.6 (continued)

Counts of point prevalent ESRD patients, by mode of therapy: all dialysis

December 31 point prevalent patients, by age, gender, race, ethnicity, & primary diagnosis

	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005
0-4	214	226	225	195	207	206	231	221	222	228	258	274
5-9	216	205	210	219	209	213	239	230	236	214	231	212
10-14	434	433	428	430	455	424	485	511	556	517	522	475
15-19	1,059	1,086	1,143	1,153	1,185	1,175	1,128	1,176	1,226	1,252	1,289	1,297
20-29	7,945	8,107	8,333	8,503	8,786	8,856	8,954	8,960	8,858	8,833	8,896	9,029
30-39	18,699	19,126	19,649	20,138	20,533	21,206	21,650	21,888	22,165	22,461	22,483	22,908
40-49	28,110	30,576	32,838	34,592	36,653	38,316	39,842	41,069	42,201	43,260	44,393	45,169
50-59	35,418	37,722	40,753	44,234	48,107	52,073	55,931	59,695	62,590	65,804	69,203	73,239
60-64	22,220	23,561	24,877	26,433	28,218	29,633	31,065	32,541	34,403	36,321	37,915	39,364
65-69	26,169	27,138	28,529	29,903	31,265	32,552	33,736	35,158	36,074	37,165	38,302	39,180
70-79	42,896	45,877	49,584	53,991	57,733	60,764	63,552	65,713	67,816	69,194	70,542	71,557
80+	14,002	15,422	17,159	19,288	21,519	23,692	26,515	29,049	31,761	34,240	36,499	38,615
Unknown	*	*	*	*	*	*	*	.
0-19	1,923	1,950	2,006	1,997	2,056	2,018	2,083	2,138	2,240	2,211	2,300	2,258
20-44	39,506	41,037	42,674	44,191	45,665	46,982	47,950	48,666	49,018	49,623	50,212	51,025
45-64	72,886	78,055	83,776	89,709	96,632	103,102	109,492	115,487	121,199	127,056	132,678	138,684
65-74	51,398	53,667	56,638	60,062	62,965	65,349	67,728	70,143	71,873	73,512	75,320	76,596
75+	31,669	34,770	38,634	43,120	47,552	51,659	56,075	59,777	63,778	67,087	70,023	72,756
Unknown	*	*	*	*	*	*	*	.
Male	102,244	108,386	116,146	124,815	133,566	141,841	149,971	157,568	164,972	171,572	178,622	185,402
Female	95,139	101,090	107,579	114,261	121,303	127,268	133,355	138,641	143,134	147,914	151,905	155,900
Unknown	.	*	*	*	*	*	*	*	*	*	*	17
White	112,775	118,017	124,951	133,056	140,982	148,867	156,976	164,310	170,284	176,069	182,570	188,885
African American	74,792	79,978	85,648	91,473	97,801	102,776	107,362	111,489	115,980	120,152	123,251	126,937
Native American	2,720	2,977	3,220	3,438	3,769	4,023	4,252	4,483	4,607	4,719	4,897	5,053
Asian	6,531	7,296	8,032	8,736	9,556	10,412	11,300	12,120	12,974	13,684	14,378	15,038
Other/unknown	565	1,211	1,877	2,376	2,762	3,033	3,439	3,810	4,264	4,866	5,438	5,406
†Hispanic			24,142	27,132	29,862	32,902	36,944	39,543	42,286	45,179	48,018	50,494
†Non-Hispanic			199,586	211,947	225,008	236,209	246,385	256,669	265,823	274,311	282,516	290,825
Diabetes	66,908	74,116	82,282	91,258	99,678	106,955	115,105	122,435	128,719	134,832	140,931	146,300
Hypertension	61,606	63,208	65,803	69,081	72,725	76,415	79,823	83,212	86,697	90,629	93,608	96,027
Glomerulonephritis	30,816	32,167	33,665	35,000	36,254	37,203	37,739	38,292	38,706	38,793	39,087	39,081
Cystic kidney	7,388	7,579	7,714	7,794	8,053	8,333	8,551	8,655	8,740	8,784	8,943	9,186
Other urologic	4,959	5,514	5,992	6,487	6,894	7,480	7,822	8,104	8,385	8,508	8,561	8,130
Other cause	18,146	19,082	19,906	20,727	21,970	22,899	23,875	24,596	25,435	26,101	26,947	28,874
Unknown cause	7,103	7,400	7,993	8,402	8,984	9,533	10,144	10,656	11,080	11,397	11,690	12,230
Missing cause	457	413	373	330	312	293	270	262	347	446	767	1,491
All	197,383	209,479	223,728	239,079	254,870	269,111	283,329	296,212	308,109	319,490	330,534	341,319

Table D.7

Counts of point prevalent ESRD patients, by mode of therapy: hemodialysis*December 31 point prevalent patients, by age, gender, race, ethnicity, & primary diagnosis*

	1980	1981	1982	1983	1984	1985	1986	1987	1988	1989	1990	1991	1992	1993
0-4	25	21	23	32	34	41	44	43	41	45	44	45	32	44
5-9	83	69	74	66	57	38	50	56	60	51	57	53	58	63
10-14	225	236	228	220	177	164	139	151	158	146	169	157	171	175
15-19	789	782	760	683	646	611	587	603	652	683	657	643	676	688
20-29	3,783	3,958	4,164	4,299	4,373	4,418	4,399	4,461	4,711	5,077	5,278	5,390	5,685	5,851
30-39	5,899	6,477	6,983	7,470	7,915	8,254	8,702	9,149	9,758	10,650	11,199	11,995	12,872	13,595
40-49	7,258	7,821	8,275	9,261	9,797	10,046	10,495	11,294	12,483	14,041	15,491	17,157	18,822	20,594
50-59	10,933	12,061	13,140	13,997	14,585	15,114	15,665	16,284	17,570	18,936	20,385	22,320	24,554	26,825
60-64	5,732	6,649	7,474	8,615	9,467	10,088	10,537	11,426	12,201	13,224	14,310	15,463	16,457	17,700
65-69	5,165	5,758	6,381	7,804	8,605	9,372	10,534	11,957	13,206	14,738	16,226	17,765	19,539	20,981
70-79	5,487	6,428	7,456	9,649	11,119	12,406	14,184	16,015	18,023	20,794	23,587	27,345	30,865	33,902
80+	786	988	1,234	1,827	2,183	2,696	3,302	4,042	4,773	5,750	6,980	8,243	9,844	11,307
Unknown	*	*	*	*	*	*	11	*	*	*	*	*	.	.
0-19	1,122	1,108	1,085	1,001	914	854	820	853	911	925	927	898	937	970
20-44	13,071	14,002	14,960	16,110	16,810	17,333	18,078	18,890	20,385	22,416	24,068	25,622	27,509	29,057
45-64	20,534	22,964	25,076	27,532	29,327	30,587	31,720	33,724	36,338	39,512	42,595	46,703	50,881	55,508
65-74	8,796	9,932	11,195	13,859	15,424	16,885	19,035	21,395	23,704	26,789	30,071	33,821	37,666	40,874
75+	2,642	3,242	3,876	5,421	6,483	7,589	8,985	10,619	12,298	14,493	16,722	19,532	22,582	25,316
Unknown	*	*	*	*	*	*	11	*	*	*	*	*	.	.
Male	25,203	27,629	30,157	33,965	36,628	38,551	41,019	44,187	48,233	53,579	58,939	65,218	71,862	78,408
Female	20,964	23,622	26,040	29,964	32,338	34,706	37,630	41,299	45,407	50,562	55,449	61,361	67,713	73,317
Unknown	.	.	.	*	*
White	30,670	33,429	36,181	40,398	42,762	44,623	47,351	50,762	54,709	60,189	65,339	71,591	78,062	83,750
African American	14,260	16,357	18,411	21,691	24,156	26,275	28,624	31,695	35,361	39,952	44,423	49,606	55,319	60,763
Native American	270	357	430	563	630	728	837	958	1,135	1,290	1,444	1,628	1,871	2,115
Asian	309	422	611	810	1,009	1,285	1,518	1,795	2,165	2,525	2,994	3,541	4,059	4,790
Other/unknown	658	686	564	468	409	346	319	276	271	185	188	213	264	307
†Hispanic	1,114	1,399	1,638	1,903	2,315	2,740	3,290	3,990	5,051	6,493	7,502	8,714	10,562	12,603
†Non-Hispanic	45,053	49,852	54,559	62,027	66,651	70,517	75,359	81,496	88,590	97,648	106,886	117,865	129,013	139,122
Diabetes	4,294	5,909	7,740	9,748	11,988	14,096	16,582	19,474	22,894	27,603	32,601	38,265	44,335	50,135
Hypertension	7,954	10,100	12,573	15,240	17,525	19,512	21,870	24,635	27,772	31,892	35,568	39,911	44,891	48,977
Glomerulonephritis	9,353	10,487	11,675	13,094	14,001	14,762	15,567	16,433	17,364	18,619	19,621	20,693	21,787	22,977
Cystic kidney	3,021	3,346	3,721	4,154	4,360	4,522	4,590	4,744	4,862	4,981	5,057	5,259	5,362	5,553
Other urologic	1,118	1,385	1,709	2,066	2,361	2,652	2,832	3,134	3,381	3,560	3,623	3,866	4,050	4,022
Other cause	4,628	5,291	6,212	7,146	7,808	8,341	8,836	9,217	9,931	10,854	11,571	12,245	13,032	13,726
Unknown cause	4,887	5,075	4,893	5,120	5,183	5,021	5,096	5,329	5,380	5,136	5,123	5,366	5,473	5,732
Missing cause	10,912	9,658	7,674	7,362	5,740	4,351	3,276	2,520	2,057	1,496	1,224	974	645	603
All	46,167	51,251	56,197	63,930	68,966	73,257	78,649	85,486	93,641	104,141	114,388	126,579	139,575	151,725



Table D.7 (continued)

Counts of point prevalent ESRD patients, by mode of therapy: hemodialysis

December 31 point prevalent patients, by age, gender, race, ethnicity, & primary diagnosis

	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005
0-4	49	56	49	44	45	48	56	54	47	41	54	55
5-9	63	64	75	81	85	84	102	91	92	72	89	80
10-14	200	197	189	182	210	205	233	256	278	259	248	227
15-19	658	711	731	761	798	808	780	809	877	902	916	930
20-29	6,134	6,317	6,606	6,830	7,192	7,368	7,485	7,520	7,506	7,503	7,587	7,763
30-39	14,450	15,084	15,796	16,579	17,207	17,911	18,527	18,825	19,164	19,528	19,628	19,999
40-49	22,441	24,660	27,123	29,106	31,530	33,350	34,980	36,389	37,594	38,633	39,762	40,567
50-59	29,383	31,526	34,520	38,131	42,220	46,166	50,094	53,793	56,644	59,723	63,093	66,842
60-64	18,947	20,186	21,680	23,403	25,421	26,950	28,438	29,936	31,709	33,458	35,058	36,422
65-69	22,551	23,572	25,151	26,824	28,467	29,847	31,109	32,513	33,445	34,555	35,690	36,575
70-79	38,039	41,134	45,001	49,578	53,566	56,817	59,736	61,898	63,930	65,259	66,589	67,474
80+	12,896	14,325	16,074	18,219	20,529	22,695	25,502	27,943	30,591	33,016	35,248	37,228
Unknown	*	*	*	.
0-19	970	1,028	1,044	1,068	1,138	1,145	1,171	1,210	1,294	1,274	1,307	1,292
20-44	30,753	32,416	34,448	36,386	38,424	39,895	41,150	42,022	42,651	43,294	43,942	44,765
45-64	60,602	65,357	71,277	77,663	85,146	91,850	98,374	104,441	109,966	115,551	121,186	126,828
65-74	44,686	47,140	50,431	54,330	57,636	60,258	62,858	65,297	67,013	68,571	70,419	71,672
75+	28,800	31,891	35,795	40,291	44,926	49,101	53,489	57,057	60,953	64,259	67,108	69,605
Unknown	*	*	*	.
Male	85,718	91,911	100,348	109,789	119,477	128,157	136,469	144,231	151,506	157,958	164,850	171,245
Female	80,093	85,919	92,645	99,947	107,792	114,090	120,572	125,795	130,371	134,988	139,109	142,911
Unknown	.	*	*	*	*	*	*	*	*	*	*	*
White	91,144	96,537	104,260	113,367	122,636	131,020	139,612	147,036	153,037	158,689	165,263	171,385
African American	66,408	71,627	77,548	83,787	90,484	95,692	100,448	104,645	109,146	113,216	116,339	119,883
Native American	2,331	2,580	2,845	3,060	3,412	3,700	3,903	4,127	4,247	4,359	4,570	4,725
Asian	5,463	6,070	6,757	7,479	8,296	9,137	9,977	10,768	11,567	12,218	12,873	13,455
Other/unknown	465	1,018	1,585	2,045	2,442	2,700	3,102	3,451	3,881	4,468	4,918	4,714
†Hispanic	14,622	17,871	21,082	24,046	26,837	29,734	33,680	36,293	38,863	41,704	44,601	47,076
†Non-Hispanic	151,189	159,961	171,913	185,692	200,433	212,515	223,362	233,734	243,015	251,246	259,362	267,086
Diabetes	56,840	63,768	71,991	81,336	90,365	97,885	106,173	113,453	119,699	125,666	131,985	137,289
Hypertension	53,538	55,474	58,657	62,424	66,525	70,305	73,833	77,388	80,690	84,471	87,372	89,580
Glomerulonephritis	24,386	25,574	27,155	28,728	30,353	31,613	32,306	33,012	33,512	33,654	33,939	34,051
Cystic kidney	5,860	6,037	6,205	6,409	6,778	7,080	7,337	7,434	7,592	7,549	7,711	7,899
Other urologic	4,198	4,725	5,198	5,693	6,121	6,701	7,033	7,305	7,560	7,694	7,737	7,391
Other cause	14,617	15,596	16,579	17,562	18,944	19,902	21,010	21,587	22,425	23,105	23,924	25,723
Unknown cause	6,014	6,330	6,915	7,334	7,942	8,538	9,162	9,659	10,126	10,450	10,728	11,222
Missing cause	358	328	295	252	242	225	188	189	274	361	567	1,007
All	165,811	177,832	192,995	209,738	227,270	242,249	257,042	270,027	281,878	292,950	303,963	314,162

Table D.8

Counts of point prevalent ESRD patients, by mode of therapy: peritoneal dialysis*December 31 point prevalent patients, by age, gender, race, ethnicity, & primary diagnosis*

	1980	1981	1982	1983	1984	1985	1986	1987	1988	1989	1990	1991	1992	1993
0-4	11	32	54	67	76	104	116	98	99	105	108	133	143	136
5-9	*	20	45	72	86	72	91	89	88	111	113	102	96	104
10-14	17	33	64	119	144	137	158	163	170	162	191	212	203	204
15-19	11	38	90	125	158	229	215	236	243	291	301	328	346	355
20-29	65	194	301	527	681	771	838	922	1,037	1,084	1,255	1,372	1,484	1,613
30-39	145	376	617	1,006	1,325	1,642	1,718	1,878	2,085	2,331	2,723	3,065	3,362	3,676
40-49	137	387	660	1,158	1,435	1,761	1,929	2,075	2,353	2,744	3,233	3,886	4,336	4,788
50-59	228	652	955	1,485	1,939	2,266	2,462	2,617	2,716	3,082	3,543	3,987	4,546	4,998
60-64	78	337	546	863	1,126	1,429	1,558	1,722	1,811	1,970	2,239	2,339	2,689	2,796
65-69	88	252	429	601	885	1,190	1,429	1,552	1,693	1,903	2,168	2,586	2,864	3,192
70-79	92	244	516	633	916	1,269	1,481	1,711	1,860	2,115	2,478	3,083	3,586	4,064
80+	13	48	93	110	153	204	251	305	349	425	505	618	711	828
Unknown	.	.	.	*	*	*	*	*
0-19	48	123	253	383	464	542	580	586	600	669	713	775	788	799
20-44	273	758	1,224	2,075	2,690	3,238	3,521	3,850	4,292	4,753	5,591	6,382	6,974	7,600
45-64	380	1,188	1,855	2,964	3,816	4,631	4,984	5,364	5,710	6,458	7,402	8,267	9,443	10,271
65-74	147	414	781	1,010	1,491	2,033	2,379	2,621	2,900	3,238	3,729	4,526	5,125	5,813
75+	46	130	257	334	463	630	782	947	1,002	1,205	1,422	1,761	2,036	2,271
Unknown	.	.	.	*	*	*	*	*
Male	467	1,345	2,285	3,687	4,775	5,904	6,427	7,023	7,592	8,587	9,871	11,486	12,779	14,026
Female	427	1,268	2,085	3,080	4,151	5,172	5,820	6,347	6,912	7,736	8,986	10,225	11,587	12,728
Unknown
White	732	2,058	3,362	5,229	6,785	8,437	9,214	9,967	10,752	11,866	13,578	15,458	17,224	18,571
African American	137	502	917	1,372	1,893	2,359	2,700	2,996	3,324	3,931	4,667	5,509	6,180	7,009
Native American	14	20	38	66	115	121	152	185	180	197	234	266	327	338
Asian	*	20	42	68	111	134	156	203	227	312	361	455	606	800
Other/unknown	*	13	11	32	22	25	25	19	21	17	17	23	29	36
†Hispanic														
†Non-Hispanic														
Diabetes	203	547	1,020	1,418	2,010	2,525	2,916	3,316	3,728	4,469	5,353	6,454	7,612	8,315
Hypertension	169	496	925	1,299	1,717	2,226	2,630	2,903	3,249	3,760	4,597	5,421	6,106	6,820
Glomerulonephritis	223	602	946	1,621	2,195	2,705	2,909	3,193	3,479	3,792	4,198	4,680	5,087	5,541
Cystic kidney	63	178	301	451	581	715	799	829	826	882	1,010	1,128	1,244	1,352
Other urologic	26	89	141	203	288	361	372	384	421	474	515	544	608	661
Other cause	106	305	547	888	1,166	1,483	1,592	1,718	1,840	2,073	2,307	2,598	2,832	3,122
Unknown cause	77	220	320	458	548	605	651	678	664	666	719	757	791	870
Missing cause	27	176	170	429	421	456	378	349	297	207	158	129	86	73
All	894	2,613	4,370	6,767	8,926	11,076	12,247	13,370	14,504	16,323	18,857	21,711	24,366	26,754



III



Table D.8 (continued)

Counts of point prevalent ESRD patients, by mode of therapy: peritoneal dialysis

December 31 point prevalent patients, by age, gender, race, ethnicity, & primary diagnosis

	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005
0-4	150	156	165	143	158	152	168	158	171	182	197	213
5-9	134	128	127	129	109	117	131	134	136	136	140	128
10-14	210	206	216	230	225	203	231	235	266	241	258	228
15-19	351	344	372	354	335	329	316	318	307	316	341	323
20-29	1,663	1,651	1,614	1,569	1,468	1,356	1,341	1,331	1,245	1,246	1,226	1,177
30-39	3,878	3,814	3,650	3,369	3,133	3,080	2,919	2,883	2,838	2,779	2,698	2,754
40-49	5,211	5,564	5,466	5,250	4,885	4,718	4,599	4,445	4,383	4,433	4,415	4,364
50-59	5,610	5,938	6,022	5,902	5,670	5,668	5,609	5,721	5,746	5,890	5,868	6,059
60-64	3,023	3,226	3,109	2,966	2,725	2,604	2,561	2,526	2,612	2,781	2,761	2,803
65-69	3,342	3,429	3,295	3,020	2,739	2,656	2,571	2,593	2,588	2,563	2,547	2,517
70-79	4,456	4,551	4,448	4,311	4,077	3,872	3,728	3,756	3,834	3,879	3,884	3,989
80+	968	1,021	1,022	1,035	951	964	968	1,070	1,129	1,194	1,225	1,340
Unknown	*	*	*
0-19	845	834	880	856	827	801	846	845	880	875	936	892
20-44	8,000	8,078	7,794	7,393	6,804	6,607	6,342	6,231	5,995	6,008	5,924	5,915
45-64	11,385	12,115	12,067	11,663	11,077	10,819	10,687	10,675	10,829	11,121	11,044	11,242
65-74	6,202	6,276	6,048	5,615	5,221	4,999	4,773	4,768	4,793	4,857	4,792	4,785
75+	2,564	2,725	2,717	2,751	2,546	2,493	2,494	2,651	2,758	2,779	2,864	3,061
Unknown	*	*	*
Male	15,119	15,590	15,151	14,456	13,461	13,055	12,856	12,779	12,942	13,118	13,195	13,426
Female	13,877	14,438	14,355	13,822	13,014	12,665	12,287	12,392	12,313	12,522	12,365	12,469
Unknown
White	19,838	20,392	19,869	18,974	17,586	17,068	16,571	16,619	16,606	16,772	16,704	16,861
African American	7,716	7,907	7,773	7,417	7,031	6,813	6,645	6,562	6,582	6,714	6,659	6,773
Native American	376	390	369	370	347	310	336	344	353	349	321	317
Asian	1,006	1,184	1,237	1,220	1,216	1,224	1,281	1,314	1,360	1,435	1,464	1,540
Other/unknown	60	155	258	297	295	305	310	332	354	370	412	404
†Hispanic			2,949	2,975	2,904	3,035	3,153	3,128	3,290	3,352	3,288	3,289
†Non-Hispanic			26,557	25,303	23,571	22,685	21,990	22,043	21,965	22,288	22,272	22,606
Diabetes	9,295	9,950	10,042	9,717	9,107	8,855	8,713	8,790	8,845	8,968	8,748	8,806
Hypertension	7,436	7,384	6,895	6,465	5,986	5,912	5,778	5,642	5,808	6,003	6,084	6,301
Glomerulonephritis	5,943	6,206	6,200	5,963	5,568	5,241	5,116	4,980	4,903	4,908	4,900	4,797
Cystic kidney	1,417	1,466	1,440	1,331	1,229	1,199	1,146	1,175	1,101	1,177	1,173	1,232
Other urologic	690	739	748	752	725	728	733	757	779	775	782	704
Other cause	3,225	3,283	3,160	3,020	2,860	2,834	2,718	2,859	2,882	2,872	2,881	3,000
Unknown cause	943	964	992	1,003	975	924	914	950	914	903	933	976
Missing cause	47	36	29	27	25	27	25	18	23	34	59	79
All	28,996	30,028	29,506	28,278	26,475	25,720	25,143	25,171	25,255	25,640	25,560	25,895



Table D.9

Counts of point prevalent ESRD patients, by mode of therapy: functioning transplant*December 31 point prevalent patients, by age, gender, race, ethnicity, & primary diagnosis*

	1980	1981	1982	1983	1984	1985	1986	1987	1988	1989	1990	1991	1992	1993
0-4	31	47	74	88	99	125	135	173	179	176	166	177	192	200
5-9	115	129	157	173	222	291	341	376	405	464	509	565	567	596
10-14	306	339	386	430	490	533	561	616	665	725	774	819	886	934
15-19	704	794	845	937	1,007	1,098	1,224	1,321	1,341	1,343	1,366	1,389	1,409	1,488
20-29	2,680	3,235	3,771	4,271	4,756	5,291	5,885	6,243	6,473	6,640	6,970	7,227	7,324	7,411
30-39	2,857	3,760	4,648	5,687	6,917	8,262	9,669	10,767	11,800	12,752	13,819	14,826	15,674	16,514
40-49	1,974	2,486	3,144	3,879	4,733	5,890	7,275	8,751	10,197	11,644	13,035	14,803	16,497	18,286
50-59	1,195	1,542	1,968	2,448	3,126	3,848	4,922	5,961	6,943	7,931	9,055	10,129	11,512	12,945
60-64	171	235	316	426	600	834	1,173	1,578	2,071	2,530	3,005	3,512	3,969	4,415
65-69	59	73	95	122	169	259	380	590	827	1,087	1,394	1,757	2,203	2,712
70-79	16	23	27	37	52	78	114	178	245	328	471	678	904	1,135
80+	*	*	*	*	*	12	17	16	17	18	26	32	32	37
Unknown	*	*	*	*	*	*	*	*	*	*	*	*	*	*
0-19	1,156	1,309	1,462	1,628	1,818	2,047	2,261	2,486	2,590	2,708	2,815	2,950	3,054	3,218
20-44	6,603	8,361	10,159	12,087	14,271	16,747	19,556	21,780	23,786	25,746	28,008	30,158	31,864	33,585
45-64	2,274	2,897	3,688	4,624	5,861	7,378	9,368	11,520	13,698	15,751	17,876	20,339	23,112	25,986
65-74	72	88	115	152	213	323	473	730	1,020	1,346	1,774	2,319	2,970	3,669
75+	*	*	11	*	13	26	38	54	69	87	117	148	169	215
Unknown	*	*	*	*	*	*	*	*	*	*	*	*	*	*
Male	6,044	7,668	9,415	11,302	13,565	16,277	19,444	22,396	25,044	27,636	30,505	33,555	36,617	39,813
Female	4,071	5,000	6,024	7,202	8,614	10,246	12,254	14,175	16,119	18,001	20,084	22,359	24,552	26,861
Unknown	.	*	.	.	*	*	*	*	*	*	*	*	*	*
White	8,201	10,292	12,503	14,908	17,827	21,211	25,286	29,122	32,649	36,269	40,168	44,357	48,406	52,681
African American	1,543	1,938	2,413	2,950	3,547	4,352	5,232	6,014	6,762	7,369	8,112	8,964	9,838	10,756
Native American	72	98	120	144	182	222	255	299	362	430	479	544	595	653
Asian	98	131	190	263	365	457	618	786	985	1,157	1,387	1,591	1,866	2,107
Other/unknown	201	210	213	239	259	282	308	351	406	414	445	459	465	478
†Hispanic														
†Non-Hispanic														
Diabetes	854	1,239	1,721	2,290	2,957	3,881	5,023	6,047	7,025	8,123	9,293	10,589	11,745	13,290
Hypertension	1,046	1,355	1,740	2,224	2,706	3,378	4,109	4,830	5,546	6,203	7,018	7,777	8,676	9,544
Glomerulonephritis	3,446	4,304	5,265	6,273	7,559	8,922	10,458	11,957	13,362	14,652	16,129	17,717	19,213	20,602
Cystic kidney	435	587	760	955	1,235	1,539	1,978	2,464	2,937	3,370	3,854	4,344	4,910	5,399
Other urologic	419	547	653	807	961	1,143	1,351	1,525	1,680	1,868	2,045	2,250	2,415	2,643
Other cause	1,664	2,088	2,524	3,032	3,647	4,322	5,150	5,823	6,316	6,871	7,469	8,226	8,966	9,786
Unknown cause	819	1,010	1,171	1,308	1,468	1,656	1,859	2,066	2,259	2,448	2,594	2,751	2,909	3,064
Missing cause	1,432	1,539	1,605	1,615	1,647	1,683	1,771	1,860	2,039	2,104	2,189	2,261	2,336	2,347
All	10,115	12,669	15,439	18,504	22,180	26,524	31,699	36,572	41,164	45,639	50,591	55,915	61,170	66,675



113



Table D.9 (continued)

Counts of point prevalent ESRD patients, by mode of therapy: functioning transplant

December 31 point prevalent patients, by age, gender, race, ethnicity, & primary diagnosis

	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005
0-4	196	184	211	208	212	235	247	264	266	256	278	287
5-9	590	612	627	691	679	696	657	697	708	758	773	808
10-14	1,016	1,099	1,167	1,160	1,238	1,291	1,311	1,311	1,365	1,374	1,361	1,398
15-19	1,651	1,782	1,817	1,909	1,939	2,085	2,176	2,211	2,236	2,380	2,486	2,611
20-29	7,402	7,488	7,634	7,726	7,775	7,788	7,792	7,805	7,873	7,971	8,176	8,367
30-39	17,299	18,036	18,615	18,927	19,430	19,617	19,923	20,100	20,202	20,077	20,027	19,909
40-49	20,039	21,890	23,461	24,787	26,275	27,650	29,161	30,388	31,481	32,542	33,641	34,402
50-59	14,383	15,843	17,610	19,880	22,170	24,432	26,690	29,202	31,641	33,988	36,743	39,602
60-64	4,904	5,509	6,235	6,915	7,719	8,532	9,323	10,275	11,480	12,734	13,928	15,137
65-69	3,176	3,661	4,206	4,721	5,259	5,896	6,548	7,308	8,117	9,035	9,948	11,025
70-79	1,449	1,797	2,190	2,701	3,254	3,786	4,517	5,225	6,175	7,094	8,125	9,422
80+	49	58	85	96	129	178	232	296	378	466	591	725
Unknown	*	*
0-19	3,453	3,677	3,822	3,968	4,068	4,307	4,391	4,483	4,575	4,768	4,898	5,104
20-44	35,027	36,638	37,926	38,942	40,156	41,032	41,807	42,458	43,019	43,320	43,830	44,183
45-64	29,000	32,128	35,629	39,293	43,213	46,987	51,082	55,312	59,658	63,992	68,685	73,234
65-74	4,391	5,123	5,966	6,862	7,799	8,807	10,009	11,273	12,733	14,258	15,887	17,757
75+	283	393	515	656	843	1,053	1,288	1,556	1,937	2,337	2,777	3,415
Unknown	*	*
Male	43,032	46,522	49,937	53,444	57,066	60,652	64,421	68,224	72,274	76,271	80,556	85,122
Female	29,122	31,435	33,918	36,274	39,010	41,530	44,151	46,853	49,643	52,399	55,516	58,566
Unknown	*	*	*	*	*	*	*	*	*	*	*	*
White	56,573	60,731	64,994	69,247	73,945	78,401	83,067	87,735	92,739	97,355	102,147	106,917
African American	11,953	13,238	14,503	15,714	16,923	18,092	19,302	20,618	21,959	23,403	24,960	26,665
Native American	731	796	851	911	970	1,065	1,131	1,190	1,251	1,302	1,374	1,427
Asian	2,418	2,706	3,010	3,325	3,683	4,029	4,412	4,809	5,193	5,680	6,200	6,751
Other/unknown	480	489	500	524	558	599	665	730	780	935	1,396	1,933
†Hispanic			7,377	8,122	9,101	10,028	11,032	12,136	13,390	14,615	15,924	17,287
†Non-Hispanic			76,481	81,599	86,978	92,158	97,545	102,946	108,532	114,060	120,153	126,406
Diabetes	14,703	16,212	17,826	19,320	20,957	22,485	24,072	25,727	27,543	29,188	30,951	32,857
Hypertension	10,534	11,517	12,491	13,367	14,269	15,050	15,909	16,917	17,932	18,855	20,059	21,411
Glomerulonephritis	22,002	23,494	24,951	26,434	28,085	29,681	31,291	32,781	34,418	36,001	37,728	39,264
Cystic kidney	5,869	6,450	6,999	7,646	8,303	8,901	9,533	10,225	10,949	11,777	12,490	13,272
Other urologic	2,805	3,014	3,227	3,436	3,672	3,893	4,140	4,405	4,693	4,978	5,243	5,451
Other cause	10,595	11,425	12,267	13,083	13,940	14,805	15,693	16,571	17,466	18,340	19,390	20,377
Unknown cause	3,220	3,343	3,520	3,770	4,015	4,314	4,626	5,018	5,398	5,813	6,162	6,597
Missing cause	2,427	2,505	2,577	2,665	2,838	3,057	3,313	3,438	3,523	3,723	4,054	4,464
All	72,155	77,960	83,858	89,721	96,079	102,186	108,577	115,082	121,922	128,675	136,077	143,693



Table D.10

Incident ESRD patients, 2005*by age, gender, race, ethnicity, primary diagnosis, & modality*

	Center hemo	Center self hemo	Home hemo	CAPD	CCPD	Other PD	Uncertain dialysis	Transplant	All
0-4	45	.	.	*	127	*	*	31	218
5-9	50	.	.	*	52	.	*	37	148
10-14	142	.	.	12	102	*	*	75	337
15-19	428	.	.	28	64	.	*	92	622
20-29	2,305	.	*	168	78	.	18	148	2,724
30-39	4,921	*	20	392	190	*	29	322	5,878
40-49	10,498	*	50	658	280	*	63	529	12,081
50-59	18,093	*	79	1,107	493	*	121	674	20,576
60-64	10,754	*	43	551	222	*	63	248	11,892
65-69	11,248	*	42	518	167	*	39	174	12,196
70-79	23,615	*	98	859	289	16	45	88	25,017
80+	14,583	*	89	364	138	*	23	*	15,221
Unknown	*	*
0-19	665	.	.	56	345	*	22	235	1,325
20-44	11,617	*	53	860	400	*	75	694	13,706
45-64	34,954	*	146	2,016	863	11	219	1,227	39,445
65-74	22,981	*	78	996	331	15	65	239	24,711
75+	26,465	14	151	745	263	14	42	29	27,723
Unknown	*	*
Male	53,733	20	242	2,536	1,183	24	234	1,381	59,353
Female	42,945	11	186	2,137	1,019	23	181	1,042	47,544
Unknown	*	*	*	15
White	62,469	22	266	3,366	1,575	36	162	1,851	69,747
African American	28,394	*	140	946	486	*	44	202	30,228
Native American	1,115	.	*	52	23	.	*	*	1,202
Asian	3,613	*	11	269	70	*	*	65	4,036
Other/unknown	1,093	.	*	40	48	.	209	299	1,699
Hispanic	12,774	*	35	441	315	*	*	126	13,709
Non-Hispanic	83,910	29	393	4,232	1,887	39	415	2,298	93,203
Diabetes	43,421	14	154	2,022	754	18	26	442	46,851
Hypertension	26,712	*	135	1,074	450	11	12	218	28,622
Glomerulonephritis	6,696	*	18	634	343	.	18	390	8,100
Cystic kidney	1,730	.	*	265	143	*	12	338	2,495
Other urologic	1,939	*	*	75	50	*	.	81	2,158
Other cause	11,013	*	48	385	370	*	29	374	12,224
Unknown cause	4,168	*	35	179	79	*	*	119	4,590
Missing cause	1,005	.	22	39	13	11	320	462	1,872
All	96,684	31	428	4,673	2,202	47	423	2,424	106,912

on our website & CD

Incident ESRD patients for selected years: by age, gender, race, ethnicity, primary diagnosis, & modality (Table D.10.1).



115



Table D.11

Point prevalent ESRD patients, 2005

December 31 point prevalent patients, by age, gender, race, ethnicity, primary diagnosis, & modality

	Center hemo	Center self hemo	Home hemo	CAPD	CCPD	Other PD	Uncertain dialysis	Transplant	All
0-4	53	.	*	13	200	.	*	287	561
5-9	77	.	*	11	117	.	*	808	1,020
10-14	223	.	*	16	212	*	18	1,398	1,873
15-19	919	.	11	44	279	*	43	2,611	3,908
20-29	7,701	*	55	423	754	.	89	8,367	17,396
30-39	19,779	16	204	1,176	1,578	*	153	19,909	42,817
40-49	40,202	24	341	1,910	2,454	*	236	34,402	79,571
50-59	66,288	33	521	2,729	3,330	*	331	39,602	112,841
60-64	36,184	26	212	1,276	1,527	*	133	15,137	54,501
65-69	36,392	*	173	1,011	1,506	*	78	11,025	50,205
70-79	67,089	14	371	1,587	2,402	*	90	9,422	80,979
80+	37,012	*	208	536	804	*	44	725	39,340
Unknown
0-19	1,272	.	20	84	808	*	71	5,104	7,362
20-44	44,328	30	407	2,451	3,464	*	342	44,183	95,208
45-64	125,826	76	926	5,063	6,179	14	600	73,234	211,918
65-74	71,266	18	388	1,912	2,873	13	126	17,757	94,353
75+	69,227	14	364	1,222	1,839	*	86	3,415	76,171
Unknown
Male	169,942	85	1,218	5,416	8,010	20	711	85,122	270,524
Female	141,972	53	886	5,316	7,153	17	503	58,566	214,466
Unknown	*	.	*	.	.	.	11	*	22
White	169,925	66	1,394	6,893	9,968	19	620	106,917	295,802
African American	119,198	62	623	2,779	3,994	*	272	26,665	153,602
Native American	4,715	*	*	155	162	*	*	1,427	6,480
Asian	13,385	*	63	759	781	*	42	6,751	21,789
Other/unknown	4,696	*	17	146	258	*	281	1,933	7,339
Hispanic	46,933	12	131	1,237	2,052	11	118	17,287	67,781
Non-Hispanic	264,986	126	1,974	9,495	13,111	26	1,107	126,406	417,231
Diabetes	136,576	40	673	3,903	4,903	12	193	32,857	179,157
Hypertension	89,037	44	499	2,701	3,600	*	145	21,411	117,438
Glomerulonephritis	33,633	24	394	1,869	2,928	*	232	39,264	78,345
Cystic kidney	7,774	*	116	524	708	*	54	13,272	22,458
Other urologic	7,304	*	83	253	451	*	32	5,451	13,581
Other cause	25,494	14	215	1,039	1,961	*	142	20,377	49,251
Unknown cause	11,125	*	94	404	572	*	30	6,597	18,827
Missing cause	976	.	31	39	40	*	397	4,464	5,955
All	311,919	138	2,105	10,732	15,163	37	1,225	143,693	485,012

on our website & CD

Prevalent ESRD patients for selected years (Table D.11.1). Counts of incident ESRD patients for selected years (Table D.a). Living ESRD patients on December 31 for selected years (D.b). Incident ESRD patients for selected years (D.c). Living ESRD patients on December 31 for selected years, by age & ESRD network of residence (D.d).

Table D.12

Treatment status two years after first ESRD service, 2001–2003 combined*incident patients with Medicare coverage*

All	% on Day 90	Count	Treatment status two years after first ESRD service (percent of patients)							
			Center hemo	Home hemo	CAPD & CCPD	Oth/unk dialysis	Transplant	Death	Lost to followup	Recover function
Age at first service										
0-19		1,521	29.3	0.5	16.4	0.4	46.5	5.7	0.4	0.8
20-44		19,349	59.9	0.3	8.5	0.1	13.6	15.1	0.6	1.9
45-64		53,037	60.8	0.4	5.6	0.0	5.3	26.0	0.4	1.5
65-74		51,353	52.0	0.3	4.3	0.0	2.4	38.5	0.5	2.1
75+		50,806	42.5	0.2	2.1	0.0	0.2	53.0	0.4	1.7
Modality on day 90										
Center hemodialysis		155,823	57.5	0.2	1.5	0.0	2.8	36.9	0.4	0.7
Home hemodialysis		667	20.1	9.7	9.4	.	3.0	56.5	0.9	0.3
CAPD/CCPD		13,937	17.9	1.2	41.1	0.1	8.4	30.7	0.2	0.3
Other/unknown dialysis		3,663	9.6	0.1	0.8	0.4	0.7	32.8	0.5	55.3
Transplant		1,976	1.8	.	0.3	0.1	93.0	4.8	0.1	.
Interaction of age at first service & modality on day 90										
0-19	Center hemodialysis	748	51.6	0.5	5.6	0.3	35.0	6.1	0.4	0.4
	Home hemodialysis	*	.	.	30.0	.	40.0	30.0	.	.
	CAPD/CCPD	529	10.4	0.6	37.8	0.8	44.0	5.7	0.6	0.2
	Other/unknown dialysis	20	10.0	.	10.0	.	15.0	25.0	.	40.0
	Transplant	214	1.4	.	0.9	.	96.3	1.4	.	.
20-44	Center hemodialysis	16,251	68.1	0.2	3.7	0.0	10.4	16.2	0.7	0.8
	Home hemodialysis	64	39.1	9.4	20.3	.	9.4	20.3	1.6	.
	CAPD/CCPD	2,122	21.0	1.3	48.7	0.0	18.4	10.3	0.1	0.2
	Other/unknown dialysis	339	11.5	.	1.8	0.6	1.2	13.0	0.3	71.7
	Transplant	573	2.6	.	0.2	0.2	94.1	3.0	.	.
45-64	Center hemodialysis	46,645	66.8	0.2	1.9	0.0	3.6	26.5	0.4	0.6
	Home hemodialysis	156	28.8	14.7	12.8	.	5.8	35.3	1.9	0.6
	CAPD/CCPD	4,544	20.8	1.5	45.0	0.1	8.3	23.8	0.2	0.3
	Other/unknown dialysis	895	11.8	0.1	0.8	0.6	0.9	25.9	1.1	58.8
	Transplant	797	1.8	.	0.1	0.1	93.0	5.0	.	.
65-74	Center hemodialysis	45,459	56.9	0.1	1.2	0.0	1.6	38.9	0.5	0.8
	Home hemodialysis	193	19.7	10.4	8.8	.	0.5	59.1	1.0	0.5
	CAPD/CCPD	4,092	17.4	1.2	39.9	0.0	3.9	36.8	0.3	0.4
	Other/unknown dialysis	1,245	8.9	.	0.6	0.4	0.6	32.8	0.4	56.3
	Transplant	364	1.1	.	0.3	.	89.3	9.1	0.3	.
75+	Center hemodialysis	46,720	45.2	0.1	0.5	.	0.1	53.0	0.4	0.7
	Home hemodialysis	244	10.7	6.6	4.1	.	.	78.7	.	.
	CAPD/CCPD	2,650	13.1	0.6	30.8	.	0.4	54.6	0.3	0.2
	Other/unknown dialysis	1,164	8.0	0.2	0.5	0.2	0.1	44.1	0.1	46.9
	Transplant	28	92.9	7.1	.	.
All		176,066	52.6	0.3	4.6	0.0	4.2	36.1	0.4	1.8



117



on our website & CD

Treatment status two years after first ESRD service, by gender, race, & interactions (Tables D.12.1-18).

Includes only incident patients with known age & patients who have Medicare as their primary payor, are receiving their first ESRD service, & survive the first 90 days.

Table D.13

Point prevalent ESRD patients, 2005: by patient vintage

December 31 point prevalent patients, by year-end treatment modality & number of years patient has had ESRD (vintage)

	Dialysis	Functioning graft	Center hemo	Center self hemo	Home hemo	CAPD	CCPD	Other PD	Uncertain dialysis	All
0	85,255	3,826	77,355	29	402	3,728	3,366	15	360	89,081
1	62,754	6,451	56,621	16	318	2,368	3,211	*	213	69,205
2	47,397	8,131	43,206	16	251	1,533	2,271	*	117	55,528
3	35,074	9,401	32,167	*	204	776	1,872	*	44	44,475
4	26,652	10,186	24,617	11	149	555	1,281	.	39	36,838
5	19,127	10,501	17,883	*	109	340	754	*	34	29,628
6	14,068	10,490	13,158	*	100	258	495	*	46	24,558
7	10,295	10,000	9,612	*	73	189	356	*	55	20,295
8	7,689	8,922	7,177	*	52	157	255	.	43	16,611
9	5,846	8,131	5,434	*	53	130	198	*	25	13,977
10	4,686	7,312	4,326	*	42	111	178	.	27	11,998
11	3,825	6,618	3,509	*	36	95	143	*	36	10,443
12	3,062	5,945	2,808	*	32	72	116	.	32	9,007
13	2,583	5,365	2,349	*	26	71	110	.	25	7,948
14	2,142	4,461	1,933	*	29	63	90	.	24	6,603
15	1,827	3,976	1,655	*	21	51	71	*	25	5,803
16	1,524	3,473	1,379	*	17	44	64	.	18	4,997
17	1,313	3,020	1,182	.	20	30	67	.	14	4,333
18	1,060	2,538	950	*	21	33	44	.	11	3,598
19	859	2,260	773	*	14	23	43	.	*	3,119
20	737	1,957	653	*	16	17	45	*	*	2,694
21	623	1,749	567	.	17	*	25	.	*	2,372
22	536	1,549	483	*	13	17	19	.	*	2,085
23	457	1,439	422	.	11	*	12	.	*	1,896
24	366	1,192	329	.	*	14	13	.	.	1,558
25	312	1,019	289	.	*	*	12	.	*	1,331
26	287	818	263	.	11	*	*	.	*	1,105
27	229	778	192	.	16	*	16	.	*	1,007
28	252	767	217	*	13	*	*	.	*	1,019
29	114	354	104	.	*	*	*	.	*	468
30	114	279	94	.	*	*	*	.	*	393
31	102	253	77	*	11	*	*	.	*	355
32	71	222	61	.	*	*	*	.	*	293
33	28	99	25	.	*	*	*	.	.	127
34	12	79	12	91
35	*	40	*	.	.	*	*	.	.	49
36	32	92	31	.	.	.	*	.	.	124
All	341,319	143,693	311,919	138	2,105	10,732	15,163	37	1,225	485,012



Table D.14

Point prevalent ESRD patients, 2005: by number of years on modality*December 31 point prevalent patients, by year-end treatment modality*

	Dialysis	Functioning graft	Center hemo	Center self hemo	Home hemo	CAPD	CCPD	Other PD	Uncertain dialysis	All
0	221,263	16,390	202,577	107	1,709	6,485	9,380	21	984	237,653
1	62,695	15,126	56,600	23	246	2,327	3,313	*	180	77,821
2	28,501	13,586	25,717	*	76	1,200	1,476	*	28	42,087
3	12,174	12,587	11,296	*	30	295	547	*	*	24,761
4	6,669	11,441	6,233	*	12	175	238	*	*	18,110
5	3,812	10,074	3,606	*	*	80	113	.	*	13,886
6	2,333	8,984	2,209	*	*	60	53	.	*	11,317
7	1,291	8,103	1,239	*	*	21	22	.	*	9,394
8	914	6,817	872	.	*	29	*	*	.	7,731
9	536	5,987	506	.	*	21	*	*	*	6,523
10	467	5,263	450	.	*	*	*	.	*	5,730
11	193	4,350	185	.	.	*	*	.	.	4,543
12	107	3,802	98	.	*	*	*	.	.	3,909
13	85	3,124	81	.	*	*	*	.	.	3,209
14	62	2,875	55	.	*	*	.	.	.	2,937
15	55	2,434	52	.	.	*	.	.	.	2,489
16	26	2,080	24	.	.	*	.	.	.	2,106
17	20	1,747	18	.	.	.	*	*	.	1,767
18	18	1,541	18	1,559
19	20	1,415	19	.	.	*	.	.	.	1,435
20	11	1,082	11	1,093
21	*	952	*	.	*	958
22	13	878	12	.	.	*	.	.	.	891
23	*	695	*	697
24	*	591	*	.	*	597
25	*	468	*	472
26	*	413	*	417
27	*	332	*	337
28	*	237	*	241
29	*	66	*	69
30	*	74	*	*	81
31	*	52	*	*	58
32	*	47	*	*	50
33	.	34	34
34	*	46	*	50
All	341,319	143,693	311,919	138	2,105	10,732	15,163	37	1,225	485,012



Table D.15

Incident ESRD patients, 2005: by modality & state/territory

	Center hemo	Center self hemo	Home hemo	CAPD	CCPD	Other PD	Uncertain dialysis	Transplant	All
Alabama	1,726	.	*	125	24	*	*	21	1,903
Alaska	68	.	.	18	*	.	.	*	91
Arizona	1,755	*	*	112	18	.	*	46	1,937
Arkansas	839	.	*	80	21	*	*	20	964
California	11,225	*	18	585	160	*	50	165	12,210
Colorado	804	.	*	58	23	.	*	36	926
Connecticut	834	.	*	80	66	.	*	21	1,006
Delaware	336	.	.	15	*	.	.	*	359
District of Columbia	424	.	*	11	*	.	*	*	446
Florida	6,006	.	136	209	98	*	14	103	6,568
Georgia	3,243	.	12	222	51	*	*	53	3,586
Hawaii	513	*	*	49	*	.	.	*	571
Idaho	235	*	.	49	*	.	.	*	304
Illinois	4,177	*	85	229	115	*	22	127	4,762
Indiana	1,906	*	*	121	23	.	*	57	2,116
Iowa	728	.	*	40	14	.	*	33	820
Kansas	693	*	.	62	19	.	.	20	797
Kentucky	1,374	.	.	99	19	*	*	27	1,528
Louisiana	2,049	.	*	54	24	*	.	21	2,150
Maine	319	.	.	*	13	.	.	*	347
Maryland	2,308	.	*	83	58	*	*	44	2,510
Massachusetts	1,569	.	*	69	39	*	*	68	1,752
Michigan	3,649	.	16	216	29	.	*	93	4,011
Minnesota	1,119	.	*	58	18	.	*	100	1,302
Mississippi	1,322	.	*	101	42	.	*	16	1,487
Missouri	1,781	.	*	127	38	.	*	38	1,989
Montana	158	.	.	25	*	.	.	*	195
Nebraska	447	*	*	30	13	.	*	18	516
Nevada	664	.	.	36	17	.	*	15	733
New Hampshire	255	.	*	25	*	.	.	11	302
New Jersey	3,271	.	*	67	68	.	*	86	3,500
New Mexico	529	.	.	37	18	*	.	13	598
New York	6,595	*	15	162	121	15	25	126	7,060
North Carolina	3,145	.	*	125	120	.	*	63	3,460
North Dakota	171	.	.	16	*	.	.	*	201
Ohio	4,151	.	36	257	73	.	14	100	4,631
Oklahoma	1,076	.	*	45	70	.	*	18	1,212
Oregon	764	.	.	50	18	.	*	23	859
Pennsylvania	4,462	*	*	97	117	.	*	115	4,803
Rhode Island	315	.	.	*	*	.	.	*	335
South Carolina	1,659	.	*	90	56	.	*	28	1,842
South Dakota	210	.	.	18	*	.	.	16	250
Tennessee	2,032	.	*	88	90	.	*	34	2,251
Texas	7,722	.	21	245	214	*	11	84	8,300
Utah	377	.	.	16	11	.	*	17	422
Vermont	143	.	.	.	*	.	*	*	154
Virginia	2,508	.	*	124	35	*	*	66	2,743
Washington	1,378	17	*	97	17	*	*	28	1,540
West Virginia	633	.	*	41	58	*	*	*	740
Wisconsin	1,506	.	*	71	50	*	*	104	1,742
Wyoming	80	.	.	*	*	.	*	*	96
American Samoa	*	*
Guam	80	.	.	*	.	.	.	*	86
Puerto Rico	1,146	.	.	*	69	*	.	*	1,225
Saipan	19	19
Virgin Islands	52	52
Foreign	17	17
Unknown	108	.	*	*	*	.	191	270	577
All	96,684	31	428	4,673	2,202	47	423	2,424	106,912

on our website & CD

Incident ESRD patients for selected years: by state (Table D.e).

Table D.16

Point prevalent ESRD patients, 2005: by modality & state/territory*December 31 point prevalent ESRD patients*

	Center hemo	Center self hemo	Home hemo	CAPD	CCPD	Other PD	Uncertain dialysis	Transplant	All
Alabama	6,088	.	26	245	343	.	*	2,314	9,022
Alaska	233	.	*	34	45	.	*	257	574
Arizona	5,719	.	25	195	204	.	18	2,519	8,680
Arkansas	2,426	.	*	168	184	.	*	1,190	3,985
California	38,445	16	116	1,571	1,810	*	186	16,230	58,376
Colorado	2,870	.	*	84	159	.	19	1,829	4,968
Connecticut	2,841	.	17	156	366	.	*	1,472	4,855
Delaware	1,070	.	*	18	48	.	*	499	1,639
District of Columbia	1,687	*	*	36	23	.	*	375	2,137
Florida	17,757	*	199	427	899	*	50	7,413	26,750
Georgia	11,864	*	65	482	560	*	16	3,498	16,489
Hawaii	2,092	.	*	91	53	.	.	607	2,845
Idaho	657	*	*	93	71	.	.	609	1,442
Illinois	13,332	*	364	488	713	*	78	7,276	22,257
Indiana	5,763	.	54	318	280	*	26	3,118	9,560
Iowa	2,112	.	21	65	95	.	13	1,724	4,030
Kansas	2,073	.	16	141	120	.	*	1,105	3,459
Kentucky	3,897	.	18	151	224	*	16	1,777	6,084
Louisiana	6,634	.	17	111	201	.	12	1,935	8,910
Maine	915	*	*	24	56	.	*	602	1,607
Maryland	7,054	*	17	254	344	*	26	3,367	11,070
Massachusetts	4,647	.	31	157	251	.	23	3,238	8,347
Michigan	11,053	*	58	525	384	.	25	5,512	17,559
Minnesota	3,219	*	23	94	136	.	14	3,489	6,976
Mississippi	4,801	.	45	205	259	.	*	1,202	6,514
Missouri	5,714	*	55	249	373	.	*	2,722	9,124
Montana	528	.	.	46	53	.	*	504	1,132
Nebraska	1,309	*	24	63	74	.	*	902	2,379
Nevada	1,841	.	*	100	97	.	*	845	2,894
New Hampshire	710	.	*	41	39	.	*	586	1,382
New Jersey	9,711	*	31	186	342	.	16	4,225	14,519
New Mexico	2,195	.	*	91	145	*	*	857	3,299
New York	21,922	36	103	526	632	*	98	8,708	32,033
North Carolina	11,406	.	38	306	659	*	11	3,649	16,071
North Dakota	434	.	.	46	*	.	.	436	924
Ohio	12,530	*	125	528	586	*	40	5,810	19,621
Oklahoma	3,278	*	14	115	294	.	*	1,425	5,134
Oregon	2,215	.	20	101	156	.	14	1,778	4,284
Pennsylvania	12,785	28	60	267	474	*	69	7,221	20,906
Rhode Island	815	.	*	15	15	.	*	561	1,410
South Carolina	6,386	.	28	198	336	.	*	1,947	8,905
South Dakota	622	.	*	34	27	.	*	502	1,189
Tennessee	6,246	.	38	192	361	*	11	2,754	9,603
Texas	27,444	*	150	618	1,312	*	50	9,546	39,125
Utah	1,062	*	11	80	25	.	*	1,201	2,388
Vermont	384	.	*	*	12	.	*	289	697
Virginia	8,471	*	73	267	413	*	17	3,685	12,928
Washington	4,225	*	95	185	212	*	*	2,869	7,605
West Virginia	1,650	*	13	74	193	.	*	951	2,886
Wisconsin	4,332	.	34	167	196	*	16	3,769	8,515
Wyoming	244	.	*	16	13	.	.	218	495
American Samoa	76	*	*	78
Guam	286	.	*	25	*	.	.	16	329
Puerto Rico	3,396	.	*	41	280	*	*	1,044	4,775
Saipan	95	.	.	*	.	.	.	12	114
Virgin Islands	160	.	.	*	*	.	.	30	196
Foreign	51	*	.	*	.	.	.	238	291
Unknown	147	.	*	*	*	.	253	1,235	1,646
All	311,919	138	2,105	10,732	15,163	37	1,225	143,693	485,012

on our website & CD

Living ESRD patients on December 31 for selected years: by state (Table D.f.)



121



