



SECTION I

Patient

Survival



Contents

- 691 I.1 **Patient counts**
patients alive on day 91
- 693 I.2 **Patient counts**
patients aged 65 & over, alive on day 1
- 693 I.3 **Patient counts**
dialysis patients alive on day 91
- 695 I.4 **Patient counts**
first renal transplant, cadaveric donor
- 696 I.5 **Patient counts**
first renal transplant, living donor
- 697 I.6 **One-year survival probabilities**
all patients, from day 91 to one year + 90 days, unadjusted
- 699 I.7 **Two-year survival probabilities**
all patients, from day 91 to two years + 90 days, unadjusted
- 701 I.8 **Five-year survival probabilities**
all patients, from day 91 to five years + 90 days, unadjusted
- 703 I.9 **Ten-year survival probabilities**
all patients, from day 91 to ten years + 90 days, unadjusted
- 705 I.10 **One-year survival probabilities**
all patients, from day 91 to one year + 90 days, adjusted for age, gender, race, & primary diagnosis
- 706 I.11 **Two-year survival probabilities**
all patients, from day 91 to two years + 90 days, adjusted for age, gender, race, & primary diagnosis
- 707 I.12 **Five-year survival probabilities**
all patients, from day 91 to five years + 90 days, adjusted for age, gender, race, & primary diagnosis
- 708 I.13 **Ten-year survival probabilities**
all patients, from day 91 to ten years + 90 days, adjusted for age, gender, race, & primary diagnosis
- 709 I.14 **One-year survival probabilities**
patients aged 65 & over, from day 1 to one year, unadjusted
- 709 I.15 **Two-year survival probabilities**
patients aged 65 & over, from day 1 to two years, unadjusted
- 710 I.16 **Five-year survival probabilities**
patients aged 65 & over, from day 1 to five years, unadjusted
- 710 I.17 **Ten-year survival probabilities**
patients aged 65 & over, from day 1 to ten years, unadjusted
- 711 I.18 **One-year survival probabilities**
patients aged 65 & over, from day 1 to one year, adjusted for age, gender, race, & primary diagnosis
- 711 I.19 **Two-year survival probabilities**
patients aged 65 & over, from day 1 to two years, adjusted for age, gender, race, & primary diagnosis

- 712 [I.20 Five-year survival probabilities](#)
patients aged 65 & over, from day 1 to five years, adjusted for age, gender, race, & primary diagnosis
- 712 [I.21 Ten-year survival probabilities](#)
patients aged 65 & over, from day 1 to ten years, adjusted for age, gender, race, & primary diagnosis
- 713 [I.22 One-year survival probabilities](#)
dialysis patients (censored at transplant), from day 91 to one year + 90 days, unadjusted
- 715 [I.23 Two-year survival probabilities](#)
dialysis patients (censored at transplant), from day 91 to two years + 90 days, unadjusted
- 717 [I.24 Five-year survival probabilities](#)
dialysis patients (censored at transplant), from day 91 to five years + 90 days, unadjusted
- 719 [I.25 Ten-year survival probabilities](#)
dialysis patients (censored at transplant), from day 91 to ten years + 90 days, unadjusted
- 721 [I.26 One-year survival probabilities](#)
dialysis patients (censored at transplant), from day 91 to one year + 90 days, adjusted for age, gender, race, & primary diagnosis
- 722 [I.27 Two-year survival probabilities](#)
dialysis patients (censored at transplant), from day 91 to two years + 90 days, adjusted for age, gender, race, & primary diagnosis
- 723 [I.28 Five-year survival probabilities](#)
dialysis patients (censored at transplant), from day 91 to five years + 90 days, adjusted for age, gender, race, & primary diagnosis
- 724 [I.29 Ten-year survival probabilities](#)
dialysis patients (censored at transplant), from day 91 to ten years + 90 days, adjusted for age, gender, race, & primary diagnosis
- 725 [I.30 One-year survival probabilities](#)
transplant patients (cadaveric donor), from day of transplant to one year, unadjusted
- 726 [I.31 Two-year survival probabilities](#)
transplant patients (cadaveric donor), from day of transplant to two years, unadjusted
- 727 [I.32 Five-year survival probabilities](#)
transplant patients (cadaveric donor), from day of transplant to five years, unadjusted
- 728 [I.33 Ten-year survival probabilities](#)
transplant patients (cadaveric donor), from day of transplant to ten years, unadjusted
- 729 [I.34 One-year survival probabilities](#)
transplant patients (cadaveric donor), from day of transplant to one year, adjusted for age, gender, race, & primary diagnosis
- 730 [I.35 Two-year survival probabilities](#)
transplant patients (cadaveric donor), from day of transplant to two years, adjusted for age, gender, race, & primary diagnosis
- 731 [I.36 Five-year survival probabilities](#)
transplant patients (cadaveric donor), from day of transplant to five years, adjusted for age, gender, race, & primary diagnosis
- 732 [I.37 Ten-year survival probabilities](#)
transplant patients (cadaveric donor), from day of transplant to ten years, adjusted for age, gender, race, & primary diagnosis
- 733 [I.38 One-year survival probabilities](#)
transplant patients (living donor), from day of transplant to one year, unadjusted
- 734 [I.39 Two-year survival probabilities](#)
transplant patients (living donor), from day of transplant to two years, unadjusted
- 735 [I.40 Five-year survival probabilities](#)
transplant patients (living donor), from day of transplant to five years, unadjusted
- 736 [I.41 Ten-year survival probabilities](#)
transplant patients (living donor), from day of transplant to ten years, unadjusted
- 737 [I.42 One-year survival probabilities](#)
transplant patients (living donor), from day of transplant to one year, adjusted for age, gender, race, & primary diagnosis

- 738 **I.43 Two-year survival probabilities**
transplant patients (living donor), from day of transplant to two years, adjusted for age, gender, race, & primary diagnosis
- 739 **I.44 Five-year survival probabilities**
transplant patients (living donor), from day of transplant to five years, adjusted for age, gender, race, & primary diagnosis
- 740 **I.45 Ten-year survival probabilities**
transplant patients (living donor), from day of transplant to ten years, adjusted for age, gender, race, & primary diagnosis

Table I.1

Patient Counts*patients alive on day 91*

	1978	1979	1980	1981	1982	1983	1984	1985	1986	1987
0-4	25	25	40	64	91	93	97	113	107	109
5-9	73	84	85	91	101	88	99	90	110	96
10-14	199	175	163	183	204	196	212	193	180	198
15-19	390	427	403	456	439	419	400	393	408	438
20-24	603	605	599	682	722	659	733	741	715	765
25-29	802	813	824	936	1,104	1,084	1,122	1,170	1,263	1,206
30-34	813	937	1,063	1,254	1,338	1,362	1,431	1,443	1,628	1,540
35-39	900	895	1,004	1,143	1,292	1,346	1,446	1,634	1,735	1,861
40-44	964	1,021	1,046	1,248	1,295	1,438	1,481	1,611	1,751	1,989
45-49	1,106	1,246	1,227	1,365	1,448	1,611	1,662	1,835	1,943	2,108
50-54	1,442	1,548	1,612	1,836	2,014	2,028	2,065	2,210	2,331	2,504
55-59	1,610	1,778	1,872	2,283	2,421	2,522	2,714	2,871	2,991	3,209
60-64	1,530	1,771	1,947	2,244	2,563	2,992	3,280	3,589	3,739	4,223
65-69	1,513	1,714	1,937	1,972	2,273	3,054	3,091	3,516	4,023	4,583
70-74	1,018	1,239	1,366	1,478	1,840	2,575	2,547	3,001	3,363	3,857
75-79	495	695	778	834	1,052	1,576	1,725	1,971	2,360	2,676
80-84	145	236	272	298	396	638	674	850	1,027	1,213
85+	38	64	86	79	141	207	236	283	354	398
White	8,599	9,437	10,069	11,681	13,816	15,540	16,333	17,928	19,606	21,274
Black	4,568	5,270	5,626	6,045	6,277	7,463	7,611	8,385	9,021	10,077
Other	499	566	629	720	641	885	1,071	1,201	1,401	1,622
Female	6,093	6,814	7,220	8,254	9,325	10,777	11,205	12,562	13,606	15,145
Male	7,573	8,459	9,104	10,192	11,409	13,111	13,810	14,952	16,422	17,828
Diabetes	1,375	1,577	2,183	3,477	4,800	5,608	6,752	7,782	8,977	10,041
Hypertension	1,821	2,000	2,464	3,819	5,150	5,596	6,245	7,143	7,771	8,803
Glomerulonephritis	1,856	1,965	2,041	2,945	3,676	3,850	4,072	4,256	4,369	4,629
Cystic Kidney	534	578	611	828	984	1,014	1,047	1,105	1,178	1,210
Other Urologic	206	230	260	382	389	471	478	603	608	609
Other Cause	1,161	1,371	1,524	2,156	2,972	3,292	3,507	3,851	3,888	4,060
Unknown Cause	1,190	1,277	1,307	1,510	1,295	1,303	1,339	1,340	1,568	1,836
Missing Cause	5,523	6,275	5,934	3,329	1,468	2,754	1,575	1,434	1,669	1,785
Total	13,666	15,273	16,324	18,446	20,734	23,888	25,015	27,514	30,028	32,973

Table I.1 (continued)

	1988	1989	1990	1991	1992	1993	1994	1995	1996	1997
0-4	112	121	102	133	119	123	112	161	188	182
5-9	86	103	121	111	89	101	138	138	139	122
10-14	186	173	183	209	198	205	225	256	268	275
15-19	427	478	435	414	443	448	468	539	573	520
20-24	796	787	825	846	913	858	897	1,019	1,024	944
25-29	1,275	1,442	1,438	1,450	1,459	1,471	1,559	1,580	1,645	1,673
30-34	1,696	1,926	1,956	2,013	2,085	2,088	2,227	2,429	2,389	2,369
35-39	2,056	2,170	2,285	2,463	2,682	2,588	2,897	3,195	3,244	3,304
40-44	2,130	2,395	2,707	3,035	3,179	3,128	3,357	3,971	4,166	4,161
45-49	2,317	2,708	2,777	3,189	3,486	3,692	4,224	4,793	5,171	5,329
50-54	2,851	3,073	3,245	3,487	4,019	4,104	4,882	5,361	5,591	6,244
55-59	3,534	3,828	4,117	4,466	4,860	4,905	5,719	6,274	6,510	7,199
60-64	4,578	5,016	5,366	5,972	6,281	6,409	7,096	7,781	8,005	8,539
65-69	5,105	5,912	6,484	7,310	7,841	8,123	8,862	8,808	9,152	9,918
70-74	4,176	4,897	5,428	6,529	7,271	7,623	8,558	8,580	9,159	10,111
75-79	3,021	3,637	3,963	4,766	5,246	5,450	6,222	6,453	7,029	8,125
80-84	1,454	1,748	1,987	2,362	2,753	2,893	3,366	3,597	4,040	4,869
85+	497	655	775	900	1,060	1,227	1,316	1,468	1,741	2,074
White	23,200	26,297	28,548	32,047	34,648	34,872	38,973	40,397	42,019	47,316
Black	11,327	13,189	13,791	15,498	16,927	17,867	19,630	21,071	22,320	23,212
Other	1,770	1,583	1,855	2,110	2,409	2,697	3,522	4,935	5,695	5,430
Female	16,735	18,834	20,323	23,030	25,376	25,775	29,153	31,221	32,554	35,635
Male	19,562	22,235	23,871	26,625	28,608	29,661	32,970	35,182	37,461	40,323
Diabetes	11,297	13,757	15,314	17,934	19,753	19,516	22,923	25,877	28,206	31,053
Hypertension	9,959	11,734	12,522	14,387	15,879	15,119	17,357	15,968	16,376	18,004
Glomerulonephritis	4,914	5,262	5,522	5,851	6,065	5,902	6,475	6,789	6,774	6,803
Cystic Kidney	1,186	1,207	1,368	1,420	1,510	1,464	1,568	1,676	1,592	1,635
Other Urologic	666	730	772	1,025	1,056	933	984	1,171	1,315	1,307
Other Cause	4,106	4,574	4,390	4,658	4,900	4,803	5,334	6,925	6,943	7,381
Unknown Cause	1,799	1,668	1,768	2,080	2,284	2,003	2,112	2,257	2,425	2,566
Missing Cause	2,370	2,137	2,538	2,300	2,537	5,696	5,372	5,740	6,403	7,209
Total	36,297	41,069	44,194	49,655	53,984	55,436	62,125	66,403	70,034	75,958

Table I.2

Patient Counts*patients aged 65 & over, alive on day 1*

	1978-79	1980-81	1982-83	1984-85	1986-87	1988-89	1990-91	1992-93	1994-95	1996-97
65-69	3,534	4,315	5,876	7,345	9,423	11,956	14,972	17,239	19,078	20,730
70-74	2,553	3,231	4,986	6,342	8,076	10,141	13,319	16,472	18,946	21,585
75-79	1,359	1,861	3,019	4,246	5,770	7,671	9,869	12,189	14,415	17,459
80-84	450	678	1,212	1,820	2,637	3,729	5,102	6,604	8,188	10,543
85 +	142	203	435	645	934	1,414	2,058	2,744	3,435	4,798
White	5,203	6,911	11,132	14,550	19,163	25,120	32,685	39,421	45,467	53,321
Black	2,544	2,996	3,974	5,117	6,619	8,591	11,104	13,665	15,160	17,333
Other	291	381	422	731	1,058	1,200	1,531	2,162	3,435	4,461
Female	3,657	4,640	7,204	9,611	12,833	16,785	21,942	27,064	31,857	36,489
Male	4,381	5,648	8,324	10,787	14,007	18,126	23,378	28,184	32,203	38,613
Diabetes	513	1,234	2,629	4,224	6,456	9,540	14,032	17,656	23,115	30,199
Hypertension	1,238	2,590	5,227	7,283	10,094	13,817	18,155	21,123	23,123	25,239
Glomerulonephritis	582	1,030	1,835	2,390	2,874	3,446	4,107	4,243	5,061	5,565
Cystic Kidney	160	270	423	465	572	599	733	754	942	988
Other Urologic	207	342	441	635	776	923	1,111	1,150	1,263	1,544
Other Cause	547	984	1,965	2,611	3,144	3,513	3,639	3,939	5,174	6,800
Unknown Cause	834	1,190	1,248	1,376	1,874	2,060	2,320	2,586	2,711	3,348
Missing Cause	3,957	2,648	1,760	1,414	1,050	1,013	1,223	3,797	2,673	1,432
Total	8,038	10,288	15,528	20,398	26,840	34,911	45,320	55,248	64,062	75,115

Table I.3

Patient Counts*dialysis patients alive on day 91*

	1978	1979	1980	1981	1982	1983	1984	1985	1986	1987
0-4	18	16	26	48	62	74	80	88	87	79
5-9	48	53	54	63	73	67	71	54	76	67
10-14	144	120	129	147	162	150	151	133	118	123
15-19	321	350	349	374	359	364	326	322	308	336
20-24	523	517	525	590	643	592	654	646	626	681
25-29	688	701	711	824	973	957	1,003	1,044	1,111	1,060
30-34	736	842	959	1,094	1,215	1,239	1,309	1,308	1,437	1,370
35-39	837	818	937	1,070	1,194	1,246	1,336	1,498	1,570	1,676
40-44	910	953	983	1,189	1,218	1,365	1,396	1,512	1,629	1,836
45-49	1,056	1,187	1,170	1,314	1,395	1,562	1,602	1,770	1,843	1,998
50-54	1,400	1,513	1,575	1,806	1,957	1,977	2,010	2,168	2,262	2,412
55-59	1,589	1,756	1,855	2,268	2,395	2,501	2,689	2,837	2,933	3,134
60-64	1,525	1,761	1,942	2,235	2,557	2,979	3,272	3,570	3,716	4,188
65-69	1,512	1,714	1,933	1,971	2,273	3,051	3,082	3,507	4,015	4,570
70-74	1,018	1,238	1,366	1,477	1,840	2,573	2,542	2,999	3,358	3,853
75-79	494	695	778	834	1,052	1,576	1,723	1,968	2,359	2,674
80-84	145	236	272	298	396	638	673	850	1,026	1,212
85 +	37	64	85	78	141	206	236	281	354	398
White	8,046	8,849	9,513	11,061	13,124	14,897	15,600	17,096	18,562	20,170
Black	4,488	5,160	5,546	5,948	6,165	7,364	7,514	8,294	8,904	9,947
Other	467	525	590	671	616	856	1,041	1,165	1,362	1,550
Female	5,861	6,561	6,984	7,984	9,031	10,508	10,876	12,185	13,160	14,649
Male	7,140	7,973	8,665	9,696	10,874	12,609	13,279	14,370	15,668	17,018
Diabetes	1,326	1,510	2,107	3,346	4,623	5,441	6,542	7,565	8,680	9,732
Hypertension	1,789	1,962	2,429	3,770	5,091	5,549	6,185	7,083	7,698	8,728
Glomerulonephritis	1,720	1,837	1,933	2,817	3,486	3,674	3,874	4,045	4,143	4,360
Cystic Kidney	516	559	602	803	960	984	1,001	1,053	1,103	1,135
Other Urologic	195	217	257	373	371	457	460	579	581	596
Other Cause	1,104	1,282	1,429	2,037	2,857	3,179	3,358	3,691	3,708	3,851
Unknown Cause	1,160	1,234	1,272	1,472	1,254	1,266	1,311	1,289	1,524	1,783
Missing Cause	5,191	5,933	5,620	3,062	1,263	2,567	1,424	1,250	1,391	1,482
Total	13,001	14,534	15,649	17,680	19,905	23,117	24,155	26,555	28,828	31,667

Table I.3 (continued)

	1988	1989	1990	1991	1992	1993	1994	1995	1996	1997
0-4	87	96	77	108	92	89	87	136	148	143
5-9	58	63	82	62	50	66	94	96	98	84
10-14	125	109	131	143	149	149	156	184	204	212
15-19	349	399	351	329	382	354	348	435	477	438
20-24	687	682	727	748	823	766	792	926	942	869
25-29	1,122	1,252	1,258	1,296	1,314	1,324	1,393	1,430	1,531	1,554
30-34	1,508	1,764	1,745	1,793	1,873	1,899	1,999	2,225	2,185	2,202
35-39	1,876	1,975	2,108	2,268	2,490	2,384	2,657	2,919	3,033	3,102
40-44	1,971	2,243	2,484	2,798	2,975	2,948	3,116	3,746	3,935	3,957
45-49	2,201	2,583	2,651	3,034	3,322	3,531	3,995	4,564	4,949	5,115
50-54	2,743	2,999	3,144	3,391	3,891	3,970	4,723	5,181	5,402	6,071
55-59	3,474	3,751	4,033	4,377	4,763	4,819	5,581	6,151	6,384	7,040
60-64	4,547	4,974	5,328	5,917	6,219	6,343	7,009	7,690	7,916	8,447
65-69	5,098	5,900	6,462	7,287	7,817	8,091	8,822	8,764	9,108	9,851
70-74	4,171	4,894	5,424	6,525	7,266	7,613	8,545	8,568	9,149	10,101
75-79	3,020	3,635	3,961	4,765	5,245	5,447	6,218	6,452	7,027	8,121
80-84	1,454	1,748	1,987	2,362	2,752	2,893	3,366	3,597	4,040	4,869
85+	497	655	774	900	1,060	1,227	1,316	1,468	1,741	2,073
White	22,122	25,184	27,318	30,734	33,380	33,600	37,554	39,008	40,656	45,886
Black	11,149	13,015	13,621	15,333	16,764	17,694	19,267	20,699	22,026	23,021
Other	1,717	1,523	1,788	2,036	2,339	2,619	3,396	4,825	5,587	5,342
Female	16,234	18,305	19,714	22,390	24,716	25,172	28,328	30,420	31,853	34,918
Male	18,754	21,417	23,013	25,713	27,767	28,741	31,887	34,112	36,397	39,331
Diabetes	11,017	13,457	15,011	17,586	19,440	19,200	22,574	25,515	27,795	30,667
Hypertension	9,863	11,663	12,429	14,282	15,761	15,038	17,257	15,872	16,284	17,908
Glomerulonephritis	4,683	5,002	5,248	5,592	5,778	5,620	6,170	6,454	6,454	6,477
Cystic Kidney	1,098	1,131	1,234	1,292	1,373	1,344	1,449	1,508	1,431	1,486
Other Urologic	648	709	741	982	1,019	888	948	1,119	1,255	1,250
Other Cause	3,930	4,377	4,197	4,439	4,701	4,578	5,104	6,663	6,684	7,095
Unknown Cause	1,758	1,624	1,704	2,022	2,232	1,951	2,052	2,208	2,372	2,517
Missing Cause	1,991	1,759	2,163	1,908	2,179	5,294	4,663	5,193	5,994	6,849
Total	34,988	39,722	42,727	48,103	52,483	53,913	60,217	64,532	68,269	74,249

Table I.4

Patient Counts*first renal transplant, cadaveric donor*

	1978	1979	1980	1981	1982	1983	1984	1985	1986	1987
0-4	7	10	10	11	12	17	29	32	30	36
5-9	17	25	19	32	27	25	34	38	45	34
10-14	53	62	61	56	66	61	73	77	81	89
15-19	121	125	116	116	120	143	149	150	165	184
20-24	183	195	212	250	246	250	292	315	351	288
25-29	221	236	279	278	350	399	432	423	542	522
30-34	243	281	303	377	455	502	597	583	774	671
35-39	246	258	266	326	399	529	609	672	803	798
40-44	230	234	275	319	379	416	518	583	736	805
45-49	195	238	254	252	343	382	450	572	682	687
50-54	166	170	207	219	288	336	409	435	599	627
55-59	97	102	111	125	153	211	271	336	468	502
60-64	32	38	25	43	61	73	150	181	256	290
65+	7	4	15	11	7	27	35	61	100	145
White	1,342	1,467	1,555	1,687	2,061	2,411	2,925	3,175	4,075	4,060
Black	393	416	505	564	753	848	972	1,112	1,300	1,325
Other	83	95	93	164	92	112	151	171	257	293
Female	669	729	796	863	1,041	1,241	1,508	1,643	2,070	2,129
Male	1,149	1,249	1,357	1,552	1,865	2,130	2,540	2,815	3,562	3,549
Diabetes	138	155	192	274	403	510	692	881	1,159	1,175
Hypertension	197	173	232	277	405	490	569	687	842	889
Glomerulonephritis	457	486	497	567	765	861	1,086	1,087	1,416	1,423
Cystic Kidney	71	100	98	124	181	214	299	331	423	488
Other Urologic	13	30	22	29	36	41	51	60	75	75
Other Cause	216	265	309	326	387	563	644	752	899	837
Unknown Cause	173	182	168	184	201	218	224	224	246	239
Missing Cause	553	587	635	634	528	474	483	436	572	552
Total	1,818	1,978	2,153	2,415	2,906	3,371	4,048	4,458	5,632	5,678

	1988	1989	1990	1991	1992	1993	1994	1995	1996	1997
0-4	29	28	41	28	31	25	33	25	33	37
5-9	45	43	31	34	26	36	39	39	53	44
10-14	74	66	60	59	50	55	86	76	76	83
15-19	162	150	154	128	121	129	213	159	120	108
20-24	291	271	309	302	268	270	241	268	222	184
25-29	458	557	552	522	479	493	473	493	437	464
30-34	678	648	687	709	711	740	727	703	648	603
35-39	806	766	824	784	790	822	885	892	824	762
40-44	771	831	887	909	869	935	963	922	961	944
45-49	750	696	775	790	814	906	976	1,017	1,061	1,064
50-54	613	597	684	666	707	779	848	863	886	966
55-59	520	535	625	580	606	664	746	724	773	867
60-64	372	349	408	477	457	536	532	588	636	653
65+	193	187	255	298	349	378	437	479	524	542
White	4,050	4,050	4,430	4,428	4,375	4,793	4,712	4,806	4,741	4,771
Black	1,378	1,354	1,498	1,496	1,517	1,610	2,026	2,040	2,043	2,044
Other	334	320	364	362	386	365	461	402	470	506
Female	2,282	2,241	2,465	2,446	2,499	2,670	2,783	2,711	2,849	2,927
Male	3,480	3,483	3,827	3,840	3,779	4,098	4,416	4,537	4,405	4,394
Diabetes	1,220	1,320	1,448	1,511	1,546	1,837	1,874	1,910	1,986	2,002
Hypertension	946	894	1,049	1,037	1,093	1,152	1,332	1,270	1,241	1,189
Glomerulonephritis	1,367	1,333	1,473	1,443	1,391	1,417	1,475	1,520	1,446	1,417
Cystic Kidney	519	464	543	504	545	572	513	579	582	584
Other Urologic	74	59	85	103	83	132	111	102	118	113
Other Cause	826	805	835	819	825	837	821	835	799	758
Unknown Cause	271	254	273	244	252	263	290	232	229	262
Missing Cause	539	595	586	625	543	558	783	800	853	996
Total	5,762	5,724	6,292	6,286	6,278	6,768	7,199	7,248	7,254	7,321

Table I.5

Patient Counts*first renal transplant, living donor*

	1978	1979	1980	1981	1982	1983	1984	1985	1986	1987
0-4	6	6	8	18	32	36	40	36	38	57
5-9	22	24	28	24	40	29	41	51	39	53
10-14	68	51	48	42	61	82	96	83	79	68
15-19	109	110	74	114	141	147	131	132	141	130
20-24	155	137	125	161	212	215	187	191	199	177
25-29	173	168	165	185	264	267	241	272	267	255
30-34	124	131	129	231	241	270	254	257	267	234
35-39	94	99	91	118	175	172	178	199	216	202
40-44	83	64	76	80	126	120	127	140	137	162
45-49	66	56	53	72	93	96	97	120	112	120
50-54	47	44	30	51	68	77	77	89	78	87
55-59	16	17	12	22	48	41	37	50	57	67
60-64	6	6	.	3	14	12	12	13	21	30
65+	4	1	4	3	1	4	3	6	5	13
White	817	753	720	935	1,276	1,315	1,266	1,378	1,403	1,399
Black	99	111	84	105	184	200	199	202	193	182
Other	57	50	39	84	56	53	56	59	60	74
Female	398	348	354	441	569	610	577	677	668	692
Male	575	566	489	683	947	958	944	962	988	963
Diabetes	82	66	88	139	260	275	296	320	369	339
Hypertension	67	56	50	86	137	140	131	136	145	146
Glomerulonephritis	250	224	225	305	452	495	445	468	446	460
Cystic Kidney	24	25	27	32	43	52	61	61	62	63
Other Urologic	9	9	5	19	37	33	27	52	47	33
Other Cause	132	144	131	175	234	292	333	327	342	339
Unknown Cause	98	91	67	84	103	94	84	92	65	82
Missing Cause	311	299	250	284	250	187	144	183	180	193
Total	973	914	843	1,124	1,516	1,568	1,521	1,639	1,656	1,655

	1988	1989	1990	1991	1992	1993	1994	1995	1996	1997
0-4	42	46	35	53	50	69	30	50	58	64
5-9	31	44	50	69	52	48	49	58	57	41
10-14	81	83	62	80	90	89	81	90	82	67
15-19	125	114	127	134	122	139	155	155	151	148
20-24	168	181	166	187	179	210	218	200	221	228
25-29	255	248	288	276	271	291	298	271	294	287
30-34	256	231	255	285	289	303	332	327	351	329
35-39	206	217	217	261	285	283	313	370	337	361
40-44	139	162	202	238	253	266	303	333	330	353
45-49	105	131	129	176	202	215	273	301	399	347
50-54	90	95	114	125	152	191	216	272	286	335
55-59	49	72	78	98	135	138	163	192	220	287
60-64	25	43	34	54	76	93	98	141	160	193
65+	8	7	23	22	48	50	55	92	124	136
White	1,299	1,393	1,484	1,667	1,782	1,884	1,991	2,237	2,357	2,453
Black	205	211	215	295	307	388	446	456	511	505
Other	76	70	81	96	115	113	147	159	202	218
Female	603	692	768	879	941	975	1,105	1,211	1,292	1,317
Male	977	982	1,012	1,179	1,263	1,410	1,479	1,641	1,778	1,859
Diabetes	311	315	345	429	407	460	512	539	627	586
Hypertension	136	155	149	213	276	297	350	344	368	358
Glomerulonephritis	461	483	521	555	601	625	699	772	827	867
Cystic Kidney	59	67	86	99	117	120	128	180	185	198
Other Urologic	26	38	50	56	75	80	73	88	100	94
Other Cause	298	338	336	383	377	431	428	491	516	535
Unknown Cause	62	71	91	85	105	122	118	123	119	116
Missing Cause	227	207	202	238	246	250	276	315	328	422
Total	1,580	1,674	1,780	2,058	2,204	2,385	2,584	2,852	3,070	3,176

Table I.6

One-year Survival Probabilities*all patients, from day 91 to one year + 90 days, unadjusted*

PROBABILITIES	1987	1988	1989	1990	1991	1992	1993	1994	1995	1996	1997
0-4	91.7	83.0	85.1	90.2	85.7	91.6	88.6	92.9	93.8	95.7	93.8
5-9	99.0	100.0	98.1	97.5	98.2	97.8	97.0	100.0	98.6	97.8	96.7
10-14	98.5	96.8	95.4	97.3	97.6	94.9	95.6	99.6	98.8	97.8	98.5
15-19	96.1	97.4	98.3	97.7	96.1	96.8	97.3	96.4	98.5	97.7	98.5
20-24	93.7	94.6	95.3	94.7	95.2	95.3	96.3	97.0	96.2	95.7	97.7
25-29	91.6	92.1	91.7	92.9	92.0	93.4	92.0	92.4	92.7	95.7	96.5
30-34	88.1	91.0	89.6	91.0	90.9	91.3	90.6	90.9	90.7	92.7	94.0
35-39	89.8	89.4	90.0	90.5	89.2	90.2	89.8	90.4	91.8	92.2	93.6
40-44	89.3	90.1	88.5	90.3	89.2	88.8	89.1	90.3	91.4	92.6	92.5
45-49	87.1	87.6	88.7	88.7	87.6	88.9	89.9	89.1	90.7	90.8	92.3
50-54	85.0	85.2	85.7	86.8	86.1	86.6	87.5	88.3	89.1	88.7	89.5
55-59	82.1	82.1	81.4	82.5	83.2	84.3	83.5	84.2	86.2	86.6	87.3
60-64	76.6	76.7	78.2	78.9	79.5	80.0	81.4	81.5	82.0	83.8	84.5
65-69	73.1	73.3	73.9	75.6	75.0	75.4	77.1	76.4	77.5	77.9	76.7
70-74	68.1	68.0	68.9	70.6	69.8	70.4	72.7	71.7	72.6	72.8	72.0
75-79	61.7	63.1	65.1	65.8	65.4	65.6	66.0	67.6	67.3	67.8	66.3
80-84	56.5	58.0	59.9	61.8	60.1	60.4	64.3	63.0	62.6	62.5	61.5
85+	52.0	47.7	51.6	53.7	55.9	53.3	52.4	55.4	55.4	53.7	55.0
White	75.7	75.8	76.2	77.1	75.9	76.1	77.0	77.3	78.1	78.1	77.4
Black	83.1	83.0	82.6	83.4	83.1	83.6	83.6	83.4	84.6	85.4	85.5
Other	80.5	82.3	85.7	86.5	87.4	85.7	87.5	86.1	86.3	86.5	87.5
Female	78.3	79.0	78.8	79.5	78.6	79.1	79.3	79.7	80.7	81.0	80.3
Male	78.1	77.8	78.4	79.4	78.6	78.7	79.9	79.8	80.8	81.1	80.8
Diabetes	74.8	75.8	76.6	76.7	76.9	77.1	78.9	78.4	79.1	79.4	78.9
Hypertension	75.6	75.4	75.9	77.3	76.0	76.2	76.8	76.6	78.6	78.0	77.0
Glomerulonephritis	86.5	86.4	86.3	87.0	85.5	87.2	88.2	86.5	88.0	88.5	88.1
Cystic Kidney	90.7	91.7	92.2	92.8	93.2	93.3	93.6	93.1	93.5	92.7	93.8
Other Urologic	74.1	73.0	73.7	77.7	79.9	78.1	80.0	78.6	80.4	80.8	82.3
Other Cause	77.2	78.9	77.5	78.4	76.4	75.7	76.3	75.9	76.3	75.8	77.3
Unknown Cause	78.4	74.3	76.1	79.5	75.3	76.7	78.2	76.9	78.3	78.6	75.9
Missing Cause	83.9	83.6	85.6	85.3	88.9	89.4	80.3	88.3	88.6	92.6	91.2
Total Crude Prob.	78.2	78.4	78.6	79.5	78.6	78.9	79.6	79.7	80.8	81.1	80.6

Table I.6 (continued)

STANDARD ERRORS											
	1987	1988	1989	1990	1991	1992	1993	1994	1995	1996	1997
0-4	2.6	3.5	3.2	2.9	3.0	2.5	2.9	2.4	1.9	1.5	1.8
5-9	1.0	0.0	1.4	1.4	1.3	1.6	1.7	0.0	1.0	1.2	1.6
10-14	0.9	1.3	1.6	1.2	1.1	1.6	1.4	0.4	0.7	0.9	0.7
15-19	0.9	0.8	0.6	0.7	0.9	0.8	0.8	0.9	0.5	0.6	0.5
20-24	0.9	0.8	0.8	0.8	0.7	0.7	0.6	0.6	0.6	0.6	0.5
25-29	0.8	0.8	0.7	0.7	0.7	0.6	0.7	0.7	0.7	0.5	0.5
30-34	0.8	0.7	0.7	0.6	0.6	0.6	0.6	0.6	0.6	0.5	0.5
35-39	0.7	0.7	0.6	0.6	0.6	0.6	0.6	0.5	0.5	0.5	0.4
40-44	0.7	0.6	0.7	0.6	0.6	0.6	0.6	0.5	0.4	0.4	0.4
45-49	0.7	0.7	0.6	0.6	0.6	0.5	0.5	0.5	0.4	0.4	0.4
50-54	0.7	0.7	0.6	0.6	0.6	0.5	0.5	0.5	0.4	0.4	0.4
55-59	0.7	0.6	0.6	0.6	0.6	0.5	0.5	0.5	0.4	0.4	0.4
60-64	0.7	0.6	0.6	0.6	0.5	0.5	0.5	0.5	0.4	0.4	0.4
65-69	0.7	0.6	0.6	0.5	0.5	0.5	0.5	0.5	0.4	0.4	0.4
70-74	0.8	0.7	0.7	0.6	0.6	0.5	0.5	0.5	0.5	0.5	0.5
75-79	0.9	0.9	0.8	0.8	0.7	0.7	0.6	0.6	0.6	0.6	0.5
80-84	1.4	1.3	1.2	1.1	1.0	0.9	0.9	0.8	0.8	0.8	0.7
85+	2.5	2.2	1.9	1.8	1.7	1.5	1.4	1.4	1.3	1.2	1.1
White	0.3	0.3	0.3	0.2	0.2	0.2	0.2	0.2	0.2	0.2	0.2
Black	0.4	0.4	0.3	0.3	0.3	0.3	0.3	0.3	0.2	0.2	0.2
Other	1.0	0.9	0.9	0.8	0.7	0.7	0.6	0.6	0.5	0.5	0.5
Female	0.3	0.3	0.3	0.3	0.3	0.3	0.3	0.2	0.2	0.2	0.2
Male	0.3	0.3	0.3	0.3	0.3	0.2	0.2	0.2	0.2	0.2	0.2
Diabetes	0.4	0.4	0.4	0.3	0.3	0.3	0.3	0.3	0.3	0.2	0.2
Hypertension	0.5	0.4	0.4	0.4	0.4	0.3	0.3	0.3	0.3	0.3	0.3
Glomerulonephritis	0.5	0.5	0.5	0.5	0.5	0.4	0.4	0.4	0.4	0.4	0.4
Cystic Kidney	0.8	0.8	0.8	0.7	0.7	0.6	0.6	0.6	0.6	0.7	0.6
Other Urologic	1.8	1.7	1.6	1.5	1.3	1.3	1.3	1.3	1.2	1.1	1.1
Other Cause	0.7	0.6	0.6	0.6	0.6	0.6	0.6	0.6	0.5	0.5	0.5
Unknown Cause	1.0	1.0	1.0	1.0	0.9	0.9	0.9	0.9	0.9	0.8	0.9
Missing Cause	0.9	0.8	0.8	0.7	0.7	0.6	0.5	0.4	0.4	0.3	0.3
Total Crude Prob.	0.2	0.2	0.2	0.2	0.2	0.2	0.2	0.2	0.2	0.1	0.1

Table I.7

Two-year Survival Probabilities*all patients, from day 91 to two years + 90 days, unadjusted*

PROBABILITIES	1986	1987	1988	1989	1990	1991	1992	1993	1994	1995	1996
0-4	91.6	88.1	78.6	81.0	85.3	83.5	91.6	87.0	88.4	90.7	93.1
5-9	93.6	96.9	97.7	97.1	94.2	98.2	97.8	91.1	97.8	97.8	97.1
10-14	95.6	98.5	95.7	93.1	94.5	96.7	93.4	94.1	98.2	97.3	96.6
15-19	93.1	94.3	95.1	96.9	95.6	93.7	95.3	95.3	94.9	97.2	94.9
20-24	91.6	88.9	91.5	91.6	91.6	90.7	92.2	92.7	94.2	93.3	93.6
25-29	86.9	84.1	86.3	87.0	86.9	86.5	87.4	87.7	87.9	87.9	91.8
30-34	82.6	80.0	83.1	82.4	82.8	83.8	83.6	83.8	83.2	85.3	87.5
35-39	79.8	81.6	82.4	82.3	82.9	80.4	82.1	83.1	83.5	85.9	86.4
40-44	78.6	80.0	80.5	79.1	82.3	79.5	80.4	79.6	82.0	83.7	85.2
45-49	75.6	76.1	77.0	77.4	78.4	76.6	78.8	79.8	79.5	82.5	82.2
50-54	70.3	70.6	72.6	74.1	73.9	73.9	74.1	76.1	78.3	78.6	78.9
55-59	67.0	67.1	66.8	67.6	68.4	69.1	70.6	70.3	71.2	73.7	75.1
60-64	60.8	59.7	60.4	62.0	62.3	63.4	63.9	65.6	66.7	67.5	70.1
65-69	54.1	55.4	56.1	56.1	56.6	56.9	58.0	59.8	58.9	60.5	61.3
70-74	48.5	48.6	49.3	50.0	50.5	50.3	51.8	52.6	52.5	54.3	54.4
75-79	43.6	41.1	42.3	44.5	44.8	44.8	45.3	46.5	47.1	46.7	48.5
80-84	37.1	34.8	34.6	39.0	39.6	37.9	39.0	39.9	40.6	41.3	41.6
85 +	29.4	27.9	28.6	32.4	29.4	30.7	30.6	31.0	32.8	33.1	31.8
White	61.5	60.7	60.6	61.0	60.9	59.6	60.3	60.8	61.3	62.7	62.9
Black	69.8	68.8	70.2	69.7	70.3	69.7	70.8	71.2	71.5	73.6	74.5
Other	68.7	67.4	69.5	74.2	74.1	77.3	74.7	76.0	74.9	75.7	76.7
Female	64.6	63.9	64.4	64.5	64.4	63.4	64.4	64.2	65.0	66.5	67.6
Male	64.1	63.1	63.6	64.1	64.4	63.7	64.1	65.4	65.5	67.6	67.8
Diabetes	56.3	56.2	57.6	59.1	58.3	58.8	59.5	61.4	61.0	62.6	63.3
Hypertension	61.8	59.7	60.5	60.9	61.0	60.0	60.9	61.4	61.5	64.1	63.5
Glomerulonephritis	76.0	75.9	76.9	75.9	76.9	75.0	77.2	78.5	76.8	79.0	79.2
Cystic Kidney	83.5	83.1	83.1	84.3	86.3	85.6	86.1	87.2	86.0	87.4	85.4
Other Urologic	64.8	62.4	61.0	62.6	63.5	68.6	65.1	68.5	65.0	68.9	69.3
Other Cause	67.4	65.8	66.1	65.5	66.9	64.0	63.4	64.0	63.9	64.7	64.9
Unknown Cause	64.9	62.3	60.2	63.2	65.5	60.9	63.5	65.4	62.9	65.3	65.1
Missing Cause	67.0	73.0	73.2	75.7	74.3	79.1	80.2	66.0	78.4	78.9	85.5
Total Crude Prob.	64.3	63.5	64.0	64.3	64.4	63.5	64.2	64.9	65.3	67.1	67.7

Table I.7 (continued)

STANDARD ERRORS											
	1986	1987	1988	1989	1990	1991	1992	1993	1994	1995	1996
0-4	2.7	3.1	3.9	3.6	3.5	3.2	2.5	3.0	3.0	2.3	1.8
5-9	2.3	1.8	1.6	1.6	2.1	1.3	1.6	2.8	1.2	1.2	1.4
10-14	1.5	0.9	1.5	1.9	1.7	1.2	1.8	1.6	0.9	1.0	1.1
15-19	1.2	1.1	1.0	0.8	1.0	1.2	1.0	1.0	1.0	0.7	0.9
20-24	1.0	1.1	1.0	1.0	1.0	1.0	0.9	0.9	0.8	0.8	0.8
25-29	0.9	1.1	1.0	0.9	0.9	0.9	0.9	0.9	0.8	0.8	0.7
30-34	0.9	1.0	0.9	0.9	0.9	0.8	0.8	0.8	0.8	0.7	0.7
35-39	1.0	0.9	0.8	0.8	0.8	0.8	0.7	0.7	0.7	0.6	0.6
40-44	1.0	0.9	0.9	0.8	0.7	0.7	0.7	0.7	0.7	0.6	0.6
45-49	1.0	0.9	0.9	0.8	0.8	0.7	0.7	0.7	0.6	0.5	0.5
50-54	0.9	0.9	0.8	0.8	0.8	0.7	0.7	0.7	0.6	0.6	0.6
55-59	0.9	0.8	0.8	0.8	0.7	0.7	0.7	0.7	0.6	0.6	0.5
60-64	0.8	0.8	0.7	0.7	0.7	0.6	0.6	0.6	0.6	0.5	0.5
65-69	0.8	0.7	0.7	0.6	0.6	0.6	0.6	0.5	0.5	0.5	0.5
70-74	0.9	0.8	0.8	0.7	0.7	0.6	0.6	0.6	0.5	0.5	0.5
75-79	1.0	1.0	0.9	0.8	0.8	0.7	0.7	0.7	0.6	0.6	0.6
80-84	1.5	1.4	1.2	1.2	1.1	1.0	0.9	0.9	0.8	0.8	0.8
85+	2.4	2.2	2.0	1.8	1.6	1.5	1.4	1.3	1.3	1.2	1.1
White	0.3	0.3	0.3	0.3	0.3	0.3	0.3	0.3	0.2	0.2	0.2
Black	0.5	0.5	0.4	0.4	0.4	0.4	0.3	0.3	0.3	0.3	0.3
Other	1.2	1.2	1.1	1.1	1.0	0.9	0.9	0.8	0.7	0.6	0.6
Female	0.4	0.4	0.4	0.3	0.3	0.3	0.3	0.3	0.3	0.3	0.3
Male	0.4	0.4	0.3	0.3	0.3	0.3	0.3	0.3	0.3	0.2	0.2
Diabetes	0.5	0.5	0.5	0.4	0.4	0.4	0.3	0.3	0.3	0.3	0.3
Hypertension	0.6	0.5	0.5	0.5	0.4	0.4	0.4	0.4	0.4	0.4	0.4
Glomerulonephritis	0.6	0.6	0.6	0.6	0.6	0.6	0.5	0.5	0.5	0.5	0.5
Cystic Kidney	1.1	1.1	1.1	1.0	0.9	0.9	0.9	0.9	0.9	0.8	0.9
Other Urologic	1.9	2.0	1.9	1.8	1.7	1.4	1.5	1.5	1.5	1.4	1.3
Other Cause	0.8	0.7	0.7	0.7	0.7	0.7	0.7	0.7	0.7	0.6	0.6
Unknown Cause	1.2	1.1	1.2	1.2	1.1	1.1	1.0	1.1	1.1	1.0	1.0
Missing Cause	1.1	1.1	0.9	0.9	0.9	0.8	0.8	0.6	0.6	0.5	0.4
Total Crude Prob.	0.3	0.3	0.3	0.2	0.2	0.2	0.2	0.2	0.2	0.2	0.2

Table I.8

Five-year Survival Probabilities*all patients, from day 91 to five years + 90 days, unadjusted*

PROBABILITIES											
	1983	1984	1985	1986	1987	1988	1989	1990	1991	1992	1993
0-4	79.6	75.3	80.5	83.2	79.8	75.9	78.5	82.4	81.2	86.6	82.1
5-9	89.8	88.9	87.8	90.9	94.8	94.2	94.2	90.1	95.5	93.3	88.0
10-14	91.8	90.6	91.7	91.1	95.5	92.5	88.4	89.6	91.4	88.4	89.6
15-19	88.5	87.0	85.0	87.7	87.7	87.8	90.4	90.1	87.9	91.0	89.9
20-24	81.9	81.2	79.4	83.8	81.4	83.7	83.4	84.4	83.9	83.1	85.2
25-29	72.0	72.5	73.3	75.5	72.8	75.5	74.8	76.3	76.2	76.9	77.5
30-34	67.2	65.7	66.5	68.9	65.1	68.8	69.8	68.6	70.4	69.6	70.7
35-39	62.9	63.6	62.1	63.3	61.7	66.1	66.3	67.5	63.3	64.8	67.0
40-44	56.8	60.6	59.5	60.6	62.3	60.7	60.3	64.0	61.2	61.9	62.4
45-49	50.5	53.0	53.1	53.6	53.3	55.8	53.8	52.8	53.3	55.9	56.7
50-54	44.8	43.5	43.3	44.4	46.4	45.9	46.4	45.9	47.4	47.2	49.5
55-59	37.0	36.9	35.8	38.2	39.1	37.9	38.1	38.3	40.5	39.6	41.3
60-64	30.0	27.8	27.9	29.3	28.3	30.5	30.6	31.0	31.1	32.0	33.0
65-69	20.9	22.4	22.6	22.4	23.3	23.3	23.5	23.4	22.9	24.5	25.1
70-74	15.7	17.5	17.3	16.0	17.4	17.9	18.1	18.5	17.6	18.6	19.3
75-79	10.3	12.6	13.0	13.4	12.5	12.3	13.4	12.2	13.8	12.4	13.4
80-84	7.2	8.2	10.1	8.3	8.5	7.5	9.7	9.7	8.5	8.6	9.3
85+	4.3	5.1	3.2	5.4	4.8	4.0	6.6	6.1	5.1	5.2	6.1
White	37.9	38.0	37.5	37.2	35.9	35.2	34.8	34.0	33.0	33.0	33.4
Black	41.8	43.5	41.4	42.2	41.7	43.7	42.7	43.3	42.3	42.8	43.1
Other	46.2	43.0	41.5	44.5	45.7	46.9	51.4	49.3	49.6	46.9	48.2
Female	39.3	40.0	39.1	38.7	37.7	38.1	36.9	36.5	35.5	36.0	35.9
Male	39.5	39.8	38.6	39.4	38.5	38.6	38.8	38.4	37.5	37.3	38.5
Diabetes	27.1	26.3	26.2	28.1	26.9	28.0	28.0	27.2	27.4	27.8	29.3
Hypertension	36.0	36.0	35.3	34.0	32.3	33.3	33.0	33.2	31.6	32.2	32.8
Glomerulonephritis	57.3	56.8	54.3	55.3	54.6	56.0	55.6	55.4	54.3	55.4	57.7
Cystic Kidney	60.9	60.7	62.5	62.2	63.1	62.1	66.3	63.6	65.3	66.0	70.0
Other Urologic	43.7	46.2	42.0	40.5	40.2	37.2	39.2	39.0	42.0	38.9	45.7
Other Cause	47.3	47.4	44.9	45.2	45.5	45.0	44.2	44.9	43.7	42.7	42.8
Unknown Cause	41.6	40.3	39.7	37.7	36.9	34.4	38.1	39.6	37.1	36.6	41.5
Missing Cause	27.5	37.8	42.5	49.1	54.0	52.9	55.9	54.3	59.3	58.9	39.2
Total Crude Prob.	39.4	39.9	38.8	39.0	38.1	38.4	38.0	37.6	36.6	36.7	37.3

Table I.8 (continued)

STANDARD ERRORS											
	1983	1984	1985	1986	1987	1988	1989	1990	1991	1992	1993
0-4	4.2	4.4	3.7	3.6	3.8	4.0	3.7	3.8	3.4	3.1	3.4
5-9	3.2	3.1	3.4	2.7	2.3	2.5	2.3	2.7	2.0	2.6	3.3
10-14	2.0	2.0	2.0	2.1	1.5	1.9	2.4	2.2	1.9	2.3	2.2
15-19	1.6	1.7	1.8	1.6	1.6	1.6	1.3	1.4	1.6	1.4	1.4
20-24	1.5	1.4	1.5	1.4	1.4	1.3	1.3	1.3	1.3	1.2	1.2
25-29	1.4	1.3	1.3	1.2	1.3	1.2	1.1	1.1	1.1	1.1	1.1
30-34	1.3	1.3	1.2	1.1	1.2	1.1	1.0	1.0	1.0	1.0	1.0
35-39	1.3	1.3	1.2	1.2	1.1	1.0	1.0	1.0	1.0	0.9	0.9
40-44	1.3	1.3	1.2	1.2	1.1	1.1	1.0	0.9	0.9	0.9	0.9
45-49	1.2	1.2	1.2	1.1	1.1	1.0	1.0	0.9	0.9	0.8	0.8
50-54	1.1	1.1	1.1	1.0	1.0	0.9	0.9	0.9	0.8	0.8	0.8
55-59	1.0	0.9	0.9	0.9	0.9	0.8	0.8	0.8	0.7	0.7	0.7
60-64	0.8	0.8	0.7	0.7	0.7	0.7	0.7	0.6	0.6	0.6	0.6
65-69	0.7	0.7	0.7	0.7	0.6	0.6	0.6	0.5	0.5	0.5	0.5
70-74	0.7	0.8	0.7	0.6	0.6	0.6	0.5	0.5	0.5	0.5	0.5
75-79	0.8	0.8	0.8	0.7	0.6	0.6	0.6	0.5	0.5	0.5	0.5
80-84	1.0	1.0	1.0	0.9	0.8	0.7	0.7	0.7	0.6	0.5	0.5
85+	1.4	1.4	1.0	1.2	1.1	0.9	1.0	0.9	0.7	0.7	0.7
White	0.4	0.4	0.4	0.3	0.3	0.3	0.3	0.3	0.3	0.3	0.3
Black	0.6	0.6	0.5	0.5	0.5	0.5	0.4	0.4	0.4	0.4	0.4
Other	1.7	1.5	1.4	1.3	1.2	1.2	1.3	1.2	1.1	1.0	1.0
Female	0.5	0.5	0.4	0.4	0.4	0.4	0.4	0.3	0.3	0.3	0.3
Male	0.4	0.4	0.4	0.4	0.4	0.3	0.3	0.3	0.3	0.3	0.3
Diabetes	0.6	0.5	0.5	0.5	0.4	0.4	0.4	0.4	0.3	0.3	0.3
Hypertension	0.6	0.6	0.6	0.5	0.5	0.5	0.4	0.4	0.4	0.4	0.4
Glomerulonephritis	0.8	0.8	0.8	0.8	0.7	0.7	0.7	0.7	0.7	0.6	0.6
Cystic Kidney	1.5	1.5	1.5	1.4	1.4	1.4	1.4	1.3	1.3	1.2	1.2
Other Urologic	2.3	2.3	2.0	2.0	2.0	1.9	1.8	1.8	1.5	1.5	1.6
Other Cause	0.9	0.8	0.8	0.8	0.8	0.8	0.7	0.8	0.7	0.7	0.7
Unknown Cause	1.4	1.3	1.3	1.2	1.1	1.1	1.2	1.2	1.1	1.0	1.1
Missing Cause	0.8	1.2	1.3	1.2	1.2	1.0	1.1	1.0	1.0	1.0	0.6
Total Crude Prob.	0.3	0.3	0.3	0.3	0.3	0.3	0.2	0.2	0.2	0.2	0.2

Table I.9

Ten-year Survival Probabilities*all patients, from day 91 to ten years + 90 days, unadjusted*

PROBABILITIES	1978	1979	1980	1981	1982	1983	1984	1985	1986	1987	1988
0-4	68.0	64.0	75.0	71.9	76.9	76.3	70.1	73.5	81.3	78.9	67.0
5-9	76.7	73.8	81.2	81.3	78.2	81.8	85.9	81.1	85.5	82.3	91.9
10-14	74.9	74.9	77.9	77.0	77.0	84.2	82.1	82.9	85.0	88.4	83.7
15-19	73.3	76.6	79.4	80.0	79.7	80.4	78.5	75.8	78.9	78.8	78.5
20-24	66.8	72.7	69.3	69.1	71.1	71.0	70.5	68.7	71.2	70.8	74.7
25-29	59.4	57.1	60.7	57.7	62.1	58.2	58.7	61.5	61.6	60.0	62.6
30-34	49.2	52.2	49.0	51.6	50.3	52.8	51.4	49.1	53.6	49.4	52.0
35-39	39.6	40.4	44.8	42.7	45.4	44.5	45.8	44.7	45.0	45.3	46.9
40-44	35.8	36.1	36.4	36.7	37.7	37.6	38.9	40.5	40.1	39.8	39.6
45-49	27.5	28.6	28.3	29.1	29.8	29.6	32.1	30.7	32.5	31.7	32.4
50-54	20.4	21.4	20.6	23.5	22.8	21.4	21.4	21.7	22.3	23.2	22.0
55-59	14.7	13.9	11.7	13.4	13.8	12.7	13.5	14.5	14.7	16.3	14.5
60-64	7.6	8.2	8.2	8.6	6.7	8.7	8.3	9.2	8.9	8.3	9.1
65-69	4.9	4.9	4.5	5.0	4.0	4.8	5.2	6.1	5.0	4.7	5.1
70-74	3.0	2.3	2.9	2.3	3.4	2.8	3.3	3.3	2.7	3.0	2.5
75-79	2.6	1.7	1.5	1.7	1.9	1.1	2.1	1.8	1.7	1.9	1.8
80-84	0.0	0.0	1.5	0.7	1.8	0.3	1.3	0.9	0.8	0.9	0.5
85 +	0.0	0.0	1.2	0.0	0.0	0.0	0.8	0.7	0.3	0.0	0.2
White	27.3	26.5	25.8	26.2	25.7	22.9	23.1	22.4	22.0	20.8	19.6
Black	23.7	23.0	22.3	23.8	22.7	20.5	21.4	20.9	20.6	19.8	21.3
Other	25.9	26.3	26.6	29.3	32.0	28.7	27.3	25.8	27.6	26.7	28.3
Female	26.3	24.8	24.7	25.7	24.1	21.5	22.2	21.5	20.5	19.6	19.7
Male	25.8	25.7	24.5	25.4	25.7	23.0	23.3	22.7	23.0	21.8	21.3
Diabetes	11.3	11.9	11.9	11.7	11.3	11.1	11.3	11.6	11.9	11.1	10.8
Hypertension	23.2	23.1	21.6	19.2	18.1	17.0	16.9	15.9	15.8	13.7	14.5
Glomerulonephritis	47.0	44.1	44.3	40.5	40.7	38.4	38.5	37.4	38.4	36.8	37.5
Cystic Kidney	39.0	39.6	36.2	38.3	36.1	37.8	38.1	41.6	39.0	40.6	41.9
Other Urologic	25.2	26.1	28.8	22.3	26.2	25.5	24.7	25.7	23.0	21.5	21.1
Other Cause	37.6	41.4	38.4	33.7	31.8	30.9	31.3	28.9	28.7	30.2	28.8
Unknown Cause	25.9	24.6	24.9	24.1	25.3	22.3	21.3	22.6	20.5	19.5	19.1
Missing Cause	20.0	18.8	18.8	26.5	32.6	17.2	26.2	29.7	33.1	35.1	32.9
Total Crude Prob.	26.0	25.3	24.6	25.5	25.0	22.3	22.8	22.1	21.9	20.8	20.6

Table I.9 (continued)

STANDARD ERRORS											
	1978	1979	1980	1981	1982	1983	1984	1985	1986	1987	1988
0-4	9.1	9.3	6.7	5.6	4.4	4.4	4.6	4.1	3.7	3.9	4.4
5-9	4.9	4.8	4.2	4.1	4.1	4.1	3.5	4.1	3.3	3.9	2.9
10-14	3.1	3.3	3.2	3.1	2.9	2.6	2.6	2.7	2.7	2.3	2.7
15-19	2.2	2.0	2.0	1.9	1.9	1.9	2.1	2.2	2.0	2.0	2.0
20-24	1.9	1.8	1.9	1.8	1.7	1.8	1.7	1.7	1.7	1.6	1.5
25-29	1.7	1.7	1.7	1.6	1.5	1.5	1.5	1.4	1.4	1.4	1.4
30-34	1.8	1.6	1.5	1.4	1.4	1.4	1.3	1.3	1.2	1.3	1.2
35-39	1.6	1.6	1.6	1.5	1.4	1.4	1.3	1.2	1.2	1.2	1.1
40-44	1.5	1.5	1.5	1.4	1.3	1.3	1.3	1.2	1.2	1.1	1.1
45-49	1.3	1.3	1.3	1.2	1.2	1.1	1.1	1.1	1.1	1.0	1.0
50-54	1.1	1.0	1.0	1.0	0.9	0.9	0.9	0.9	0.9	0.8	0.8
55-59	0.9	0.8	0.7	0.7	0.7	0.7	0.7	0.7	0.6	0.7	0.6
60-64	0.7	0.7	0.6	0.6	0.5	0.5	0.5	0.5	0.5	0.4	0.4
65-69	0.6	0.5	0.5	0.5	0.4	0.4	0.4	0.4	0.3	0.3	0.3
70-74	0.5	0.4	0.5	0.4	0.4	0.3	0.4	0.3	0.3	0.3	0.2
75-79	0.7	0.5	0.4	0.4	0.4	0.3	0.3	0.3	0.3	0.3	0.2
80-84	0.0	0.0	0.7	0.4	0.6	0.2	0.4	0.3	0.3	0.3	0.2
85+	0.0	0.0	0.9	0.0	0.0	0.0	0.5	0.4	0.2	0.0	0.2
White	0.5	0.5	0.4	0.4	0.4	0.3	0.3	0.3	0.3	0.3	0.3
Black	0.6	0.6	0.6	0.5	0.5	0.5	0.5	0.4	0.4	0.4	0.4
Other	2.0	1.8	1.8	1.7	1.8	1.5	1.4	1.3	1.2	1.1	1.1
Female	0.6	0.5	0.5	0.5	0.4	0.4	0.4	0.4	0.3	0.3	0.3
Male	0.5	0.5	0.5	0.4	0.4	0.4	0.4	0.3	0.3	0.3	0.3
Diabetes	0.9	0.8	0.7	0.5	0.5	0.4	0.4	0.4	0.3	0.3	0.3
Hypertension	1.0	0.9	0.8	0.6	0.5	0.5	0.5	0.4	0.4	0.4	0.4
Glomerulonephritis	1.2	1.1	1.1	0.9	0.8	0.8	0.8	0.7	0.7	0.7	0.7
Cystic Kidney	2.1	2.0	1.9	1.7	1.5	1.5	1.5	1.5	1.4	1.4	1.4
Other Urologic	3.0	2.9	2.8	2.1	2.2	2.0	2.0	1.8	1.7	1.7	1.6
Other Cause	1.4	1.3	1.2	1.0	0.9	0.8	0.8	0.7	0.7	0.7	0.7
Unknown Cause	1.3	1.2	1.2	1.1	1.2	1.2	1.1	1.1	1.0	0.9	0.9
Missing Cause	0.5	0.5	0.5	0.8	1.2	0.7	1.1	1.2	1.2	1.1	1.0
Total Crude Prob.	0.4	0.4	0.3	0.3	0.3	0.3	0.3	0.3	0.2	0.2	0.2

Table I.10

One-year Survival Probabilities*all patients, from day 91 to one year + 90 days, adjusted for age, gender, race, & primary diagnosis*

PROBABILITIES	1987	1988	1989	1990	1991	1992	1993	1994	1995	1996	1997
0-19	96.5	95.5	95.8	96.6	95.1	95.8	95.7	97.2	98.0	97.4	97.8
20-44	91.0	91.7	91.3	92.3	91.6	91.9	91.6	92.1	92.7	93.6	94.5
45-64	82.2	82.5	83.4	84.3	84.5	85.2	85.7	85.9	87.1	87.7	88.5
65-74	69.1	69.6	70.7	72.5	72.2	72.8	74.5	74.0	75.1	75.6	74.7
75 +	57.3	58.2	61.1	62.3	61.9	61.9	63.3	64.3	64.1	64.2	63.4
White	74.1	75.1	76.3	77.8	77.3	77.8	78.8	79.5	80.4	80.5	80.9
Black	83.9	83.9	83.7	84.9	84.7	85.4	85.3	85.0	85.9	87.3	87.8
Other	82.4	84.6	87.8	89.2	89.7	88.5	90.4	88.7	89.1	89.4	90.5
Male	77.8	78.2	79.4	80.9	80.6	80.9	82.1	82.1	83.0	83.6	84.3
Female	78.4	79.5	80.0	81.2	80.9	81.4	81.6	82.1	82.9	83.6	83.9
Diabetes	71.9	73.7	74.9	75.7	76.2	76.8	78.8	78.8	79.6	80.5	80.9
Hypertension	74.9	75.2	76.1	78.0	77.1	77.4	78.3	78.3	80.6	80.6	80.6
Glomerulonephritis	88.5	88.6	89.2	90.0	88.5	89.8	90.8	90.5	91.4	92.2	92.0
Other Cause	84.6	84.2	84.2	85.8	85.1	85.3	85.0	85.6	85.8	86.6	87.8
Total Adjusted Prob.	78.1	78.8	79.7	81.0	80.7	81.2	81.9	82.1	83.0	83.6	84.1
STANDARD ERRORS											
0-19	0.6	0.7	0.7	0.6	0.7	0.7	0.7	0.5	0.4	0.5	0.5
20-44	0.3	0.3	0.3	0.3	0.3	0.3	0.3	0.3	0.2	0.2	0.2
45-64	0.3	0.3	0.3	0.3	0.3	0.3	0.3	0.2	0.2	0.2	0.2
65-74	0.5	0.5	0.4	0.4	0.4	0.4	0.4	0.3	0.3	0.3	0.3
75 +	0.8	0.7	0.6	0.6	0.6	0.5	0.5	0.5	0.4	0.4	0.4
White	0.3	0.3	0.3	0.2	0.2	0.2	0.2	0.2	0.2	0.2	0.2
Black	0.4	0.3	0.3	0.3	0.3	0.3	0.3	0.3	0.2	0.2	0.2
Other	0.9	0.8	0.8	0.7	0.6	0.6	0.5	0.5	0.4	0.4	0.4
Male	0.3	0.3	0.3	0.3	0.2	0.2	0.2	0.2	0.2	0.2	0.2
Female	0.3	0.3	0.3	0.3	0.3	0.2	0.2	0.2	0.2	0.2	0.2
Diabetes	0.5	0.4	0.4	0.4	0.3	0.3	0.3	0.3	0.3	0.2	0.2
Hypertension	0.5	0.4	0.4	0.4	0.4	0.3	0.3	0.3	0.3	0.3	0.3
Glomerulonephritis	0.5	0.4	0.4	0.4	0.4	0.4	0.4	0.4	0.3	0.3	0.4
Other Cause	0.4	0.3	0.3	0.3	0.3	0.3	0.3	0.3	0.3	0.3	0.3
Total Adjusted Prob.	0.2	0.2	0.2	0.2	0.2	0.2	0.2	0.2	0.1	0.1	0.1

Table I.11

Two-year Survival Probabilities*all patients, from day 91 to two years + 90 days, adjusted for age, gender, race, & primary diagnosis*

PROBABILITIES	1986	1987	1988	1989	1990	1991	1992	1993	1994	1995	1996
0-19	93.6	94.8	93.1	93.8	93.9	93.4	94.6	93.4	95.3	96.4	95.3
20-44	83.6	83.4	84.8	84.6	85.7	84.5	85.1	85.2	85.8	87.1	88.3
45-64	67.3	67.4	68.3	70.0	70.7	71.2	72.0	72.9	73.9	75.6	76.8
65-74	49.0	49.8	50.9	51.9	52.4	53.0	54.4	55.4	55.4	57.4	57.9
75 +	37.8	35.3	36.1	40.2	40.1	39.7	40.5	41.7	42.5	42.7	43.7
White	56.5	56.8	57.9	60.0	60.6	60.4	61.5	62.3	63.5	64.9	65.5
Black	70.5	69.7	71.2	71.1	72.1	71.9	73.2	73.5	73.6	75.3	77.2
Other	69.5	68.9	72.0	77.1	78.1	80.8	78.8	80.6	78.7	79.8	80.9
Male	62.0	61.3	63.0	64.6	65.5	65.5	66.4	67.8	68.1	70.0	70.7
Female	62.5	63.0	64.0	65.3	66.0	65.9	67.0	66.9	67.9	69.1	70.7
Diabetes	50.4	51.3	53.5	55.8	55.9	57.1	58.2	60.5	60.8	62.7	64.2
Hypertension	58.9	57.4	58.8	59.9	60.8	60.4	61.3	62.4	62.9	66.0	66.4
Glomerulonephritis	78.3	78.4	79.8	79.8	81.0	79.1	80.8	82.2	82.5	83.9	85.0
Other Cause	74.0	74.0	73.4	74.2	75.7	74.8	75.4	74.4	75.6	76.1	77.8
Total Adjusted Prob.	62.2	62.1	63.5	64.9	65.7	65.7	66.7	67.4	68.1	69.6	70.7
STANDARD ERRORS											
0-19	0.9	0.8	0.9	0.8	0.8	0.8	0.8	0.8	0.7	0.6	0.6
20-44	0.4	0.4	0.4	0.4	0.4	0.4	0.4	0.4	0.3	0.3	0.3
45-64	0.5	0.4	0.4	0.4	0.4	0.3	0.3	0.3	0.3	0.3	0.3
65-74	0.6	0.6	0.5	0.5	0.5	0.4	0.4	0.4	0.4	0.4	0.4
75 +	0.9	0.8	0.7	0.7	0.6	0.6	0.5	0.5	0.5	0.5	0.4
White	0.4	0.4	0.3	0.3	0.3	0.3	0.3	0.3	0.3	0.3	0.3
Black	0.5	0.5	0.4	0.4	0.4	0.4	0.3	0.3	0.3	0.3	0.3
Other	1.3	1.2	1.1	1.1	1	0.9	0.8	0.8	0.7	0.6	0.6
Male	0.4	0.4	0.4	0.3	0.3	0.3	0.3	0.3	0.3	0.3	0.3
Female	0.4	0.4	0.4	0.4	0.3	0.3	0.3	0.3	0.3	0.3	0.3
Diabetes	0.6	0.5	0.5	0.4	0.4	0.4	0.4	0.4	0.3	0.3	0.3
Hypertension	0.6	0.5	0.5	0.5	0.5	0.4	0.4	0.4	0.4	0.4	0.4
Glomerulonephritis	0.6	0.6	0.6	0.6	0.5	0.5	0.5	0.5	0.5	0.5	0.5
Other Cause	0.5	0.5	0.4	0.4	0.4	0.4	0.4	0.4	0.4	0.3	0.3
Total Adjusted Prob.	0.3	0.3	0.3	0.2	0.2	0.2	0.2	0.2	0.2	0.2	0.2

Table I.12

Five-year Survival Probabilities*all patients, from day 91 to five years + 90 days, adjusted for age, gender, race, & primary diagnosis*

PROBABILITIES	1983	1984	1985	1986	1987	1988	1989	1990	1991	1992	1993
0-19	88.6	86.6	86.3	88.4	89.4	87.8	88.6	88.9	88.6	90.0	88.5
20-44	66.2	67.7	67.1	69.1	67.9	69.9	70.3	71.8	70.2	70.5	71.5
45-64	35.7	35.3	35.3	38.0	38.5	39.5	40.2	40.4	42.0	42.3	43.6
65-74	14.9	16.6	17.3	16.5	17.6	18.0	18.8	18.9	18.9	20.2	20.6
75 +	6.5	8.8	9.6	9.0	8.4	7.9	10.3	9.3	9.5	9.1	10.2
White	22.5	24.2	25.0	26.1	26.2	26.8	28.3	28.1	28.6	29.4	30.4
Black	38.0	39.7	38.5	40.5	40.4	42.3	41.9	43.3	42.9	43.7	44.3
Other	38.1	35.1	37.2	40.2	43.2	45.7	52.0	50.7	51.8	49.9	52.2
Male	28.1	29.5	29.9	32.0	31.7	33.0	35.0	35.0	35.4	35.5	37.2
Female	29.0	30.5	30.7	31.7	32.3	33.1	33.4	33.9	34.3	35.2	35.3
Diabetes	16.9	17.1	17.8	19.4	19.5	20.8	21.5	21.7	22.8	23.6	25.6
Hypertension	24.6	25.4	26.1	25.7	25.7	26.6	27.5	28.2	27.4	27.8	29.0
Glomerulonephritis	54.1	54.1	52.7	54.7	54.1	56.3	57.3	57.2	56.9	57.3	59.9
Other Cause	45.3	48.6	48.3	50.1	51.4	50.0	51.7	52.2	52.2	52.4	50.6
Total Adjusted Prob.	28.6	30.0	30.3	31.9	32.0	33.0	34.3	34.5	34.9	35.4	36.3
STANDARD ERRORS											
0-19	1.1	1.2	1.2	1.1	1.1	1.2	1.1	1.1	1.1	1	1.1
20-44	0.7	0.6	0.6	0.6	0.6	0.5	0.5	0.5	0.5	0.5	0.5
45-64	0.5	0.5	0.5	0.5	0.5	0.4	0.4	0.4	0.4	0.4	0.4
65-74	0.6	0.6	0.5	0.5	0.4	0.4	0.4	0.4	0.3	0.3	0.3
75 +	0.6	0.6	0.6	0.5	0.4	0.4	0.4	0.4	0.3	0.3	0.3
White	0.4	0.4	0.4	0.4	0.3	0.3	0.3	0.3	0.3	0.3	0.3
Black	0.6	0.6	0.6	0.5	0.5	0.5	0.5	0.4	0.4	0.4	0.4
Other	1.8	1.6	1.5	1.4	1.4	1.3	1.4	1.3	1.2	1.1	1.1
Male	0.5	0.4	0.4	0.4	0.4	0.4	0.4	0.3	0.3	0.3	0.3
Female	0.5	0.5	0.5	0.4	0.4	0.4	0.4	0.4	0.3	0.3	0.3
Diabetes	0.6	0.5	0.5	0.5	0.4	0.4	0.4	0.3	0.3	0.3	0.3
Hypertension	0.6	0.6	0.6	0.5	0.5	0.5	0.4	0.4	0.4	0.4	0.4
Glomerulonephritis	0.9	0.9	0.8	0.8	0.8	0.8	0.8	0.7	0.7	0.7	0.7
Other Cause	0.6	0.6	0.6	0.6	0.6	0.5	0.5	0.5	0.5	0.5	0.5
Total Adjusted Prob.	0.3	0.3	0.3	0.3	0.3	0.3	0.3	0.2	0.2	0.2	0.2

Table I.13

Ten-year Survival Probabilities*all patients, from day 91 to ten years + 90 days, adjusted for age, gender, race, & primary diagnosis*

PROBABILITIES	1978	1979	1980	1981	1982	1983	1984	1985	1986	1987	1988
0-19	74.8	74.2	79.1	79.0	78.7	81.3	79.1	77.8	81.6	81.5	79.3
20-44	40.4	42.5	44.1	45.0	48.0	48.5	49.8	49.8	50.9	50.0	51.7
45-64	10.0	9.9	10.7	11.3	11.5	12.3	12.5	13.2	14.5	14.9	14.8
65-74	2.6	1.6	2.2	2.0	2.2	2.3	2.7	3.4	2.6	2.6	2.5
75 +	0.7	0.7	0.8	0.6	0.9	0.4	1.2	1.0	0.8	0.8	0.6
White	4.5	5.1	5.3	5.5	5.9	6.1	7.4	8.0	8.0	8.1	7.7
Black	12.5	11.6	12.4	13.2	13.2	13.2	14.0	14.6	14.9	14.7	15.7
Other	6.2	5.9	9.4	12.7	14.9	14.3	14.7	16.3	16.6	17.2	19.7
Male	6.6	7.1	7.5	8.1	8.9	8.8	9.9	11.1	11.6	11.2	11.1
Female	7.5	7.0	7.7	8.1	8.4	8.5	9.7	9.9	9.6	10.2	10.3
Diabetes	4.1	3.2	3.8	3.7	3.7	3.6	4.2	4.8	4.5	4.7	4.5
Hypertension	4.7	4.6	5.5	4.6	5.2	5.4	5.6	5.7	5.7	5.6	5.5
Glomerulonephritis	25.4	26.2	26.2	26.2	26.3	27.2	28.6	28.9	30.6	29.2	30.0
Other Cause	18.7	19.4	19.4	22.1	23.9	22.8	25.8	26.6	26.2	27.7	26.9
Total Adjusted Prob.	7.0	7.1	7.6	8.1	8.7	8.7	9.9	10.5	10.7	10.7	10.7
STANDARD ERRORS											
0-19	1.8	1.7	1.6	1.5	1.4	1.4	1.4	1.5	1.4	1.4	1.4
20-44	0.9	0.8	0.8	0.7	0.7	0.7	0.7	0.7	0.6	0.6	0.6
45-64	0.6	0.5	0.5	0.4	0.4	0.4	0.4	0.4	0.4	0.3	0.3
65-74	0.4	0.3	0.3	0.2	0.2	0.2	0.2	0.2	0.2	0.2	0.2
75 +	0.3	0.3	0.3	0.2	0.2	0.1	0.2	0.2	0.1	0.1	0.1
White	0.3	0.3	0.3	0.2	0.2	0.2	0.2	0.2	0.2	0.2	0.2
Black	0.7	0.6	0.5	0.5	0.5	0.4	0.4	0.4	0.4	0.4	0.4
Other	1.4	1.3	1.5	1.5	1.6	1.4	1.2	1.2	1.2	1.1	1.1
Male	0.4	0.4	0.3	0.3	0.3	0.3	0.3	0.3	0.3	0.3	0.3
Female	0.5	0.4	0.4	0.4	0.3	0.3	0.3	0.3	0.3	0.3	0.3
Diabetes	0.6	0.5	0.4	0.3	0.3	0.3	0.3	0.3	0.2	0.2	0.2
Hypertension	0.6	0.5	0.5	0.4	0.3	0.3	0.3	0.3	0.3	0.3	0.2
Glomerulonephritis	1.3	1.2	1.2	1.0	0.9	0.8	0.8	0.8	0.8	0.8	0.8
Other Cause	0.5	0.4	0.4	0.5	0.6	0.5	0.6	0.6	0.5	0.5	0.5
Total Adjusted Prob.	0.3	0.3	0.3	0.2	0.2	0.2	0.2	0.2	0.2	0.2	0.2

Table I.14

One-year Survival Probabilities*patients aged 65 & over, from day 1 to one year, unadjusted*

	PROBABILITIES					STANDARD ERRORS				
	1988-89	1990-91	1992-93	1994-95	1996-97	1988-89	1990-91	1992-93	1994-95	1996-97
65-69	73.1	73.8	75.2	75.8	75.1	0.4	0.4	0.3	0.3	0.3
70-74	66.7	68.2	70.1	70.3	69.4	0.5	0.4	0.4	0.3	0.3
75-79	61.2	63.6	63.5	64.6	63.6	0.6	0.5	0.4	0.4	0.4
80-84	57.4	57.6	59.1	59.0	58.2	0.8	0.7	0.6	0.5	0.5
85 +	46.8	51.1	50.9	52.2	48.5	1.3	1.1	1.0	0.9	0.7
White	63.0	64.3	65.0	65.6	64.3	0.3	0.3	0.2	0.2	0.2
Black	73.2	73.9	74.8	74.3	72.3	0.5	0.4	0.4	0.4	0.3
Other	72.4	76.8	77.7	75.6	74.0	1.3	1.1	0.9	0.7	0.7
Female	67.7	68.4	68.9	69.2	67.5	0.4	0.3	0.3	0.3	0.2
Male	64.2	65.8	67.1	67.2	66.0	0.4	0.3	0.3	0.3	0.2
Diabetes	66.8	68.0	69.7	69.8	68.9	0.5	0.4	0.3	0.3	0.3
Hypertension	64.4	65.4	65.4	66.7	65.8	0.4	0.4	0.3	0.3	0.3
Glomerulonephritis	72.4	72.5	74.7	73.9	73.9	0.8	0.7	0.7	0.6	0.6
Cystic Kidney	82.1	83.4	83.7	84.4	83.4	1.6	1.4	1.3	1.2	1.2
Other Urologic	62.1	68.6	68.6	66.3	69.3	1.6	1.4	1.4	1.3	1.2
Other Cause	62.1	61.9	63.4	61.1	56.4	0.8	0.8	0.8	0.7	0.6
Unknown Cause	62.2	62.8	62.3	63.6	61.0	1.1	1.0	1.0	0.9	0.8
Missing Cause	69.9	75.1	71.5	70.7	57.5	1.4	1.2	0.7	0.9	1.3
Total Crude Prob.	65.9	67.1	68.0	68.2	66.7	0.3	0.2	0.2	0.2	0.2

Table I.15

Two-year Survival Probabilities*patients aged 65 & over, from day 1 to two years, unadjusted*

	PROBABILITIES						STANDARD ERRORS					
	1986-87	1988-89	1990-91	1992-93	1994-95	1996-97	1986-87	1988-89	1990-91	1992-93	1994-95	1996-97
65-69	53.6	55.4	56.0	58.1	59.0	59.1	0.5	0.5	0.4	0.4	0.4	0.4
70-74	47.2	48.0	49.2	50.9	51.9	51.8	0.6	0.5	0.4	0.4	0.4	0.4
75-79	40.5	41.7	43.5	44.4	45.2	45.1	0.6	0.6	0.5	0.4	0.4	0.4
80-84	33.9	35.5	36.4	38.0	38.9	38.0	0.9	0.8	0.7	0.6	0.5	0.5
85 +	26.6	28.2	28.2	29.2	30.5	29.2	1.4	1.2	1.0	0.9	0.8	0.7
White	42.7	43.8	44.4	45.4	46.4	45.7	0.4	0.3	0.3	0.3	0.2	0.2
Black	55.0	55.5	56.2	57.8	57.4	56.2	0.6	0.5	0.5	0.4	0.4	0.4
Other	48.5	54.3	60.7	61.8	58.9	58.2	1.5	1.4	1.2	1.0	0.8	0.8
Female	48.3	49.0	49.1	50.5	50.8	49.7	0.4	0.4	0.3	0.3	0.3	0.3
Male	43.8	45.2	46.6	47.8	48.6	48.1	0.4	0.4	0.3	0.3	0.3	0.3
Diabetes	42.4	45.0	46.4	49.2	49.5	49.3	0.6	0.5	0.4	0.4	0.3	0.3
Hypertension	45.1	46.1	46.4	47.0	48.7	48.5	0.5	0.4	0.4	0.3	0.3	0.3
Glomerulonephritis	53.7	54.4	55.2	57.3	56.6	57.3	0.9	0.8	0.8	0.8	0.7	0.7
Cystic Kidney	64.3	66.9	67.5	70.6	70.1	71.4	2.0	1.9	1.7	1.7	1.5	1.6
Other Urologic	49.0	45.6	52.1	51.9	48.6	51.9	1.8	1.6	1.5	1.5	1.4	1.4
Other Cause	46.1	44.7	45.1	45.7	44.9	40.8	0.9	0.8	0.8	0.8	0.7	0.7
Unknown Cause	48.0	43.5	44.4	46.0	45.2	42.9	1.2	1.1	1.0	1.0	1.0	0.9
Missing Cause	39.1	58.0	58.6	52.2	54.3	45.8	1.5	1.5	1.4	0.8	1.0	1.4
Total Crude Prob.	46.0	47.0	47.8	49.1	49.7	48.9	0.3	0.3	0.2	0.2	0.2	0.2

Table I.16

Five-year Survival Probabilities*patients aged 65 & over, from day 1 to five years, unadjusted*

	PROBABILITIES					STANDARD ERRORS				
	1984-85	1986-87	1988-89	1990-91	1992-93	1984-85	1986-87	1988-89	1990-91	1992-93
65-69	21.9	22.4	23.2	23.1	24.6	0.5	0.4	0.4	0.3	0.3
70-74	16.8	16.5	17.5	17.6	18.7	0.5	0.4	0.4	0.3	0.3
75-79	12.0	12.6	12.5	12.7	12.7	0.5	0.4	0.4	0.3	0.3
80-84	8.7	7.9	8.4	8.8	8.7	0.7	0.5	0.5	0.4	0.3
85 +	3.9	4.6	5.2	5.1	5.4	0.8	0.7	0.6	0.5	0.4
White	14.8	14.4	14.6	14.2	14.8	0.3	0.3	0.2	0.2	0.2
Black	21.4	22.1	22.5	23.3	23.5	0.6	0.5	0.5	0.4	0.4
Other	16.3	19.1	23.2	25.1	24.8	1.4	1.2	1.2	1.1	0.9
Female	18.7	18.2	17.9	17.6	18.6	0.4	0.3	0.3	0.3	0.2
Male	14.6	14.9	16.0	16.1	16.1	0.3	0.3	0.3	0.2	0.2
Diabetes	12.0	12.3	13.0	13.6	14.9	0.5	0.4	0.3	0.3	0.3
Hypertension	16.3	16.0	16.3	16.1	16.2	0.4	0.4	0.3	0.3	0.3
Glomerulonephritis	21.1	21.5	22.8	22.7	23.9	0.8	0.8	0.7	0.7	0.7
Cystic Kidney	33.1	29.4	30.9	32.7	38.7	2.2	1.9	1.9	1.7	1.8
Other Urologic	23.3	22.2	21.5	22.0	22.5	1.7	1.5	1.3	1.2	1.2
Other Cause	17.5	17.4	17.7	17.6	18.6	0.7	0.7	0.6	0.6	0.6
Unknown Cause	19.7	16.0	15.5	16.8	15.9	1.1	0.8	0.8	0.8	0.7
Missing Cause	10.0	19.8	28.8	28.0	22.0	0.8	1.2	1.4	1.3	0.7
Total Crude Prob.	16.5	16.5	16.9	16.8	17.4	0.3	0.2	0.2	0.2	0.2

Table I.17

Ten-year Survival Probabilities*patients aged 65 & over, from day 1 to ten years, unadjusted*

	PROBABILITIES						STANDARD ERRORS					
	1978-79	1980-81	1982-83	1984-85	1986-87	1988-89	1978-79	1980-81	1982-83	1984-85	1986-87	1988-89
65-69	4.9	4.8	4.4	5.7	4.9	4.9	0.4	0.3	0.3	0.3	0.2	0.2
70-74	2.7	2.7	2.9	3.1	2.8	2.5	0.3	0.3	0.2	0.2	0.2	0.2
75-79	2.1	1.5	1.4	1.8	1.8	1.4	0.4	0.3	0.2	0.2	0.2	0.1
80-84	0.0	0.9	0.8	1.0	0.8	0.6	0.0	0.3	0.3	0.2	0.2	0.1
85 +	0.0	0.5	0.0	0.6	0.1	0.4	0.0	0.4	0.0	0.3	0.1	0.2
White	3.1	2.9	2.6	3.2	2.7	2.2	0.2	0.2	0.1	0.1	0.1	0.1
Black	4.2	4.0	3.8	4.3	3.7	4.1	0.4	0.4	0.3	0.3	0.2	0.2
Other	1.0	1.8	4.5	3.4	4.6	5.0	0.5	0.7	1.0	0.7	0.6	0.7
Female	4.3	3.7	3.7	3.8	3.1	3.1	0.3	0.3	0.2	0.2	0.2	0.1
Male	2.6	2.7	2.3	3.2	3.0	2.5	0.2	0.2	0.2	0.2	0.1	0.1
Diabetes	1.9	1.9	1.9	2.2	1.7	1.6	0.6	0.4	0.3	0.2	0.2	0.1
Hypertension	3.5	2.8	2.6	3.0	2.6	2.3	0.5	0.3	0.2	0.2	0.2	0.1
Glomerulonephritis	5.3	3.7	4.4	4.6	4.9	4.8	0.9	0.6	0.5	0.4	0.4	0.4
Cystic Kidney	7.5	8.5	7.8	7.3	5.8	6.5	2.0	1.7	1.3	1.2	1.0	1.0
Other Urologic	4.8	3.8	4.1	6.3	4.9	4.2	1.5	1.0	0.9	1.0	0.8	0.7
Other Cause	2.7	3.5	3.0	4.5	3.8	3.7	0.7	0.6	0.4	0.4	0.3	0.3
Unknown Cause	3.7	3.6	3.8	4.4	2.9	2.6	0.6	0.5	0.5	0.6	0.4	0.4
Missing Cause	2.9	3.1	2.0	2.8	5.5	7.1	0.3	0.3	0.3	0.4	0.7	0.8
Total Crude Prob.	3.3	3.2	2.9	3.5	3.0	2.8	0.2	0.2	0.1	0.1	0.1	0.1

Table I.18

One-year Survival Probabilities*patients aged 65 & over, from day 1 to one year, adjusted for age, gender, race, & primary diagnosis*

	PROBABILITIES					STANDARD ERRORS				
	1988-89	1990-91	1992-93	1994-95	1996-97	1988-89	1990-91	1992-93	1994-95	1996-97
65-74	69.3	70.8	72.5	73.2	72.6	0.3	0.3	0.2	0.2	0.2
75 +	57.3	59.7	60.2	61.0	60.0	0.5	0.4	0.3	0.3	0.3
White	61.1	63.0	64.2	65.4	64.7	0.3	0.3	0.2	0.2	0.2
Black	72.0	73.2	74.5	74.3	73.1	0.5	0.4	0.4	0.4	0.3
Other	71.8	75.9	77.4	75.5	74.7	1.3	1.1	0.9	0.7	0.7
Male	62.6	64.9	66.4	67.2	66.6	0.4	0.3	0.3	0.3	0.2
Female	66.2	67.6	68.5	69.2	68.2	0.4	0.3	0.3	0.3	0.2
Primary diagnosis										
Hypertension	65.6	67.2	69.3	69.8	69.5	0.5	0.4	0.4	0.3	0.3
Glomerulonephritis	62.9	64.5	64.9	66.6	66.5	0.4	0.4	0.3	0.3	0.3
Other Cause	71.4	72.1	74.4	74.2	74.7	0.8	0.7	0.7	0.6	0.6
Total Adjusted Prob.	63.3	65.4	66.6	65.5	61.4	0.5	0.5	0.4	0.4	0.4
	64.3	66.2	67.4	68.2	67.4	0.3	0.2	0.2	0.2	0.2

Table I.19

Two-year Survival Probabilities*patients aged 65 & over, from day 1 to two years, adjusted for age, gender, race, & primary diagnosis*

	PROBABILITIES						STANDARD ERRORS					
	1986-87	1988-89	1990-91	1992-93	1994-95	1996-97	1986-87	1988-89	1990-91	1992-93	1994-95	1996-97
65-74	48.5	50.6	52.0	54.1	55.4	55.6	0.4	0.3	0.3	0.3	0.3	0.3
75 +	35.0	36.7	38.4	39.8	40.8	40.5	0.5	0.4	0.4	0.3	0.3	0.3
White	39.1	41.0	42.3	44.0	45.8	45.8	0.4	0.3	0.3	0.3	0.2	0.2
Black	52.5	53.4	54.8	57.0	57.0	56.7	0.6	0.6	0.5	0.4	0.4	0.4
Other	45.4	53.0	59.1	61.1	58.4	58.7	1.6	1.5	1.3	1.1	0.9	0.8
Male	40.7	42.7	44.8	46.4	48.1	48.4	0.4	0.4	0.3	0.3	0.3	0.3
Female	44.8	46.6	47.6	49.5	50.3	50.1	0.5	0.4	0.3	0.3	0.3	0.3
Diabetes	40.1	43.1	44.9	48.1	49.1	49.6	0.6	0.5	0.4	0.4	0.3	0.3
Hypertension	42.1	43.8	44.7	45.8	48.1	48.9	0.5	0.4	0.4	0.3	0.3	0.4
Glomerulonephritis	51.0	52.7	54.1	56.4	56.6	58.0	1.0	0.9	0.8	0.8	0.7	0.7
Other Cause	45.2	45.9	47.6	48.4	48.3	45.4	0.6	0.6	0.5	0.5	0.4	0.5
Total Adjusted Prob.	42.6	44.6	46.2	47.9	49.2	49.2	0.3	0.3	0.2	0.2	0.2	0.2

Table I.20

Five-year Survival Probabilities*patients aged 65 & over, from day 1 to five years, adjusted for age, gender, race, & primary diagnosis*

	PROBABILITIES					STANDARD ERRORS				
	1984-85	1986-87	1988-89	1990-91	1992-93	1984-85	1986-87	1988-89	1990-91	1992-93
65-74	16.8	17.1	18.5	19.0	20.4	0.4	0.3	0.3	0.2	0.2
75 +	8.7	8.5	9.0	9.3	9.5	0.4	0.3	0.3	0.2	0.2
White	11.2	11.0	11.7	11.9	12.8	0.3	0.2	0.2	0.2	0.2
Black	18.4	19.1	19.5	21.0	21.5	0.6	0.5	0.5	0.4	0.4
Other	14.4	15.2	20.7	22.2	22.6	1.4	1.2	1.2	1.1	0.9
Male	11.6	11.8	13.1	13.6	13.9	0.3	0.3	0.3	0.2	0.2
Female	14.5	14.2	14.7	15.0	16.5	0.4	0.3	0.3	0.3	0.2
Diabetes	10.2	10.1	10.8	11.6	13.0	0.5	0.4	0.3	0.3	0.3
Hypertension	12.4	12.6	13.3	13.6	14.0	0.4	0.3	0.3	0.3	0.2
Glomerulonephritis	17.4	17.7	19.7	20.2	21.8	0.8	0.7	0.7	0.6	0.6
Other Cause	15.6	16.1	17.4	17.9	18.3	0.5	0.4	0.4	0.4	0.4
Total Adjusted Prob.	12.9	12.9	13.8	14.3	15.1	0.3	0.2	0.2	0.2	0.2

Table I.21

Ten-year Survival Probabilities*patients aged 65 & over, from day 1 to ten years, adjusted for age, gender, race, & primary diagnosis*

	PROBABILITIES						STANDARD ERRORS					
	1978-79	1980-81	1982-83	1984-85	1986-87	1988-89	1978-79	1980-81	1982-83	1984-85	1986-87	1988-89
65-74	2.2	2.3	2.3	3.2	2.7	2.7	0.2	0.2	0.1	0.2	0.1	0.1
75 +	0.8	0.7	0.6	1.0	0.8	0.6	0.2	0.1	0.1	0.1	0.1	0.1
White	1.2	1.2	1.0	1.7	1.4	1.1	0.2	0.1	0.1	0.1	0.1	0.1
Black	1.9	1.9	2.0	2.8	2.4	2.6	0.3	0.3	0.2	0.2	0.2	0.2
Other	0.1	0.9	3.0	2.4	2.5	3.4	0.1	0.5	0.9	0.6	0.5	0.5
Male	0.9	1.2	1.0	1.9	1.6	1.3	0.1	0.1	0.1	0.1	0.1	0.1
Female	1.8	1.4	1.6	2.0	1.5	1.7	0.3	0.2	0.1	0.1	0.1	0.1
Diabetes	1.2	1.2	1.0	1.6	1.1	0.9	0.5	0.3	0.2	0.2	0.1	0.1
Hypertension	1.0	1.2	1.0	1.5	1.3	1.2	0.3	0.2	0.1	0.1	0.1	0.1
Glomerulonephritis	2.4	1.3	2.6	2.8	2.9	2.8	0.6	0.3	0.4	0.3	0.3	0.3
Other Cause	1.7	2.1	1.9	3.2	2.7	2.7	0.2	0.2	0.2	0.2	0.2	0.2
Total Adjusted Prob.	1.3	1.3	1.2	2.0	1.6	1.5	0.1	0.1	0.1	0.1	0.1	0.1

Table I.22

One-year Survival Probabilities*dialysis patients (censored at transplant), from day 91 to one year + 90 days, unadjusted*

PROBABILITIES	1987	1988	1989	1990	1991	1992	1993	1994	1995	1996	1997
0-4	85.5	77.2	77.5	86.3	76.9	85.7	81.7	90.1	91.9	92.9	91.0
5-9	82.5	90.5	90.3	92.3	87.7	91.6	92.9	98.6	97.8	94.9	95.8
10-14	94.0	90.1	83.2	94.1	91.3	90.7	90.3	98.4	98.0	96.9	97.4
15-19	86.3	90.8	95.3	94.1	93.0	95.3	94.9	93.9	96.6	96.5	98.0
20-24	85.4	87.4	91.0	91.0	91.3	92.5	93.3	95.4	94.7	94.5	97.3
25-29	81.1	83.6	83.9	86.9	87.0	89.6	88.6	89.9	90.7	94.8	96.0
30-34	76.5	83.3	82.0	85.2	85.1	87.1	86.9	88.6	88.7	91.4	93.2
35-39	79.3	81.1	83.5	84.7	84.6	87.0	87.0	87.9	90.1	90.9	92.7
40-44	78.3	81.6	82.1	85.9	84.6	85.8	86.4	88.1	90.0	91.7	91.7
45-49	78.9	80.8	84.1	84.8	84.1	86.0	88.1	87.0	89.3	89.9	91.8
50-54	79.8	80.8	81.2	83.7	83.8	84.3	86.1	86.9	88.1	87.9	88.9
55-59	78.7	78.6	78.9	80.6	81.3	82.2	82.3	83.2	85.5	86.1	86.7
60-64	74.9	75.2	76.7	77.6	78.2	78.9	80.4	80.8	81.3	83.5	84.2
65-69	72.5	73.0	73.3	75.2	74.6	74.9	76.8	75.9	77.2	77.6	76.4
70-74	67.9	67.8	68.8	70.3	69.7	70.3	72.6	71.7	72.5	72.8	71.9
75-79	61.7	63.0	65.0	65.7	65.4	65.5	66.0	67.6	67.3	67.8	66.3
80-84	56.4	58.0	59.9	61.7	60.1	60.4	64.3	63.0	62.6	62.5	61.5
85 +	52.0	47.7	51.6	53.7	55.9	53.3	52.4	55.4	55.4	53.7	54.9
White	69.7	70.8	71.8	73.6	72.7	73.4	74.8	75.4	76.5	76.7	76.1
Black	80.8	81.1	81.4	82.3	82.2	82.9	83.0	82.7	84.0	85.0	85.2
Other	75.7	79.3	82.4	83.9	85.6	84.1	86.4	85.0	85.6	86.0	87.1
Female	74.9	76.1	76.2	77.4	76.7	77.4	78.0	78.5	79.7	80.2	79.6
Male	72.5	73.2	74.6	76.4	75.9	76.6	78.1	78.1	79.5	80.1	79.9
Diabetes	69.5	71.7	73.2	74.3	74.7	75.3	77.6	77.4	78.3	78.7	78.4
Hypertension	73.3	73.6	74.6	76.1	75.0	75.4	76.2	76.0	78.1	77.5	76.7
Glomerulonephritis	80.4	81.3	82.0	83.2	82.2	84.5	86.0	84.7	86.4	87.1	86.9
Cystic Kidney	83.1	84.8	86.5	88.5	89.7	90.2	91.2	90.5	91.6	91.3	92.5
Other Urologic	70.7	70.4	71.1	75.4	78.3	76.3	77.8	77.0	78.9	79.3	81.0
Other Cause	72.6	75.0	73.5	75.5	73.2	72.9	73.7	73.7	74.5	74.3	76.0
Unknown Cause	75.2	71.3	73.4	77.2	73.1	74.6	76.7	75.4	77.3	77.6	74.9
Missing Cause	77.5	77.8	80.6	81.1	84.9	87.0	78.3	86.2	86.9	91.9	90.5
Total Crude Prob.	73.6	74.5	75.4	76.9	76.3	77.0	78.1	78.3	79.6	80.1	79.7

Table I.22 (continued)

STANDARD ERRORS											
	1987	1988	1989	1990	1991	1992	1993	1994	1995	1996	1997
0-4	4.5	4.7	4.7	4.2	4.5	4.0	4.5	3.3	2.4	2.3	2.6
5-9	5.6	5.2	4.7	3.3	5.8	4.7	3.4	1.4	1.5	2.5	2.4
10-14	2.6	3.0	4.0	2.3	2.6	2.7	2.8	1.1	1.2	1.4	1.3
15-19	2.1	1.8	1.2	1.4	1.5	1.2	1.3	1.4	0.9	0.9	0.8
20-24	1.5	1.4	1.2	1.1	1.1	1.0	1.0	0.8	0.8	0.8	0.6
25-29	1.3	1.2	1.1	1.0	1.0	0.9	0.9	0.8	0.8	0.6	0.5
30-34	1.2	1.0	1.0	0.9	0.9	0.8	0.8	0.7	0.7	0.6	0.6
35-39	1.0	0.9	0.9	0.8	0.8	0.7	0.7	0.7	0.6	0.5	0.5
40-44	1.0	0.9	0.8	0.7	0.7	0.7	0.7	0.6	0.5	0.4	0.5
45-49	0.9	0.9	0.7	0.7	0.7	0.6	0.6	0.5	0.5	0.4	0.4
50-54	0.8	0.8	0.7	0.7	0.6	0.6	0.6	0.5	0.5	0.4	0.4
55-59	0.7	0.7	0.7	0.6	0.6	0.6	0.6	0.5	0.5	0.4	0.4
60-64	0.7	0.6	0.6	0.6	0.5	0.5	0.5	0.5	0.4	0.4	0.4
65-69	0.7	0.6	0.6	0.5	0.5	0.5	0.5	0.5	0.4	0.4	0.4
70-74	0.8	0.7	0.7	0.6	0.6	0.5	0.5	0.5	0.5	0.5	0.5
75-79	0.9	0.9	0.8	0.8	0.7	0.7	0.6	0.6	0.6	0.6	0.5
80-84	1.4	1.3	1.2	1.1	1.0	0.9	0.9	0.8	0.8	0.8	0.7
85+	2.5	2.2	1.9	1.8	1.7	1.5	1.4	1.4	1.3	1.2	1.1
White	0.3	0.3	0.3	0.3	0.3	0.2	0.2	0.2	0.2	0.2	0.2
Black	0.4	0.4	0.3	0.3	0.3	0.3	0.3	0.3	0.3	0.2	0.2
Other	1.1	1.0	1.0	0.9	0.8	0.8	0.7	0.6	0.5	0.5	0.5
Female	0.4	0.3	0.3	0.3	0.3	0.3	0.3	0.2	0.2	0.2	0.2
Male	0.3	0.3	0.3	0.3	0.3	0.3	0.2	0.2	0.2	0.2	0.2
Diabetes	0.5	0.4	0.4	0.4	0.3	0.3	0.3	0.3	0.3	0.2	0.2
Hypertension	0.5	0.4	0.4	0.4	0.4	0.3	0.3	0.3	0.3	0.3	0.3
Glomerulonephritis	0.6	0.6	0.6	0.5	0.5	0.5	0.5	0.5	0.4	0.4	0.4
Cystic Kidney	1.1	1.1	1.1	0.9	0.9	0.8	0.8	0.8	0.7	0.8	0.7
Other Urologic	1.9	1.8	1.7	1.6	1.3	1.4	1.4	1.4	1.2	1.2	1.1
Other Cause	0.7	0.7	0.7	0.7	0.7	0.7	0.7	0.6	0.5	0.5	0.5
Unknown Cause	1.0	1.1	1.1	1.0	1.0	0.9	1.0	1.0	0.9	0.9	0.9
Missing Cause	1.1	0.9	1.0	0.9	0.8	0.7	0.6	0.5	0.5	0.4	0.4
Total Crude Prob.	0.3	0.2	0.2	0.2	0.2	0.2	0.2	0.2	0.2	0.2	0.2

Table I.23

Two-year Survival Probabilities*dialysis patients (censored at transplant), from day 91 to two years + 90 days, unadjusted*

PROBABILITIES	1986	1987	1988	1989	1990	1991	1992	1993	1994	1995	1996
0-4	70.3	66.7	60.5	68.9	75.1	71.7	85.7	79.9	81.5	86.4	89.2
5-9	55.8	76.5	81.0	82.0	79.9	87.7	91.6	75.8	94.4	94.8	93.3
10-14	76.5	87.7	83.1	79.9	83.7	83.7	86.4	86.2	92.8	93.0	93.6
15-19	76.5	78.9	82.4	89.5	87.1	84.7	90.5	90.8	90.3	94.5	92.1
20-24	74.0	73.3	80.1	81.8	84.4	82.5	86.3	87.6	91.1	90.6	91.6
25-29	67.5	66.0	70.2	73.2	75.3	76.3	79.0	81.5	82.8	83.6	89.5
30-34	62.5	60.8	68.9	69.0	71.5	73.4	74.7	76.9	77.4	81.6	84.7
35-39	57.6	64.7	68.6	69.4	72.3	71.7	74.6	77.1	78.6	82.4	83.7
40-44	60.6	62.7	66.0	67.6	73.1	70.7	73.8	73.6	77.2	80.5	83.0
45-49	60.9	61.9	65.4	68.1	70.9	69.8	72.4	75.3	75.5	79.7	80.0
50-54	60.5	61.5	64.1	66.0	67.6	68.3	69.2	72.4	75.2	76.2	77.1
55-59	60.2	60.6	60.8	62.3	64.3	65.0	66.2	67.6	68.7	72.0	73.7
60-64	57.9	56.8	57.5	59.0	59.6	61.1	61.8	63.7	65.1	66.1	69.2
65-69	53.2	54.4	55.4	54.9	55.6	56.0	57.2	59.0	58.1	59.8	60.7
70-74	48.3	48.3	49.0	49.7	50.1	50.0	51.5	52.3	52.3	54.1	54.2
75-79	43.5	41.1	42.3	44.5	44.7	44.8	45.2	46.5	47.0	46.7	48.5
80-84	37.0	34.7	34.6	39.0	39.5	37.9	39.1	39.9	40.6	41.3	41.6
85 +	29.4	27.9	28.6	32.4	29.5	30.7	30.6	31.0	32.8	33.1	31.8
White	50.1	50.7	51.7	52.9	54.0	53.3	54.7	55.9	57.0	58.9	59.7
Black	65.0	64.0	66.3	66.7	67.8	67.6	68.9	69.6	69.9	72.3	73.6
Other	61.2	59.4	63.6	67.7	68.7	72.9	71.3	73.6	72.4	74.1	75.4
Female	57.8	57.7	58.9	59.5	60.2	59.5	60.9	61.4	62.4	64.3	65.8
Male	53.3	53.5	55.4	56.9	58.2	58.2	59.3	61.3	61.9	64.6	65.3
Diabetes	46.9	48.2	50.8	53.2	53.8	54.8	55.9	58.6	58.5	60.7	61.6
Hypertension	57.3	55.5	56.7	58.0	58.3	57.8	58.9	59.7	60.1	62.9	62.6
Glomerulonephritis	62.9	64.3	66.9	66.5	69.1	67.8	71.0	73.0	72.2	75.0	75.9
Cystic Kidney	71.5	70.0	70.9	73.3	77.3	77.4	78.7	81.8	80.8	83.0	81.3
Other Urologic	59.2	57.0	56.5	58.5	58.8	65.1	61.1	64.0	61.7	65.6	66.2
Other Cause	58.1	57.5	59.0	58.3	60.8	57.9	57.6	58.4	59.6	61.3	62.0
Unknown Cause	57.9	55.7	54.7	58.2	60.8	55.9	59.6	62.0	59.6	63.1	63.1
Missing Cause	53.8	62.5	63.9	65.4	65.9	71.6	75.0	62.2	73.9	75.3	83.5
Total Crude Prob.	55.4	55.4	57.0	58.1	59.1	58.8	60.1	61.4	62.1	64.4	65.6

Table I.23 (continued)

STANDARD ERRORS											
	1986	1987	1988	1989	1990	1991	1992	1993	1994	1995	1996
0-4	6.3	6.8	6.4	5.8	6.0	5.1	4.0	4.7	4.7	3.3	2.8
5-9	8.9	6.6	7.8	6.9	5.9	5.8	4.7	8.2	3.2	3.2	2.9
10-14	4.8	4.3	4.4	4.5	4.1	3.8	3.5	3.6	2.9	2.4	2.1
15-19	3.0	2.7	2.5	1.9	2.1	2.4	1.8	1.8	1.9	1.2	1.4
20-24	2.1	2.0	1.8	1.8	1.6	1.6	1.4	1.4	1.1	1.1	1.0
25-29	1.6	1.7	1.6	1.4	1.4	1.3	1.3	1.2	1.1	1.1	0.9
30-34	1.4	1.5	1.3	1.2	1.2	1.2	1.1	1.1	1.0	0.9	0.8
35-39	1.4	1.3	1.2	1.1	1.1	1.0	0.9	0.9	0.9	0.8	0.7
40-44	1.3	1.2	1.1	1.1	1.0	0.9	0.9	0.9	0.8	0.7	0.6
45-49	1.2	1.1	1.1	1.0	0.9	0.9	0.8	0.8	0.7	0.6	0.6
50-54	1.1	1.0	0.9	0.9	0.9	0.8	0.8	0.7	0.7	0.6	0.6
55-59	0.9	0.9	0.8	0.8	0.8	0.7	0.7	0.7	0.6	0.6	0.6
60-64	0.8	0.8	0.7	0.7	0.7	0.6	0.6	0.6	0.6	0.5	0.5
65-69	0.8	0.7	0.7	0.6	0.6	0.6	0.6	0.5	0.5	0.5	0.5
70-74	0.9	0.8	0.8	0.7	0.7	0.6	0.6	0.6	0.5	0.5	0.5
75-79	1.0	1.0	0.9	0.8	0.8	0.7	0.7	0.7	0.6	0.6	0.6
80-84	1.5	1.4	1.2	1.2	1.1	1.0	0.9	0.9	0.8	0.8	0.8
85+	2.4	2.2	2.0	1.8	1.6	1.5	1.4	1.3	1.3	1.2	1.1
White	0.4	0.4	0.3	0.3	0.3	0.3	0.3	0.3	0.3	0.3	0.3
Black	0.5	0.5	0.5	0.4	0.4	0.4	0.4	0.4	0.3	0.3	0.3
Other	1.4	1.3	1.2	1.2	1.1	1.0	1.0	0.9	0.8	0.6	0.6
Female	0.4	0.4	0.4	0.4	0.4	0.3	0.3	0.3	0.3	0.3	0.3
Male	0.4	0.4	0.4	0.4	0.3	0.3	0.3	0.3	0.3	0.3	0.3
Diabetes	0.5	0.5	0.5	0.4	0.4	0.4	0.4	0.4	0.3	0.3	0.3
Hypertension	0.6	0.5	0.5	0.5	0.4	0.4	0.4	0.4	0.4	0.4	0.4
Glomerulonephritis	0.8	0.8	0.7	0.7	0.7	0.7	0.6	0.6	0.6	0.6	0.6
Cystic Kidney	1.4	1.4	1.5	1.4	1.3	1.3	1.2	1.1	1.1	1.1	1.1
Other Urologic	2.1	2.1	2.0	1.9	1.9	1.6	1.6	1.7	1.6	1.5	1.4
Other Cause	0.8	0.8	0.8	0.8	0.8	0.8	0.8	0.8	0.7	0.6	0.6
Unknown Cause	1.3	1.2	1.2	1.3	1.2	1.1	1.1	1.1	1.1	1.0	1.0
Missing Cause	1.4	1.3	1.1	1.2	1.1	1.1	1.0	0.7	0.7	0.6	0.5
Total Crude Prob.	0.3	0.3	0.3	0.3	0.2	0.2	0.2	0.2	0.2	0.2	0.2

Table I.24

Five-year Survival Probabilities*dialysis patients (censored at transplant), from day 91 to five years + 90 days, unadjusted*

PROBABILITIES	1983	1984	1985	1986	1987	1988	1989	1990	1991	1992	1993
0-4	55.7	42.2	50.2	55.1	63.4	51.4	56.8	63.0	64.4	66.2	63.6
5-9	43.0	53.6	37.0	45.4	51.3	74.2	73.8	67.6	87.7	69.3	68.9
10-14	62.0	57.1	53.1	61.2	66.1	72.4	63.9	67.0	66.3	73.2	75.1
15-19	57.7	57.3	48.2	61.1	62.6	60.0	73.4	75.4	74.2	79.4	77.6
20-24	49.5	46.8	47.4	57.2	53.0	61.9	61.6	67.7	68.5	70.2	72.8
25-29	36.9	40.0	36.6	46.2	44.5	52.0	49.6	55.2	56.4	60.2	63.8
30-34	35.5	31.6	34.2	39.4	36.9	45.6	48.3	47.5	51.0	53.4	56.4
35-39	34.8	33.4	32.8	32.6	35.6	43.2	43.9	47.8	46.7	49.0	53.7
40-44	33.3	33.3	33.1	36.6	38.4	39.4	41.1	46.3	44.6	47.6	49.4
45-49	31.4	32.0	32.2	33.9	33.4	37.4	36.6	37.8	41.1	42.3	45.4
50-54	31.4	30.2	30.0	30.1	33.0	33.0	33.5	33.5	36.4	37.3	40.6
55-59	30.8	28.5	27.6	29.5	29.8	29.5	29.6	30.6	33.0	32.2	35.1
60-64	27.7	24.8	24.3	25.3	24.4	26.2	26.2	26.8	27.1	28.0	29.3
65-69	20.3	21.8	21.8	21.5	22.0	22.1	21.9	21.8	21.4	23.1	23.6
70-74	15.7	17.2	17.0	15.8	17.1	17.5	17.7	18.2	17.2	18.3	18.9
75-79	10.3	12.7	12.9	13.3	12.5	12.3	13.4	12.1	13.8	12.3	13.3
80-84	7.2	8.0	10.1	8.2	8.5	7.5	9.7	9.6	8.5	8.6	9.3
85+	4.4	5.1	2.5	5.2	4.8	4.0	6.6	6.1	5.1	5.2	6.1
White	22.0	21.3	20.6	20.5	20.6	20.9	21.0	21.1	21.1	21.8	22.7
Black	32.5	33.1	32.6	34.0	33.6	36.2	35.9	37.2	37.0	37.7	38.6
Other	32.7	27.0	26.3	30.3	32.4	34.7	38.0	37.6	38.2	38.3	40.7
Female	28.4	27.9	27.6	27.6	27.1	28.1	27.4	28.0	27.5	28.5	29.1
Male	23.6	23.2	22.3	23.4	23.9	25.3	26.0	26.4	26.7	27.2	29.0
Diabetes	16.3	14.7	14.8	17.1	16.9	18.8	19.0	19.4	20.3	21.0	23.1
Hypertension	28.2	27.1	27.0	26.3	25.4	26.7	27.3	27.6	26.6	27.3	28.1
Glomerulonephritis	34.9	34.5	31.5	32.9	33.7	37.2	36.6	38.1	38.9	40.2	43.3
Cystic Kidney	42.5	39.6	42.2	40.6	41.9	40.1	46.3	43.3	43.7	48.9	54.7
Other Urologic	33.2	34.5	29.8	30.2	30.4	28.6	31.2	29.1	32.8	29.9	36.4
Other Cause	29.7	29.5	28.2	29.2	30.8	31.1	30.4	31.9	31.0	30.9	31.5
Unknown Cause	30.1	28.5	27.9	26.8	25.8	23.9	28.3	29.1	27.4	27.9	33.5
Missing Cause	16.4	21.8	26.2	30.1	36.9	37.5	37.5	39.0	46.2	48.2	31.4
Total Crude Prob.	25.8	25.4	24.7	25.4	25.4	26.6	26.7	27.1	27.1	27.8	29.0

Table I.24 (continued)

STANDARD ERRORS											
	1983	1984	1985	1986	1987	1988	1989	1990	1991	1992	1993
0-4	8.2	8.7	8.8	8.6	7.2	7.2	7.8	8.1	6.6	8.6	7.4
5-9	9.4	9.3	10.6	9.6	10.9	9.5	9.7	8.1	5.8	11.6	9.7
10-14	5.6	6.2	6.8	7.3	7.7	6.2	6.8	6.6	6.5	5.9	5.6
15-19	3.6	4.1	4.1	4.0	3.8	4.0	3.4	3.3	3.4	3.1	3.2
20-24	2.8	2.7	2.7	2.7	2.8	2.6	2.6	2.4	2.3	2.1	2.3
25-29	1.9	1.9	2.0	1.9	2.0	1.9	1.9	1.8	1.8	1.7	1.7
30-34	1.6	1.6	1.6	1.6	1.6	1.5	1.5	1.5	1.5	1.4	1.4
35-39	1.5	1.5	1.4	1.4	1.4	1.4	1.3	1.3	1.2	1.2	1.2
40-44	1.4	1.4	1.4	1.3	1.3	1.2	1.2	1.2	1.1	1.1	1.1
45-49	1.2	1.3	1.2	1.2	1.2	1.1	1.1	1.0	1.0	1.0	1.0
50-54	1.1	1.1	1.0	1.0	1.0	1.0	0.9	0.9	0.9	0.8	0.9
55-59	0.9	0.9	0.9	0.9	0.8	0.8	0.8	0.8	0.7	0.7	0.7
60-64	0.8	0.8	0.7	0.7	0.7	0.7	0.6	0.6	0.6	0.6	0.6
65-69	0.7	0.7	0.7	0.6	0.6	0.6	0.5	0.5	0.5	0.5	0.5
70-74	0.7	0.7	0.7	0.6	0.6	0.6	0.5	0.5	0.5	0.5	0.5
75-79	0.8	0.8	0.8	0.7	0.6	0.6	0.6	0.5	0.5	0.5	0.5
80-84	1.0	1.0	1.0	0.9	0.8	0.7	0.7	0.7	0.6	0.5	0.5
85+	1.4	1.4	0.9	1.2	1.1	0.9	1.0	0.9	0.7	0.7	0.7
White	0.4	0.4	0.3	0.3	0.3	0.3	0.3	0.3	0.3	0.2	0.3
Black	0.6	0.6	0.5	0.5	0.5	0.5	0.4	0.4	0.4	0.4	0.4
Other	1.7	1.5	1.4	1.4	1.3	1.3	1.4	1.3	1.2	1.1	1.0
Female	0.5	0.5	0.4	0.4	0.4	0.4	0.3	0.3	0.3	0.3	0.3
Male	0.4	0.4	0.4	0.4	0.4	0.3	0.3	0.3	0.3	0.3	0.3
Diabetes	0.5	0.5	0.4	0.4	0.4	0.4	0.4	0.3	0.3	0.3	0.3
Hypertension	0.6	0.6	0.5	0.5	0.5	0.5	0.4	0.4	0.4	0.4	0.4
Glomerulonephritis	0.9	0.9	0.8	0.8	0.8	0.8	0.8	0.8	0.8	0.8	0.8
Cystic Kidney	1.7	1.7	1.7	1.7	1.7	1.7	1.7	1.7	1.7	1.6	1.7
Other Urologic	2.4	2.3	2.0	2.0	2.0	1.9	1.8	1.8	1.6	1.5	1.7
Other Cause	0.9	0.9	0.8	0.8	0.8	0.8	0.8	0.8	0.8	0.8	0.8
Unknown Cause	1.4	1.3	1.3	1.2	1.1	1.1	1.2	1.2	1.1	1.0	1.2
Missing Cause	0.8	1.2	1.3	1.3	1.4	1.2	1.3	1.1	1.2	1.1	0.7
Total Crude Prob.	0.3	0.3	0.3	0.3	0.3	0.3	0.2	0.2	0.2	0.2	0.2

Table I.25

Ten-year Survival Probabilities*dialysis patients (censored at transplant), from day 91 to ten years + 90 days, unadjusted*

PROBABILITIES											
	1978	1979	1980	1981	1982	1983	1984	1985	1986	1987	1988
0-4	47.7	35.9	47.5	48.6	31.3	46.4	36.9	16.7	47.2	63.4	38.6
5-9	50.8	31.9	37.4	39.6	38.7	43.0	53.6	24.6	45.4	51.3	74.2
10-14	48.8	33.0	42.4	31.9	34.5	47.5	40.0	35.3	55.6	61.7	55.5
15-19	43.4	45.6	48.8	45.3	31.4	45.5	40.9	33.1	42.7	44.8	47.4
20-24	35.9	39.7	32.2	33.4	30.4	36.3	32.2	32.9	38.0	36.5	47.1
25-29	29.2	27.0	29.6	21.4	23.9	22.8	25.1	24.4	28.7	27.5	31.8
30-34	24.3	26.0	23.2	21.4	17.8	21.8	18.1	19.8	22.7	20.7	25.5
35-39	20.6	20.2	22.0	16.3	16.9	17.3	17.5	17.4	16.8	18.8	20.8
40-44	18.8	18.7	16.9	15.5	15.4	15.8	16.0	16.3	15.8	17.6	19.4
45-49	17.3	16.0	15.1	13.8	12.8	12.7	14.0	11.9	14.2	14.1	16.0
50-54	14.1	14.8	13.0	14.2	13.0	11.4	10.3	11.8	9.6	11.2	11.0
55-59	12.9	12.2	9.2	9.3	9.7	8.3	7.9	8.6	8.1	8.5	7.9
60-64	7.5	7.6	7.5	7.4	5.7	6.9	6.4	6.7	6.3	5.6	6.3
65-69	4.8	4.8	4.5	4.8	3.8	4.6	4.8	5.6	4.4	4.0	4.2
70-74	3.0	2.3	2.9	2.3	3.4	2.8	3.3	3.3	2.6	2.7	2.4
75-79	2.6	1.7	1.5	1.7	1.9	1.1	2.1	1.7	1.7	1.9	1.8
80-84	0.0	0.0	1.5	0.7	1.8	0.3	1.2	0.9	0.8	0.9	0.5
85+	0.0	0.0	1.2	0.0	0.0	0.0	0.8	0.4	0.0	0.0	0.2
White	11.9	11.0	9.7	8.3	7.8	6.9	6.8	6.4	5.9	5.9	5.6
Black	16.7	15.1	14.4	14.7	12.3	11.9	11.7	12.3	11.6	11.3	13.0
Other	11.0	11.3	8.9	10.5	11.4	13.1	10.4	10.0	11.7	11.4	12.5
Female	15.2	13.8	12.8	12.4	10.6	9.9	9.5	9.4	8.4	8.1	8.9
Male	12.3	11.5	10.3	9.2	8.4	7.8	7.8	7.6	7.7	7.7	7.9
Diabetes	5.2	4.7	5.2	4.1	3.6	3.5	3.2	3.4	3.4	3.4	3.8
Hypertension	14.0	12.3	11.0	10.0	8.9	9.2	8.9	8.6	8.1	7.0	7.7
Glomerulonephritis	22.1	18.9	18.6	15.1	13.8	13.7	14.0	12.9	13.4	13.3	14.6
Cystic Kidney	22.9	22.7	19.4	20.6	17.2	17.4	15.8	18.7	14.4	15.7	17.8
Other Urologic	10.3	13.4	11.8	9.2	10.9	10.6	10.0	10.7	9.7	9.8	10.2
Other Cause	16.7	16.6	14.8	11.9	11.8	12.0	11.6	10.9	10.5	12.3	12.5
Unknown Cause	14.9	12.2	11.6	10.2	9.7	9.1	8.7	9.7	9.1	8.7	8.8
Missing Cause	11.6	11.3	10.3	12.7	12.7	6.7	9.5	12.5	12.4	15.6	14.4
Total Crude Prob.	13.6	12.5	11.4	10.7	9.4	8.8	8.6	8.5	8.0	7.9	8.4

Table I.25 (continued)

STANDARD ERRORS											
	1978	1979	1980	1981	1982	1983	1984	1985	1986	1987	1988
0-4	13.4	12.0	12.1	9.6	11.2	10.3	8.9	8.7	10.0	7.2	8.2
5-9	8.9	8.6	9.3	9.2	9.8	9.4	9.3	10.8	9.6	10.9	9.5
10-14	5.5	5.8	6.4	6.6	6.5	7.1	7.6	7.8	8.4	8.3	8.7
15-19	3.5	3.5	3.7	3.6	4.1	4.1	5.0	4.6	4.9	4.5	4.4
20-24	2.6	2.8	2.7	2.6	2.8	2.9	2.8	2.8	3.1	3.1	3.1
25-29	2.1	2.0	2.1	1.9	1.9	1.8	1.9	2.0	2.0	2.0	2.1
30-34	1.8	1.7	1.6	1.5	1.4	1.5	1.4	1.4	1.5	1.5	1.5
35-39	1.5	1.5	1.5	1.3	1.3	1.3	1.3	1.2	1.2	1.2	1.2
40-44	1.4	1.3	1.3	1.2	1.2	1.1	1.1	1.1	1.1	1.1	1.1
45-49	1.2	1.1	1.1	1.0	1.0	0.9	1.0	0.9	0.9	0.9	0.9
50-54	0.9	0.9	0.9	0.8	0.8	0.7	0.7	0.7	0.7	0.7	0.7
55-59	0.8	0.8	0.7	0.6	0.6	0.6	0.5	0.5	0.5	0.5	0.5
60-64	0.7	0.6	0.6	0.6	0.5	0.5	0.4	0.4	0.4	0.4	0.4
65-69	0.5	0.5	0.5	0.5	0.4	0.4	0.4	0.4	0.3	0.3	0.3
70-74	0.5	0.4	0.5	0.4	0.4	0.3	0.4	0.3	0.3	0.3	0.2
75-79	0.7	0.5	0.4	0.4	0.4	0.3	0.3	0.3	0.3	0.3	0.2
80-84	0.0	0.0	0.7	0.4	0.6	0.2	0.4	0.3	0.3	0.3	0.2
85+	0.0	0.0	0.9	0.0	0.0	0.0	0.5	0.3	0.0	0.0	0.2
White	0.4	0.4	0.3	0.3	0.3	0.2	0.2	0.2	0.2	0.2	0.2
Black	0.6	0.5	0.5	0.5	0.4	0.4	0.4	0.4	0.4	0.3	0.3
Other	1.6	1.5	1.3	1.3	1.5	1.3	1.1	1.0	1.0	0.9	0.9
Female	0.5	0.4	0.4	0.4	0.4	0.3	0.3	0.3	0.3	0.2	0.2
Male	0.4	0.4	0.4	0.3	0.3	0.3	0.3	0.2	0.2	0.2	0.2
Diabetes	0.6	0.6	0.5	0.4	0.3	0.3	0.2	0.2	0.2	0.2	0.2
Hypertension	0.8	0.8	0.7	0.5	0.4	0.4	0.4	0.4	0.3	0.3	0.3
Glomerulonephritis	1.1	1.0	1.0	0.8	0.7	0.7	0.7	0.6	0.7	0.6	0.6
Cystic Kidney	2.0	1.9	1.8	1.6	1.3	1.4	1.3	1.4	1.3	1.3	1.4
Other Urologic	2.3	2.4	2.2	1.6	1.7	1.6	1.5	1.4	1.3	1.3	1.3
Other Cause	1.3	1.2	1.1	0.8	0.7	0.7	0.6	0.6	0.6	0.6	0.6
Unknown Cause	1.1	1.0	1.0	0.9	0.9	0.9	0.8	0.9	0.8	0.7	0.7
Missing Cause	0.5	0.4	0.4	0.6	1.1	0.5	0.9	1.0	1.0	1.1	0.9
Total Crude Prob.	0.3	0.3	0.3	0.3	0.2	0.2	0.2	0.2	0.2	0.2	0.2

Table I.26

One-year Survival Probabilities*dialysis patients (censored at transplant), from day 91 to one year + 90 days, adjusted for age, gender, race, & primary diagnosis*

PROBABILITIES	1987	1988	1989	1990	1991	1992	1993	1994	1995	1996	1997
0-19	87.3	88.6	90.1	92.9	89.8	92.7	91.9	95.1	96.4	95.9	96.9
20-44	81.9	84.7	85.5	88.0	87.4	89.1	88.9	90.0	91.1	92.5	93.7
45-64	77.7	78.5	80.1	81.7	82.1	83.0	84.2	84.6	86.1	87.0	88.0
65-74	68.8	69.4	70.4	72.2	72.0	72.5	74.3	73.7	75.0	75.4	74.5
75 +	57.3	58.2	61.1	62.3	61.9	61.8	63.3	64.2	64.1	64.2	63.4
White	67.7	69.3	71.0	73.2	72.9	73.9	75.6	76.7	78.1	78.7	79.4
Black	81.0	81.4	81.9	83.2	83.2	84.1	84.3	84.1	85.2	86.9	87.6
Other	76.5	80.3	83.4	85.7	86.9	86.3	88.7	87.3	88.2	88.8	90.1
Male	71.5	72.8	74.7	77.0	76.9	77.8	79.5	79.8	81.3	82.3	83.3
Female	74.3	75.8	76.6	78.3	78.2	78.9	79.6	80.4	81.6	82.6	83.1
Diabetes	68.5	70.6	72.2	73.5	74.1	74.9	77.4	77.6	78.6	79.7	80.3
Hypertension	72.2	72.9	74.2	76.3	75.6	76.1	77.2	77.4	80.0	80.1	80.3
Glomerulonephritis	81.2	82.1	83.3	84.9	84.0	86.0	87.8	88.0	89.4	90.8	90.9
Other Cause	77.7	78.0	78.3	81.0	80.4	81.2	81.4	82.5	83.5	85.1	86.5
Total Adjusted Prob.	72.8	74.2	75.6	77.6	77.5	78.4	79.6	80.1	81.4	82.4	83.2
STANDARD ERRORS											
0-19	1.6	1.5	1.3	1.1	1.3	1.1	1.2	0.9	0.7	0.7	0.7
20-44	0.5	0.4	0.4	0.4	0.4	0.3	0.3	0.3	0.3	0.3	0.2
45-64	0.4	0.4	0.3	0.3	0.3	0.3	0.3	0.2	0.2	0.2	0.2
65-74	0.5	0.5	0.5	0.4	0.4	0.4	0.4	0.3	0.3	0.3	0.3
75 +	0.8	0.7	0.6	0.6	0.6	0.5	0.5	0.5	0.4	0.4	0.4
White	0.3	0.3	0.3	0.3	0.3	0.2	0.2	0.2	0.2	0.2	0.2
Black	0.4	0.4	0.3	0.3	0.3	0.3	0.3	0.3	0.2	0.2	0.2
Other	1.1	1.0	1.0	0.8	0.7	0.7	0.6	0.6	0.5	0.4	0.5
Male	0.4	0.3	0.3	0.3	0.3	0.3	0.2	0.2	0.2	0.2	0.2
Female	0.4	0.3	0.3	0.3	0.3	0.3	0.3	0.2	0.2	0.2	0.2
Diabetes	0.5	0.5	0.4	0.4	0.3	0.3	0.3	0.3	0.3	0.2	0.2
Hypertension	0.5	0.5	0.4	0.4	0.4	0.3	0.3	0.3	0.3	0.3	0.3
Glomerulonephritis	0.6	0.6	0.5	0.5	0.5	0.5	0.4	0.4	0.4	0.4	0.4
Other Cause	0.4	0.4	0.4	0.4	0.4	0.4	0.3	0.3	0.3	0.3	0.3
Total Adjusted Prob.	0.3	0.2	0.2	0.2	0.2	0.2	0.2	0.2	0.2	0.2	0.2

Table I.27

Two-year Survival Probabilities*dialysis patients (censored at transplant), from day 91 to two years + 90 days, adjusted for age, gender, race, & primary diagnosis*

PROBABILITIES	1986	1987	1988	1989	1990	1991	1992	1993	1994	1995	1996
0-19	73.4	78.7	79.0	84.1	84.0	82.4	88.8	87.2	90.3	93.2	92.0
20-44	65.5	67.8	72.0	73.3	76.9	76.0	78.5	79.5	81.0	83.7	85.7
45-64	59.3	59.5	61.2	63.6	65.5	66.4	67.5	69.6	71.1	73.3	75.1
65-74	48.5	49.3	50.5	51.2	51.8	52.5	53.9	54.9	54.9	56.9	57.5
75 +	37.6	35.3	36.1	40.1	40.0	39.7	40.5	41.7	42.5	42.7	43.7
White	46.8	47.8	49.5	51.8	53.3	53.5	55.2	56.9	58.8	61.0	62.3
Black	65.0	64.2	66.6	67.3	69.0	69.2	70.7	71.6	71.9	74.0	76.3
Other	61.1	60.3	65.0	69.4	71.8	75.3	74.8	77.6	76.1	78.1	79.7
Male	51.6	51.8	54.5	56.8	58.7	59.3	60.8	63.1	64.2	66.9	68.2
Female	55.9	56.6	58.2	59.7	61.2	61.3	62.9	63.5	64.9	66.7	68.9
Diabetes	45.2	46.5	49.0	51.5	52.3	53.6	54.9	58.0	58.5	60.9	62.6
Hypertension	54.7	53.5	55.0	56.8	57.9	57.8	59.1	60.4	61.4	64.8	65.5
Glomerulonephritis	64.1	65.7	68.5	68.9	71.9	70.9	73.7	76.3	77.8	80.0	82.1
Other Cause	62.2	63.1	63.5	64.4	67.3	66.8	68.4	68.2	70.3	72.1	74.9
Total Adjusted Prob.	53.6	54.0	56.2	58.2	59.9	60.3	61.8	63.3	64.5	66.8	68.5
STANDARD ERRORS											
0-19	2.4	2.1	2.1	1.8	1.8	1.9	1.5	1.6	1.4	1.0	1.0
20-44	0.7	0.7	0.6	0.6	0.5	0.5	0.5	0.5	0.4	0.4	0.4
45-64	0.5	0.5	0.4	0.4	0.4	0.4	0.4	0.3	0.3	0.3	0.3
65-74	0.6	0.6	0.5	0.5	0.5	0.4	0.4	0.4	0.4	0.4	0.4
75 +	0.9	0.8	0.7	0.7	0.6	0.6	0.5	0.5	0.5	0.5	0.4
White	0.4	0.4	0.4	0.3	0.3	0.3	0.3	0.3	0.3	0.3	0.3
Black	0.5	0.5	0.5	0.4	0.4	0.4	0.4	0.3	0.3	0.3	0.3
Other	1.4	1.3	1.2	1.2	1.1	1.0	0.9	0.9	0.8	0.6	0.6
Male	0.4	0.4	0.4	0.4	0.3	0.3	0.3	0.3	0.3	0.3	0.3
Female	0.5	0.4	0.4	0.4	0.4	0.3	0.3	0.3	0.3	0.3	0.3
Diabetes	0.6	0.5	0.5	0.5	0.4	0.4	0.4	0.4	0.3	0.3	0.3
Hypertension	0.6	0.6	0.5	0.5	0.5	0.4	0.4	0.4	0.4	0.4	0.4
Glomerulonephritis	0.8	0.8	0.7	0.7	0.7	0.7	0.6	0.6	0.6	0.6	0.6
Other Cause	0.6	0.5	0.5	0.5	0.5	0.5	0.5	0.4	0.4	0.4	0.4
Total Adjusted Prob.	0.3	0.3	0.3	0.3	0.2	0.2	0.2	0.2	0.2	0.2	0.2

Table I.28

Five-year Survival Probabilities*dialysis patients (censored at transplant), from day 91 to five years + 90 days, adjusted for age, gender, race, & primary diagnosis*

PROBABILITIES	1983	1984	1985	1986	1987	1988	1989	1990	1991	1992	1993
0-19	56.7	54.8	47.9	58.4	62.1	60.9	68.5	70.7	71.1	75.3	73.7
20-44	37.8	37.4	37.3	41.6	42.1	47.7	49.1	53.2	52.8	55.4	58.1
45-64	27.1	25.4	25.2	27.4	27.8	29.3	30.0	31.1	33.4	33.7	36.1
65-74	14.7	16.3	16.9	16.1	17.0	17.4	18.0	18.0	18.0	19.4	19.7
75 +	6.4	8.8	9.3	8.7	8.4	7.9	10.3	9.3	9.5	9.1	10.2
White	15.7	16.1	16.3	16.9	17.2	18.0	19.0	19.3	20.3	21.2	22.7
Black	30.4	31.0	30.9	33.3	33.1	35.8	35.9	37.8	38.2	39.1	40.4
Other	29.1	24.0	25.3	29.0	32.2	35.1	40.0	40.8	41.8	42.7	45.9
Male	18.8	18.9	19.2	20.7	21.0	22.9	24.6	25.1	26.5	27.0	29.3
Female	23.1	23.4	23.4	24.3	24.9	25.8	26.2	27.3	27.7	29.1	29.8
Diabetes	14.5	13.4	13.7	14.9	14.8	16.3	16.6	17.0	18.3	19.0	21.3
Hypertension	21.4	21.2	21.8	21.8	21.7	22.5	23.8	24.4	24.1	24.7	26.0
Glomerulonephritis	34.3	34.4	32.0	33.9	35.1	38.9	39.3	41.3	42.5	43.6	47.5
Other Cause	29.8	31.8	32.0	33.5	35.7	35.0	36.6	38.0	39.0	40.4	40.0
Total Adjusted Prob.	20.8	20.9	21.1	22.4	22.8	24.2	25.3	26.1	27.0	28.0	29.5
STANDARD ERRORS											
0-19	2.9	3.1	3.2	3.2	3.1	3.1	2.8	2.7	2.7	2.7	2.7
20-44	0.8	0.8	0.8	0.8	0.8	0.8	0.7	0.7	0.7	0.7	0.7
45-64	0.5	0.5	0.5	0.5	0.4	0.4	0.4	0.4	0.4	0.4	0.4
65-74	0.6	0.6	0.5	0.5	0.4	0.4	0.4	0.4	0.3	0.3	0.3
75 +	0.6	0.6	0.6	0.5	0.4	0.4	0.4	0.4	0.3	0.3	0.3
Race											
White	0.3	0.3	0.3	0.3	0.3	0.3	0.3	0.3	0.3	0.3	0.3
Black	0.6	0.6	0.5	0.5	0.5	0.5	0.5	0.4	0.4	0.4	0.4
Other	1.7	1.5	1.4	1.4	1.3	1.3	1.4	1.4	1.3	1.2	1.2
Gender											
Male	0.4	0.4	0.4	0.4	0.3	0.3	0.3	0.3	0.3	0.3	0.3
Female	0.5	0.5	0.4	0.4	0.4	0.4	0.4	0.4	0.3	0.3	0.3
Primary diagnosis											
Diabetes	0.5	0.5	0.4	0.4	0.4	0.4	0.3	0.3	0.3	0.3	0.3
Hypertension	0.6	0.6	0.5	0.5	0.5	0.5	0.4	0.4	0.4	0.4	0.4
Glomerulonephritis	0.9	0.9	0.8	0.9	0.9	0.9	0.8	0.8	0.8	0.8	0.9
Other Cause	0.6	0.6	0.6	0.6	0.6	0.6	0.6	0.6	0.6	0.6	0.5
Total Adjusted Prob.	0.3	0.3	0.3	0.3	0.3	0.3	0.2	0.2	0.2	0.2	0.2

Table I.29

Ten-year Survival Probabilities*dialysis patients (censored at transplant), from day 91 to ten years + 90 days, adjusted for age, gender, race, & primary diagnosis*

PROBABILITIES	1978	1979	1980	1981	1982	1983	1984	1985	1986	1987	1988
0-19	46.4	38.0	45.2	41.7	32.3	45.1	40.6	30.6	43.5	49.3	48.2
20-44	20.0	20.3	20.0	17.6	17.9	20.0	19.7	19.8	21.5	21.7	24.9
45-64	7.5	7.0	7.2	6.7	6.8	6.8	6.4	6.8	6.9	7.2	7.5
65-74	2.6	1.6	2.1	2.0	2.1	2.2	2.6	3.2	2.4	2.3	2.2
75 +	0.7	0.7	0.8	0.6	0.9	0.3	1.2	0.9	0.7	0.8	0.6
White	3.9	3.8	3.6	3.3	3.4	3.1	3.5	3.7	3.4	3.4	3.3
Black	10.3	9.3	9.7	10.2	9.2	9.4	9.4	10.0	9.7	9.7	11.1
Other	3.9	3.4	4.9	6.5	8.0	8.9	7.4	8.3	9.1	9.2	10.6
Male	4.5	4.5	4.6	4.5	4.6	4.3	4.6	5.2	5.2	5.0	5.0
Female	6.8	5.7	5.8	5.6	5.8	5.6	5.8	5.9	5.2	5.6	6.0
Diabetes	3.9	2.9	3.7	2.7	2.7	2.5	2.4	2.7	2.3	2.3	2.4
Hypertension	5.2	5.0	5.1	4.2	4.4	4.2	4.5	4.6	4.3	4.0	3.7
Glomerulonephritis	15.8	15.1	13.9	12.3	11.9	11.9	12.7	12.2	12.7	12.6	13.8
Other Cause	11.7	11.5	10.7	11.3	11.5	10.8	11.7	12.5	11.8	13.4	13.2
Total Adjusted Prob.	5.5	5.1	5.2	5.0	5.1	4.9	5.1	5.5	5.2	5.2	5.4
STANDARD ERRORS											
0-19	3.1	3.0	3.3	3.1	3.3	3.3	3.7	3.6	3.9	3.7	3.5
20-44	0.8	0.8	0.8	0.7	0.7	0.7	0.7	0.7	0.7	0.7	0.7
45-64	0.5	0.4	0.4	0.3	0.3	0.3	0.3	0.3	0.3	0.3	0.3
65-74	0.4	0.3	0.3	0.2	0.2	0.2	0.2	0.2	0.2	0.2	0.2
75 +	0.3	0.3	0.3	0.2	0.2	0.1	0.2	0.2	0.1	0.1	0.1
White	0.3	0.3	0.2	0.2	0.2	0.2	0.2	0.2	0.1	0.1	0.1
Black	0.6	0.5	0.5	0.4	0.4	0.4	0.4	0.4	0.3	0.3	0.3
Other	1.1	1.0	1.1	1.1	1.3	1.2	0.9	0.9	0.9	0.9	0.9
Male	0.3	0.3	0.3	0.2	0.2	0.2	0.2	0.2	0.2	0.2	0.2
Female	0.5	0.4	0.3	0.3	0.3	0.3	0.2	0.2	0.2	0.2	0.2
Diabetes	0.6	0.5	0.5	0.3	0.3	0.2	0.2	0.2	0.2	0.2	0.2
Hypertension	0.6	0.6	0.5	0.3	0.3	0.3	0.3	0.3	0.2	0.2	0.2
Glomerulonephritis	1.1	1.0	0.9	0.7	0.7	0.7	0.7	0.6	0.7	0.6	0.7
Other Cause	0.4	0.4	0.4	0.4	0.4	0.4	0.4	0.5	0.4	0.5	0.4
Total Adjusted Prob.	0.3	0.2	0.2	0.2	0.2	0.2	0.2	0.2	0.1	0.1	0.1

Table I.31

Two-year Survival Probabilities*transplant patients (cadaveric donor), from day of transplant to two years, unadjusted*

PROBABILITIES	1986	1987	1988	1989	1990	1991	1992	1993	1994	1995	1996
0-4	86.7	91.7	82.8	89.3	82.9	89.3	93.5	96.0	93.9	100.0	97.0
5-9	93.3	97.1	93.3	97.7	96.8	94.1	96.2	97.2	97.4	97.4	100.0
10-14	97.5	95.5	95.9	93.9	95.0	100.0	100.0	98.2	98.8	97.4	98.7
15-19	95.2	95.7	96.9	95.3	96.1	96.9	98.3	96.9	93.4	99.4	97.5
20-24	92.0	95.5	94.8	94.8	96.4	99.0	96.3	94.8	97.1	96.6	97.3
25-29	91.5	92.9	93.0	94.3	94.2	93.5	93.5	94.7	97.3	96.3	96.6
30-34	90.1	91.7	91.9	91.7	94.0	94.5	93.2	95.0	93.4	95.3	94.4
35-39	88.9	89.7	90.1	92.4	92.1	93.6	92.9	94.5	93.7	95.3	95.5
40-44	90.9	86.7	90.4	89.9	90.4	92.6	91.1	93.0	93.0	93.5	93.4
45-49	84.3	86.2	88.1	88.4	88.6	91.0	91.6	90.8	90.8	91.7	92.2
50-54	85.0	86.1	83.4	87.3	88.3	90.5	88.7	86.8	89.9	90.5	89.3
55-59	81.4	81.3	84.0	84.3	86.2	87.2	85.5	87.5	85.5	88.4	89.5
60-64	81.6	80.3	81.7	79.4	83.1	86.0	83.8	85.3	87.0	86.9	86.6
65+	78.0	77.2	74.1	78.6	80.4	79.5	82.2	78.0	81.9	80.4	82.4
White	88.0	87.9	88.4	89.4	89.8	91.3	90.2	90.4	90.4	91.2	91.4
Black	87.8	88.2	88.6	89.2	90.5	91.6	90.6	91.2	91.9	93.3	92.5
Other	89.1	90.4	89.2	89.1	91.2	92.8	92.5	93.2	94.6	93.5	92.8
Female	89.2	89.9	89.8	90.6	90.7	91.9	90.5	90.9	92.1	92.6	91.4
Male	87.3	87.0	87.6	88.5	89.7	91.2	90.4	90.7	90.5	91.5	92.0
Diabetes	81.3	81.4	83.1	86.1	85.9	86.2	86.0	87.0	87.6	88.5	88.5
Hypertension	89.1	87.9	88.9	88.0	88.4	91.9	88.6	89.4	91.1	91.0	91.1
Glomerulonephritis	90.8	91.1	90.4	91.3	92.7	93.4	94.0	94.2	94.1	94.1	93.3
Cystic Kidney	86.5	90.0	90.2	89.9	91.3	92.5	92.7	93.4	90.3	94.5	94.8
Other Urologic	93.3	90.7	90.5	93.2	92.9	91.3	96.4	93.9	90.1	92.2	97.5
Other Cause	91.4	89.6	90.9	90.6	92.5	94.1	91.3	91.5	91.0	93.9	92.7
Unknown Cause	89.8	88.7	89.3	93.3	92.3	91.8	93.3	90.1	91.0	91.8	93.9
Missing Cause	87.1	90.2	88.9	89.9	91.0	94.2	92.4	93.0	94.6	93.3	93.2
Total Crude Prob.	88.0	88.1	88.5	89.3	90.1	91.4	90.5	90.8	91.1	91.9	91.8
STANDARD ERRORS											
0-4	6.1	4.5	6.9	5.7	5.8	5.7	4.3	3.8	4.1	0.0	2.9
5-9	3.7	2.9	3.7	2.3	3.1	4.0	3.7	2.7	2.5	2.5	0.0
10-14	1.7	2.2	2.3	2.9	2.8	0.0	0.0	1.8	1.1	1.8	1.3
15-19	1.7	1.5	1.4	1.7	1.6	1.5	1.2	1.5	1.7	0.6	1.4
20-24	1.4	1.2	1.3	1.3	1.1	0.6	1.2	1.3	1.1	1.1	1.1
25-29	1.2	1.1	1.2	1.0	1.0	1.1	1.1	1.0	0.8	0.8	0.9
30-34	1.1	1.1	1.0	1.1	0.9	0.9	0.9	0.8	0.9	0.8	0.9
35-39	1.1	1.1	1.1	1.0	0.9	0.9	0.9	0.8	0.8	0.7	0.7
40-44	1.1	1.2	1.1	1.0	1.0	0.9	1.0	0.8	0.8	0.8	0.8
45-49	1.4	1.3	1.2	1.2	1.1	1.0	1.0	1.0	0.9	0.9	0.8
50-54	1.5	1.4	1.5	1.4	1.2	1.1	1.2	1.2	1.0	1.0	1.0
55-59	1.8	1.7	1.6	1.6	1.4	1.4	1.4	1.3	1.3	1.2	1.1
60-64	2.4	2.3	2.0	2.2	1.9	1.6	1.7	1.5	1.5	1.4	1.3
65+	4.1	3.5	3.1	3.0	2.5	2.3	2.0	2.1	1.8	1.8	1.7
White	0.5	0.5	0.5	0.5	0.5	0.4	0.4	0.4	0.4	0.4	0.4
Black	0.9	0.9	0.9	0.8	0.8	0.7	0.7	0.7	0.6	0.6	0.6
Other	1.9	1.7	1.7	1.7	1.5	1.4	1.3	1.3	1.1	1.2	1.2
Female	0.7	0.7	0.6	0.6	0.6	0.6	0.6	0.6	0.5	0.5	0.5
Male	0.6	0.6	0.6	0.5	0.5	0.5	0.5	0.5	0.4	0.4	0.4
Diabetes	1.1	1.1	1.1	1.0	0.9	0.9	0.9	0.8	0.8	0.7	0.7
Hypertension	1.1	1.1	1.0	1.1	1.0	0.8	1.0	0.9	0.8	0.8	0.8
Glomerulonephritis	0.8	0.8	0.8	0.8	0.7	0.7	0.6	0.6	0.6	0.6	0.7
Cystic Kidney	1.7	1.4	1.3	1.4	1.2	1.2	1.1	1.0	1.3	0.9	0.9
Other Urologic	2.9	3.3	3.4	3.2	2.8	2.8	2.0	2.1	2.8	2.6	1.4
Other Cause	0.9	1.1	1.0	1.0	0.9	0.8	1.0	1.0	1.0	0.8	0.9
Unknown Cause	1.9	2.0	1.9	1.6	1.6	1.8	1.6	1.8	1.7	1.8	1.6
Missing Cause	1.4	1.3	1.4	1.2	1.2	0.9	1.1	1.1	0.8	0.9	0.9
Total Crude Prob.	0.4	0.4	0.4	0.4	0.4	0.4	0.4	0.4	0.3	0.3	0.3

Table I.32

Five-year Survival Probabilities*transplant patients (cadaveric donor), from day of transplant to five years, unadjusted*

PROBABILITIES	1983	1984	1985	1986	1987	1988	1989	1990	1991	1992	1993
0-4	76.5	75.9	75.0	83.3	91.7	75.9	85.7	80.5	85.7	90.3	92.0
5-9	92.0	91.2	92.1	86.7	91.2	91.1	90.7	96.8	88.2	96.2	91.7
10-14	85.2	94.5	89.6	87.7	94.4	91.9	90.9	95.0	94.9	98.0	96.4
15-19	90.2	93.3	84.7	89.7	93.5	89.5	90.7	92.2	91.4	93.4	93.8
20-24	84.4	88.7	87.3	83.2	89.2	91.1	88.2	92.6	95.4	91.4	88.9
25-29	80.5	79.9	84.4	84.1	86.2	86.2	86.7	87.3	87.0	87.5	89.0
30-34	74.9	74.4	81.6	80.5	80.5	84.4	84.1	87.0	87.4	85.2	85.9
35-39	70.7	76.2	77.2	77.3	77.6	77.7	83.6	84.1	83.3	84.6	86.1
40-44	70.7	70.5	74.4	75.4	75.4	77.0	80.4	81.6	83.2	81.4	85.6
45-49	62.6	69.6	72.4	70.4	71.5	76.1	77.4	76.0	78.9	76.8	79.7
50-54	61.3	69.2	67.6	68.4	73.0	67.9	73.2	74.3	76.7	74.8	73.0
55-59	56.9	63.8	66.7	68.8	64.1	67.7	68.2	70.4	70.0	68.3	71.1
60-64	50.7	54.7	64.1	63.7	63.4	64.8	63.9	68.4	64.8	69.4	65.5
65+	40.7	42.9	42.6	57.0	57.9	53.4	60.4	58.8	58.7	59.9	56.1
White	71.7	74.6	77.2	75.9	76.4	76.7	78.9	79.0	80.1	78.7	79.3
Black	69.5	73.4	72.3	73.7	74.9	75.5	77.8	80.4	78.9	78.3	78.8
Other	78.6	72.2	78.9	82.5	81.9	79.3	82.2	83.8	81.5	83.9	84.9
Female	72.6	77.1	78.0	79.2	79.8	79.0	81.0	81.3	82.4	79.4	80.6
Male	70.7	72.5	74.9	73.6	74.2	74.9	77.4	78.5	78.3	78.6	78.8
Diabetes	53.5	57.5	63.6	62.8	63.4	65.7	70.6	70.2	71.6	68.7	71.0
Hypertension	72.2	74.9	74.2	75.9	73.1	74.6	75.5	77.5	79.1	76.4	76.5
Glomerulonephritis	79.4	80.6	83.7	80.9	81.2	81.2	83.8	84.4	83.0	85.1	86.2
Cystic Kidney	77.1	76.9	79.5	76.4	81.6	80.3	81.7	84.7	83.3	85.0	84.8
Other Urologic	92.7	94.1	81.7	82.7	82.7	82.4	86.4	81.2	82.5	88.0	86.4
Other Cause	75.7	79.8	81.0	82.3	81.2	82.2	81.9	84.1	86.2	82.5	82.9
Unknown Cause	76.6	76.8	80.8	79.7	80.8	78.6	83.1	80.2	79.9	84.1	80.6
Missing Cause	63.1	70.6	70.6	74.8	81.3	78.5	81.8	83.4	82.6	81.8	83.5
Total Crude Prob.	71.4	74.2	76.0	75.7	76.3	76.6	78.8	79.6	79.9	78.9	79.5
STANDARD ERRORS											
0-4	9.9	7.8	7.5	6.7	4.5	7.8	6.5	6.1	6.5	5.2	5.3
5-9	5.3	4.8	4.3	5.0	4.8	4.2	4.4	3.1	5.4	3.7	4.5
10-14	4.5	2.6	3.5	3.6	2.4	3.2	3.5	2.8	2.8	2.0	2.5
15-19	2.5	2.0	2.9	2.4	1.8	2.4	2.4	2.2	2.5	2.2	2.1
20-24	2.3	1.8	1.9	2.0	1.8	1.7	2.0	1.5	1.2	1.7	1.9
25-29	2.0	1.9	1.8	1.6	1.5	1.6	1.4	1.4	1.5	1.5	1.4
30-34	1.9	1.8	1.6	1.4	1.5	1.4	1.4	1.3	1.2	1.3	1.3
35-39	2.0	1.7	1.6	1.5	1.5	1.5	1.3	1.3	1.3	1.3	1.2
40-44	2.2	2.0	1.8	1.6	1.5	1.5	1.4	1.3	1.2	1.3	1.1
45-49	2.5	2.2	1.9	1.7	1.7	1.6	1.6	1.5	1.5	1.5	1.3
50-54	2.7	2.3	2.2	1.9	1.8	1.9	1.8	1.7	1.6	1.6	1.6
55-59	3.4	2.9	2.6	2.1	2.1	2.0	2.0	1.8	1.9	1.9	1.8
60-64	5.8	4.0	3.6	3.0	2.8	2.5	2.6	2.3	2.2	2.2	2.1
65+	9.2	8.2	6.2	4.9	4.1	3.6	3.6	3.1	2.8	2.6	2.5
White	0.9	0.8	0.7	0.7	0.7	0.7	0.6	0.6	0.6	0.6	0.6
Black	1.6	1.4	1.3	1.2	1.2	1.2	1.1	1.0	1.1	1.1	1.0
Other	3.9	3.6	3.1	2.4	2.2	2.2	2.1	1.9	2.0	1.9	1.9
Female	1.3	1.1	1.0	0.9	0.9	0.9	0.8	0.8	0.8	0.8	0.8
Male	1.0	0.9	0.8	0.7	0.7	0.7	0.7	0.7	0.7	0.7	0.6
Diabetes	2.2	1.9	1.6	1.4	1.4	1.4	1.3	1.2	1.2	1.2	1.1
Hypertension	2.0	1.8	1.7	1.5	1.5	1.4	1.4	1.3	1.3	1.3	1.2
Glomerulonephritis	1.4	1.2	1.1	1.0	1.0	1.1	1.0	0.9	1.0	1.0	0.9
Cystic Kidney	2.9	2.4	2.2	2.1	1.8	1.7	1.8	1.5	1.7	1.5	1.5
Other Urologic	4.0	3.3	4.9	4.3	4.3	4.4	4.4	4.2	3.7	3.6	3.0
Other Cause	1.8	1.6	1.4	1.3	1.3	1.3	1.4	1.3	1.2	1.3	1.3
Unknown Cause	2.9	2.8	2.6	2.6	2.5	2.5	2.3	2.4	2.6	2.3	2.4
Missing Cause	2.2	2.1	2.2	1.8	1.7	1.8	1.6	1.5	1.5	1.7	1.6
Total Crude Prob.	0.8	0.7	0.6	0.6	0.6	0.6	0.5	0.5	0.5	0.5	0.5

Table I.33

Ten-year Survival Probabilities*transplant patients (cadaveric donor), from day of transplant to ten years, unadjusted*

PROBABILITIES	1978	1979	1980	1981	1982	1983	1984	1985	1986	1987	1988
0-4	71.4	50.0	80.0	72.7	83.3	70.6	69.0	71.9	83.3	88.9	69.0
5-9	64.7	60.0	84.2	75.0	59.3	84.0	79.4	84.2	82.2	76.5	84.4
10-14	66.0	59.7	82.0	78.6	71.2	80.3	79.5	83.1	79.0	85.4	81.1
15-19	69.4	71.2	78.4	70.7	79.2	81.1	86.6	74.0	81.2	82.1	85.2
20-24	67.2	69.2	69.3	72.4	74.4	74.0	75.7	78.4	72.4	77.1	78.4
25-29	60.2	62.3	62.7	65.5	61.7	66.4	65.0	68.3	68.6	73.4	74.7
30-34	51.0	54.4	52.8	53.3	60.9	59.4	61.6	65.9	64.0	66.2	68.9
35-39	44.3	47.7	47.7	46.3	50.9	52.9	54.8	58.6	58.0	58.9	61.8
40-44	40.4	38.0	47.6	47.3	50.1	50.0	48.8	53.5	55.8	55.8	58.2
45-49	30.8	37.0	38.6	42.5	40.8	40.6	46.2	49.1	48.1	50.7	50.5
50-54	21.1	32.4	35.3	36.1	37.8	40.8	43.8	41.4	46.2	48.8	43.9
55-59	21.6	20.6	24.3	24.0	26.1	33.2	37.3	31.3	37.4	38.8	41.0
60-64	25.0	23.7	20.0	14.0	31.1	24.7	24.0	38.7	34.0	33.8	31.5
65+	0.0	25.0	6.7	0.0	14.3	11.1	17.1	16.4	27.0	29.0	20.2
White	47.8	48.8	52.8	53.5	53.3	55.5	55.4	57.9	56.4	57.3	57.0
Black	40.7	47.8	47.5	45.6	52.3	48.5	52.5	50.5	51.4	54.6	53.5
Other	48.2	54.7	51.6	52.4	57.6	61.6	59.6	59.1	72.0	65.2	62.9
Female	48.3	52.3	53.0	54.8	56.1	57.3	59.0	58.5	61.0	61.0	59.7
Male	45.1	46.9	50.6	49.8	51.6	51.9	52.4	54.7	53.1	54.8	54.5
Diabetes	27.5	29.0	29.7	33.2	30.8	27.8	31.6	36.3	35.5	38.8	39.4
Hypertension	49.2	51.4	54.7	48.4	51.9	54.5	54.7	51.7	50.7	52.3	50.2
Glomerulonephritis	56.0	58.0	64.0	61.0	62.7	62.3	62.5	67.8	66.7	64.3	65.6
Cystic Kidney	42.3	45.0	56.1	52.4	54.1	61.7	59.9	59.8	59.3	63.1	61.8
Other Urologic	84.6	80.0	63.6	79.3	66.7	87.8	80.4	71.7	65.3	72.0	68.9
Other Cause	61.6	63.0	64.4	67.8	62.3	62.3	62.0	65.2	64.2	65.9	65.5
Unknown Cause	44.5	49.5	48.2	50.5	58.7	61.0	59.8	57.6	61.4	61.5	60.1
Missing Cause	36.0	38.3	40.6	43.1	47.5	46.4	53.6	52.8	59.8	62.5	60.9
Total Crude Prob.	46.3	48.9	51.5	51.6	53.2	53.9	54.9	56.1	56.0	57.1	56.5
STANDARD ERRORS											
0-4	15.7	14.7	12.0	12.7	10.3	10.7	8.4	7.8	6.7	5.2	8.4
5-9	11.2	9.5	8.1	7.5	9.2	7.2	6.8	5.8	5.6	7.2	5.3
10-14	6.4	6.2	4.9	5.4	5.5	5.0	4.7	4.2	4.5	3.7	4.5
15-19	4.2	4.0	3.8	4.2	3.7	3.3	2.8	3.6	3.0	2.8	2.8
20-24	3.5	3.3	3.2	2.8	2.8	2.8	2.5	2.3	2.4	2.5	2.4
25-29	3.3	3.1	2.9	2.8	2.6	2.4	2.3	2.3	2.0	1.9	2.0
30-34	3.2	3.0	2.9	2.6	2.3	2.2	2.0	2.0	1.7	1.8	1.8
35-39	3.2	3.1	3.1	2.8	2.5	2.2	2.0	1.9	1.7	1.7	1.7
40-44	3.2	3.2	3.0	2.8	2.6	2.4	2.2	2.1	1.8	1.7	1.8
45-49	3.3	3.1	3.0	3.1	2.6	2.5	2.3	2.1	1.9	1.9	1.8
50-54	3.1	3.6	3.3	3.2	2.8	2.7	2.4	2.4	2.0	2.0	2.0
55-59	4.1	3.9	4.0	3.8	3.5	3.2	2.9	2.5	2.2	2.2	2.2
60-64	7.4	6.7	7.5	5.0	5.8	5.0	3.5	3.6	2.9	2.8	2.4
65+	0.0	16.3	5.1	0.0	10.2	5.5	6.1	4.6	4.4	3.7	2.9
White	1.4	1.3	1.3	1.2	1.1	1.0	0.9	0.9	0.8	0.8	0.8
Black	2.5	2.4	2.2	2.1	1.8	1.7	1.6	1.5	1.4	1.4	1.3
Other	5.4	5.1	5.1	3.9	5.1	4.6	4.0	3.7	2.8	2.8	2.6
Female	1.9	1.8	1.8	1.7	1.5	1.4	1.3	1.2	1.1	1.1	1.0
Male	1.5	1.4	1.4	1.3	1.2	1.1	1.0	0.9	0.8	0.8	0.8
Diabetes	3.8	3.6	3.3	2.8	2.3	2.0	1.8	1.6	1.4	1.4	1.4
Hypertension	3.5	3.8	3.3	3.0	2.5	2.2	2.1	1.9	1.7	1.7	1.6
Glomerulonephritis	2.3	2.2	2.2	2.0	1.7	1.7	1.5	1.4	1.3	1.3	1.3
Cystic Kidney	5.8	4.9	5.0	4.5	3.7	3.3	2.8	2.7	2.4	2.2	2.1
Other Urologic	9.6	7.2	10.0	7.4	7.7	5.0	5.5	5.8	5.4	5.1	5.3
Other Cause	3.3	3.0	2.7	2.6	2.5	2.0	1.9	1.7	1.6	1.6	1.7
Unknown Cause	3.8	3.7	3.8	3.7	3.5	3.3	3.3	3.3	3.1	3.1	3.0
Missing Cause	2.0	2.0	1.9	2.0	2.2	2.3	2.3	2.4	2.0	2.1	2.1
Total Crude Prob.	1.2	1.1	1.1	1.0	0.9	0.9	0.8	0.7	0.7	0.7	0.7

Table I.34

One-year Survival Probabilities*transplant patients (cadaveric donor), from day of transplant to one year, adjusted for age, gender, race, & primary diagnosis*

PROBABILITIES	1987	1988	1989	1990	1991	1992	1993	1994	1995	1996	1997
0-19	95.7	96.0	95.6	94.9	97.1	99.1	97.9	97.3	99.0	99.7	100.0
20-44	94.6	94.9	95.3	95.7	96.7	96.1	96.8	96.8	97.2	97.1	97.4
45 +	88.6	88.6	89.9	90.0	92.2	91.6	91.8	92.5	92.7	93.2	92.9
White	92.0	92.3	93.2	93.2	95.0	94.5	94.9	94.8	95.0	95.5	95.8
Black	92.5	93.2	93.4	94.2	94.8	94.3	95.2	95.4	96.5	96.3	95.3
Other	93.6	92.3	93.3	94.3	95.6	97.1	95.6	97.4	97.4	96.8	98.1
Male	91.7	92.1	92.6	93.2	94.8	94.4	95.1	94.9	95.4	95.7	96.0
Female	93.0	93.0	94.1	93.9	95.2	94.9	94.8	95.6	95.6	95.9	95.6
Diabetes	87.4	87.4	89.9	89.9	90.2	91.0	91.1	91.9	92.0	93.0	93.8
Hypertension	91.2	92.8	92.3	91.9	95.2	92.6	93.7	94.5	95.0	95.7	94.2
Glomerulonephritis	94.4	93.6	94.8	95.8	96.1	96.6	97.0	97.2	97.0	96.2	97.0
Other Cause	93.4	94.0	94.1	94.3	96.5	95.7	95.8	95.7	96.5	96.8	96.6
Total Adjusted Prob.	92.2	92.5	93.2	93.5	95.0	94.6	95.0	95.1	95.5	95.8	95.8
STANDARD ERRORS											
0-19	1.1	1.1	1.2	1.3	1.1	0.6	0.9	0.9	0.6	0.3	1.2
20-44	0.4	0.4	0.4	0.3	0.3	0.3	0.3	0.3	0.3	0.3	0.3
45 +	0.7	0.7	0.6	0.6	0.5	0.5	0.5	0.4	0.4	0.4	0.4
White	0.4	0.4	0.4	0.4	0.3	0.3	0.3	0.3	0.3	0.3	0.3
Black	0.7	0.7	0.7	0.6	0.6	0.6	0.5	0.5	0.4	0.4	0.5
Other	1.4	1.4	1.3	1.2	1.0	0.8	1.0	0.7	0.8	0.9	0.6
Male	0.5	0.5	0.4	0.4	0.3	0.4	0.3	0.3	0.3	0.3	0.3
Female	0.6	0.5	0.5	0.5	0.4	0.4	0.4	0.4	0.4	0.4	0.4
Diabetes	1.0	1.0	0.9	0.8	0.8	0.7	0.7	0.6	0.6	0.6	0.6
Hypertension	1.0	0.9	0.9	0.8	0.7	0.8	0.7	0.6	0.6	0.6	0.7
Glomerulonephritis	0.6	0.6	0.6	0.5	0.5	0.5	0.4	0.4	0.4	0.5	0.5
Other Cause	0.5	0.5	0.5	0.5	0.4	0.4	0.4	0.4	0.4	0.4	0.4
Total Adjusted Prob.	0.4	0.3	0.3	0.3	0.3	0.3	0.3	0.2	0.2	0.2	0.3

Table I.35

Two-year Survival Probabilities*transplant patients (cadaveric donor), from day of transplant to two years, adjusted for age, gender, race, & primary diagnosis*

PROBABILITIES	1986	1987	1988	1989	1990	1991	1992	1993	1994	1995	1996
0-19	94.8	95.4	94.7	95.0	93.9	96.3	97.8	97.1	95.7	98.7	98.3
20-44	91.4	91.4	92.3	93.0	93.7	94.8	93.7	95.0	95.0	95.9	95.4
45 +	82.0	82.8	83.4	84.7	86.3	88.3	87.6	87.5	88.3	89.2	89.3
White	87.7	87.9	88.9	89.9	90.7	92.4	91.8	92.2	92.0	92.9	92.8
Black	86.9	87.7	88.5	89.3	90.7	91.8	90.8	92.1	92.4	93.9	93.7
Other	89.5	91.4	90.4	90.8	92.7	94.0	93.6	93.9	95.7	96.4	95.8
Male	87.0	87.0	88.1	89.1	90.5	92.2	91.5	92.3	91.8	93.1	93.3
Female	88.5	89.7	89.9	90.8	91.3	92.6	91.8	92.2	93.2	93.8	93.1
Diabetes	78.6	79.1	81.0	84.3	84.5	85.2	85.2	86.5	87.4	88.4	88.9
Hypertension	87.5	86.3	88.5	87.8	88.3	92.0	88.5	90.1	91.9	92.3	92.4
Glomerulonephritis	91.0	91.8	91.5	92.5	94.3	94.4	95.2	95.3	95.3	95.6	94.3
Other Cause	89.9	90.5	91.0	91.4	92.6	94.7	93.6	93.6	93.1	94.9	94.9
Total Adjusted Prob.	87.7	88.1	88.9	89.8	90.8	92.3	91.6	92.3	92.4	93.4	93.2
STANDARD ERRORS											
0-19	1.2	1.1	1.3	1.3	1.4	1.2	1.0	1.1	1.2	0.7	0.8
20-44	0.5	0.5	0.5	0.4	0.4	0.4	0.4	0.4	0.4	0.4	0.4
45 +	0.9	0.8	0.8	0.8	0.7	0.6	0.6	0.6	0.6	0.5	0.5
White	0.5	0.5	0.5	0.5	0.4	0.4	0.4	0.4	0.4	0.4	0.4
Black	1.0	0.9	0.9	0.8	0.8	0.7	0.7	0.7	0.6	0.5	0.6
Other	1.9	1.6	1.6	1.6	1.3	1.2	1.2	1.2	0.9	1.0	1.1
Male	0.6	0.6	0.5	0.5	0.5	0.4	0.4	0.4	0.4	0.4	0.4
Female	0.7	0.7	0.6	0.6	0.6	0.5	0.5	0.5	0.5	0.5	0.5
Diabetes	1.3	1.3	1.2	1.1	1.0	0.9	0.9	0.8	0.8	0.7	0.7
Hypertension	1.2	1.2	1.1	1.1	1.0	0.8	1.0	0.9	0.8	0.8	0.8
Glomerulonephritis	0.8	0.7	0.7	0.7	0.6	0.6	0.5	0.5	0.5	0.5	0.6
Other Cause	0.6	0.6	0.6	0.6	0.5	0.4	0.5	0.5	0.5	0.5	0.5
Total Adjusted Prob.	0.4	0.4	0.4	0.4	0.4	0.3	0.3	0.3	0.3	0.3	0.3

Table I.36

Five-year Survival Probabilities*transplant patients (cadaveric donor), from day of transplant to five years, adjusted for age, gender, race, & primary diagnosis*

PROBABILITIES	1983	1984	1985	1986	1987	1988	1989	1990	1991	1992	1993
0-19	88.4	91.2	85.8	88.4	93.4	88.6	90.6	91.3	91.0	94.4	93.8
20-44	75.2	77.1	81.0	81.1	81.9	83.0	85.3	86.9	87.5	86.4	88.1
45 +	53.4	61.2	65.0	65.6	66.4	67.4	70.1	71.0	72.1	71.5	72.2
White	66.1	71.4	75.2	75.0	75.9	77.2	79.5	80.3	82.2	81.5	82.4
Black	63.3	68.4	70.0	71.6	73.6	74.9	77.7	80.4	78.9	78.4	80.4
Other	74.3	71.5	80.0	82.7	83.3	81.1	84.7	86.2	83.9	85.9	86.4
Male	65.2	68.8	73.3	72.6	73.7	75.4	78.0	79.7	80.1	80.5	81.8
Female	66.6	73.4	75.9	77.8	79.0	78.9	81.2	82.3	83.6	81.7	82.8
Diabetes	46.5	50.9	57.7	58.1	59.3	61.8	66.9	67.2	69.4	66.7	69.5
Hypertension	65.8	70.9	69.9	72.5	69.3	73.3	74.4	77.0	78.8	76.1	77.3
Glomerulonephritis	77.4	79.3	83.0	81.1	82.2	82.8	85.7	87.3	85.1	87.8	88.5
Other Cause	71.2	76.1	78.3	79.9	82.4	81.9	83.3	84.9	86.4	85.9	85.9
Total Adjusted Prob.	65.9	70.7	74.2	74.7	75.8	76.9	79.3	80.7	81.5	81.0	82.2
STANDARD ERRORS											
0-19	2.1	1.7	2.1	1.8	1.4	1.8	1.8	1.7	1.8	1.6	1.6
20-44	1.0	0.9	0.8	0.7	0.7	0.7	0.6	0.6	0.6	0.6	0.6
45 +	1.8	1.5	1.3	1.1	1.0	1.0	1.0	0.9	0.9	0.8	0.8
White	1.1	0.9	0.8	0.7	0.7	0.7	0.6	0.6	0.6	0.6	0.6
Black	1.9	1.7	1.4	1.3	1.3	1.2	1.2	1.1	1.1	1.1	1.0
Other	4.6	3.9	3.1	2.4	2.2	2.2	2.0	1.8	2.0	1.8	1.9
Male	1.1	1.0	0.9	0.8	0.8	0.7	0.7	0.7	0.7	0.7	0.6
Female	1.5	1.3	1.1	1.0	0.9	0.9	0.8	0.8	0.8	0.8	0.8
Diabetes	2.8	2.4	1.9	1.7	1.6	1.5	1.4	1.3	1.2	1.2	1.1
Hypertension	2.5	2.1	1.9	1.7	1.7	1.5	1.5	1.3	1.3	1.3	1.3
Glomerulonephritis	1.5	1.3	1.2	1.1	1.0	1.0	1.0	0.9	1.0	0.9	0.9
Other Cause	1.2	1.1	1.0	0.9	0.8	0.8	0.8	0.8	0.7	0.8	0.8
Total Adjusted Prob.	0.9	0.8	0.7	0.6	0.6	0.6	0.5	0.5	0.5	0.5	0.5

Table I.37

Ten-year Survival Probabilities*transplant patients (cadaveric donor), from day of transplant to ten years, adjusted for age, gender, race, & primary diagnosis*

PROBABILITIES	1978	1979	1980	1981	1982	1983	1984	1985	1986	1987	1988
0-19	69.1	62.5	80.4	75.6	74.0	80.7	81.3	77.4	81.3	83.4	82.0
20-44	47.0	47.5	50.6	52.3	56.1	57.7	58.9	63.7	64.1	65.7	67.6
45 +	13.4	24.3	24.4	24.6	27.3	29.3	33.3	35.5	37.9	40.3	38.6
White	33.5	35.8	39.0	41.4	42.5	46.0	48.6	53.0	53.3	55.0	56.0
Black	31.0	35.0	35.1	36.3	44.3	38.9	44.4	45.3	46.9	51.3	51.0
Other	43.3	50.0	41.5	45.8	50.1	50.4	57.5	58.2	71.1	66.1	63.6
Male	29.8	34.1	37.6	38.1	41.4	42.2	45.6	50.0	49.6	52.3	53.2
Female	36.9	39.9	39.3	43.8	45.9	48.0	51.7	53.9	57.5	58.5	58.2
Diabetes	12.6	19.9	17.6	20.1	20.0	20.3	24.2	28.1	29.2	33.1	33.2
Hypertension	33.6	34.0	40.7	41.7	38.9	44.7	47.1	43.5	43.6	45.3	46.1
Glomerulonephritis	46.1	49.5	56.2	52.8	57.4	57.3	58.9	65.4	66.0	64.6	66.9
Other Cause	39.0	42.5	44.7	47.2	51.6	54.8	57.7	59.8	62.5	65.2	64.5
Total Adjusted Prob.	32.5	36.3	38.5	40.3	43.2	44.7	48.0	51.5	52.9	54.9	55.2
STANDARD ERRORS											
0-19	3.7	3.5	3.0	3.1	3.1	2.7	2.4	2.5	2.3	2.1	2.2
20-44	1.6	1.6	1.5	1.3	1.2	1.2	1.0	1.0	0.9	0.9	0.9
45 +	2.3	2.6	2.5	2.3	2.0	1.8	1.5	1.3	1.2	1.1	1.0
White	1.6	1.5	1.5	1.4	1.3	1.1	1.0	1.0	0.8	0.8	0.8
Black	3.7	3.2	2.7	2.5	2.2	1.9	1.9	1.6	1.5	1.4	1.4
Other	8.9	6.1	5.8	4.4	5.7	5.5	4.3	4.0	3.0	2.9	2.8
Male	1.7	1.6	1.6	1.4	1.3	1.2	1.1	1.0	0.9	0.9	0.9
Female	2.5	2.4	2.2	2.0	1.8	1.6	1.5	1.3	1.2	1.1	1.1
Diabetes	4.5	4.3	3.9	3.3	2.8	2.4	2.1	1.8	1.6	1.6	1.5
Hypertension	4.5	5.0	4.1	3.7	3.2	2.6	2.3	2.1	1.9	1.8	1.7
Glomerulonephritis	2.6	2.6	2.5	2.4	1.9	1.8	1.6	1.5	1.3	1.3	1.3
Other Cause	1.6	1.5	1.5	1.5	1.4	1.3	1.2	1.2	1.1	1.1	1.1
Total Adjusted Prob.	1.4	1.4	1.3	1.2	1.1	1.0	0.9	0.8	0.7	0.7	0.7

Table I.38

One-year Survival Probabilities*transplant patients (living donor), from day of transplant to one year, unadjusted*

PROBABILITIES	1987	1988	1989	1990	1991	1992	1993	1994	1995	1996	1997
0-4	96.5	92.9	95.7	94.3	96.2	96.0	97.1	100.0	96.0	98.3	100.0
5-9	100.0	96.8	100.0	96.0	100.0	100.0	97.9	98.0	94.8	98.2	100.0
10-14	100.0	95.1	97.6	98.4	98.8	95.6	98.9	98.8	100.0	98.8	100.0
15-19	96.9	97.6	99.1	98.4	97.8	96.7	99.3	99.4	99.4	99.3	98.6
20-24	98.9	99.4	98.3	98.8	98.9	97.8	99.0	100.0	99.5	100.0	99.6
25-29	97.6	99.2	97.2	99.7	99.3	98.5	97.9	99.7	98.5	99.7	98.3
30-34	95.7	96.1	97.4	98.4	97.9	98.6	97.4	97.6	98.2	99.4	97.9
35-39	97.5	97.6	99.1	99.1	99.2	97.9	97.5	98.4	98.9	98.5	99.2
40-44	93.8	97.1	97.5	96.5	97.5	98.0	98.9	98.3	98.2	98.5	99.2
45-49	96.7	95.2	98.5	94.6	97.7	98.0	97.2	97.1	98.3	97.7	95.7
50-54	93.1	95.6	97.9	97.4	97.6	95.4	95.8	97.2	96.3	96.5	95.5
55-59	100.0	87.8	90.3	92.3	95.9	95.6	93.5	96.3	93.8	94.5	96.2
60-64	90.0	92.0	93.0	91.2	87.0	94.7	96.8	90.8	95.7	94.4	95.3
65+	92.3	87.5	85.7	87.0	90.9	81.3	88.0	100.0	93.5	93.5	94.9
White	97.0	96.5	97.5	97.2	97.9	97.3	97.6	98.3	97.5	97.9	97.4
Black	96.2	98.0	96.7	98.1	96.9	95.8	96.1	97.5	98.2	97.5	97.4
Other	94.6	98.7	100.0	100.0	100.0	98.3	98.2	95.9	98.7	99.0	99.1
Female	96.8	96.8	98.0	97.3	98.0	96.9	97.3	98.8	97.9	97.9	97.0
Male	96.8	96.7	97.1	97.6	97.8	97.3	97.4	97.4	97.6	97.9	98.0
Diabetes	94.7	95.5	98.1	95.9	95.6	95.3	95.0	96.9	96.3	96.8	94.9
Hypertension	98.6	94.9	96.1	94.0	97.2	96.7	96.6	97.1	96.8	97.6	96.6
Glomerulonephritis	97.0	97.8	96.5	98.3	98.6	98.0	98.7	98.6	98.4	98.9	98.7
Cystic Kidney	98.4	98.3	98.5	100.0	99.0	100.0	96.7	96.9	98.3	97.8	96.5
Other Urologic	97.0	96.2	100.0	98.0	96.4	98.7	95.0	97.3	98.9	98.0	100.0
Other Cause	97.1	97.7	98.5	98.5	98.7	96.8	98.4	98.8	97.4	98.3	98.5
Unknown Cause	96.3	96.8	97.2	96.7	98.8	95.2	98.4	100.0	97.6	98.3	97.4
Missing Cause	97.9	96.0	97.6	98.0	99.2	98.0	98.0	98.6	99.0	97.0	98.3
Total Crude Prob.	96.8	96.8	97.5	97.5	97.9	97.1	97.4	98.0	97.7	97.9	97.5

STANDARD ERRORS

0-4	2.4	3.9	3.0	3.9	2.6	2.7	2.0	0.0	2.7	1.7	0.0
5-9	0.0	3.1	0.0	2.7	0.0	0.0	2.0	2.0	2.9	1.7	0.0
10-14	0.0	2.4	1.7	1.6	1.2	2.2	1.1	1.2	0.0	1.2	0.0
15-19	1.5	1.4	0.9	1.1	1.3	1.6	0.7	0.6	0.6	0.7	0.9
20-24	0.8	0.6	0.9	0.8	0.8	1.1	0.7	0.0	0.5	0.0	0.4
25-29	0.9	0.6	1.0	0.3	0.5	0.7	0.8	0.3	0.7	0.3	0.8
30-34	1.3	1.2	1.0	0.8	0.8	0.7	0.9	0.8	0.7	0.4	0.8
35-39	1.1	1.1	0.6	0.6	0.5	0.8	0.9	0.7	0.5	0.7	0.5
40-44	1.9	1.4	1.2	1.3	1.0	0.9	0.6	0.7	0.7	0.7	0.5
45-49	1.6	2.1	1.1	2.0	1.1	1.0	1.1	1.0	0.7	0.7	1.1
50-54	2.7	2.2	1.5	1.5	1.4	1.7	1.4	1.1	1.1	1.1	1.1
55-59	0.0	4.6	3.5	3.0	2.0	1.8	2.1	1.5	1.7	1.5	1.1
60-64	5.4	5.3	3.8	4.8	4.5	2.5	1.8	2.9	1.7	1.8	1.5
65+	7.1	10.9	12.2	6.9	6.0	5.6	4.5	0.0	2.6	2.2	1.9
White	0.5	0.5	0.4	0.4	0.4	0.4	0.4	0.3	0.3	0.3	0.3
Black	1.4	1.0	1.2	0.9	1.0	1.1	1.0	0.7	0.6	0.7	0.7
Other	2.6	1.3	0.0	0.0	0.0	1.2	1.2	1.6	0.9	0.7	0.6
Female	0.7	0.7	0.5	0.6	0.5	0.6	0.5	0.3	0.4	0.4	0.5
Male	0.6	0.6	0.5	0.5	0.4	0.5	0.4	0.4	0.4	0.3	0.3
Diabetes	1.2	1.2	0.8	1.1	1.0	1.0	1.0	0.8	0.8	0.7	0.9
Hypertension	1.0	1.9	1.5	1.9	1.1	1.1	1.0	0.9	0.9	0.8	0.9
Glomerulonephritis	0.8	0.7	0.8	0.6	0.5	0.6	0.4	0.4	0.4	0.4	0.4
Cystic Kidney	1.6	1.7	1.5	0.0	1.0	0.0	1.6	1.5	1.0	1.1	1.3
Other Urologic	2.9	3.7	0.0	2.0	2.5	1.3	2.4	1.9	1.1	1.4	0.0
Other Cause	0.9	0.9	0.7	0.7	0.6	0.9	0.6	0.5	0.7	0.6	0.5
Unknown Cause	2.1	2.2	1.9	1.9	1.2	2.1	1.1	0.0	1.4	1.2	1.5
Missing Cause	1.0	1.3	1.1	1.0	0.6	0.9	0.9	0.7	0.5	0.9	0.6
Total Crude Prob.	0.4	0.4	0.4	0.4	0.3	0.4	0.3	0.3	0.3	0.3	0.3

Table I.39

Two-year Survival Probabilities*transplant patients (living donor), from day of transplant to two years, unadjusted*

PROBABILITIES	1986	1987	1988	1989	1990	1991	1992	1993	1994	1995	1996
0-4	97.4	94.7	90.5	95.7	94.3	96.2	96.0	97.1	100.0	96.0	96.6
5-9	94.9	98.1	96.8	100.0	96.0	100.0	100.0	93.7	98.0	94.8	98.2
10-14	97.5	98.5	93.8	97.6	98.4	98.8	95.6	98.9	98.8	100.0	98.8
15-19	97.9	95.4	96.0	99.1	97.6	97.8	94.3	98.6	98.7	98.1	98.7
20-24	97.5	96.6	98.2	97.2	98.8	98.9	97.8	98.1	99.5	98.5	99.1
25-29	96.3	95.3	97.3	94.8	99.3	98.2	97.4	97.9	99.3	97.4	99.0
30-34	98.1	92.7	94.1	93.9	96.1	95.8	97.2	96.4	97.0	96.9	98.0
35-39	93.1	96.5	96.6	98.6	97.7	96.6	96.8	95.8	97.1	97.0	96.4
40-44	93.4	92.0	97.1	96.3	95.5	95.8	96.4	98.5	96.7	97.0	96.7
45-49	93.8	87.5	93.3	95.4	90.7	95.5	97.0	96.7	93.4	96.7	95.7
50-54	87.2	88.5	93.3	95.8	93.9	96.0	94.7	92.7	94.4	95.6	93.4
55-59	94.7	92.5	83.7	87.5	88.5	93.9	93.3	90.6	92.6	89.6	92.3
60-64	90.5	90.0	84.0	93.0	88.2	83.3	93.4	95.7	87.8	90.8	92.5
65+	80.0	84.6	87.5	85.7	82.6	86.4	79.2	84.0	90.9	85.9	90.3
White	96.0	94.0	94.9	96.0	95.8	96.3	96.3	96.5	96.8	95.8	96.3
Black	93.3	93.4	96.1	94.3	96.3	95.6	93.5	94.1	94.8	96.5	94.7
Other	90.0	93.2	96.1	98.6	98.8	100.0	98.3	98.2	94.6	96.2	98.5
Female	96.0	93.8	95.5	96.7	95.4	96.5	95.9	95.8	96.7	95.9	96.1
Male	95.1	94.0	94.9	95.3	96.3	96.3	96.1	96.5	96.1	95.9	96.2
Diabetes	92.4	88.5	91.6	94.6	93.3	91.8	92.6	93.0	94.5	92.4	93.8
Hypertension	92.4	93.8	93.4	96.1	91.9	96.2	95.3	94.3	95.4	95.1	94.6
Glomerulonephritis	96.9	96.1	97.0	95.2	97.1	97.7	97.3	98.4	97.3	97.3	97.8
Cystic Kidney	98.4	93.7	98.3	98.5	98.8	96.0	100.0	96.7	93.8	96.1	97.3
Other Urologic	100.0	97.0	96.2	97.4	96.0	96.4	97.3	93.8	97.3	97.7	98.0
Other Cause	96.2	94.7	96.0	97.3	97.6	98.4	96.0	97.9	97.2	96.7	96.7
Unknown Cause	93.8	96.3	96.8	94.4	96.7	98.8	93.3	96.7	96.6	96.7	97.5
Missing Cause	97.8	95.3	94.7	96.1	96.0	97.5	98.0	96.0	97.8	97.1	95.7
Total Crude Prob.	95.5	93.9	95.1	95.9	96.0	96.4	96.0	96.2	96.3	95.9	96.2
STANDARD ERRORS											
0-4	2.6	2.9	4.5	3.0	3.9	2.6	2.7	2.0	0.0	2.7	2.4
5-9	3.5	1.9	3.1	0.0	2.7	0.0	0.0	3.5	2.0	2.9	1.7
10-14	1.8	1.4	2.7	1.7	1.6	1.2	2.2	1.1	1.2	0.0	1.2
15-19	1.2	1.8	1.7	0.9	1.3	1.3	2.1	1.0	0.9	1.1	0.9
20-24	1.1	1.4	1.0	1.2	0.8	0.8	1.1	0.9	0.5	0.9	0.6
25-29	1.2	1.3	1.0	1.4	0.5	0.8	1.0	0.8	0.5	1.0	0.6
30-34	0.8	1.7	1.5	1.6	1.2	1.2	1.0	1.1	0.9	1.0	0.7
35-39	1.7	1.3	1.3	0.8	1.0	1.1	1.0	1.2	0.9	0.9	1.0
40-44	2.1	2.1	1.4	1.5	1.4	1.3	1.2	0.7	1.0	0.9	1.0
45-49	2.3	3.0	2.4	1.8	2.5	1.6	1.2	1.2	1.5	1.0	1.0
50-54	3.8	3.4	2.6	2.0	2.2	1.7	1.8	1.9	1.6	1.2	1.5
55-59	2.9	3.2	5.2	3.9	3.6	2.4	2.1	2.5	2.0	2.2	1.8
60-64	6.2	5.4	7.2	3.8	5.4	5.0	2.8	2.1	3.3	2.4	2.1
65+	16.0	9.6	10.9	12.2	7.7	7.1	5.8	5.1	3.8	3.6	2.6
White	0.5	0.6	0.6	0.5	0.5	0.5	0.4	0.4	0.4	0.4	0.4
Black	1.8	1.8	1.3	1.6	1.3	1.2	1.4	1.2	1.0	0.9	1.0
Other	3.8	2.9	2.2	1.4	1.2	0.0	1.2	1.2	1.9	1.5	0.8
Female	0.8	0.9	0.8	0.7	0.8	0.6	0.6	0.6	0.5	0.6	0.5
Male	0.7	0.8	0.7	0.7	0.6	0.6	0.5	0.5	0.5	0.5	0.5
Diabetes	1.4	1.7	1.6	1.3	1.3	1.3	1.3	1.2	1.0	1.1	1.0
Hypertension	2.2	2.0	2.1	1.5	2.2	1.3	1.3	1.3	1.1	1.2	1.2
Glomerulonephritis	0.8	0.9	0.8	1.0	0.7	0.6	0.7	0.5	0.6	0.6	0.5
Cystic Kidney	1.6	3.0	1.7	1.5	1.1	2.0	0.0	1.6	2.1	1.4	1.2
Other Urologic	0.0	2.9	3.7	2.6	2.7	2.5	1.8	2.7	1.9	1.6	1.4
Other Cause	1.0	1.2	1.1	0.9	0.8	0.6	1.0	0.7	0.8	0.8	0.8
Unknown Cause	3.0	2.1	2.2	2.7	1.9	1.2	2.4	1.6	1.7	1.6	1.4
Missing Cause	1.1	1.5	1.5	1.3	1.4	1.0	0.9	1.2	0.9	0.9	1.1
Total Crude Prob.	0.5	0.6	0.5	0.5	0.5	0.4	0.4	0.4	0.4	0.4	0.3

Table I.40

Five-year Survival Probabilities*transplant patients (living donor), from day of transplant to five years, unadjusted*

PROBABILITIES	1983	1984	1985	1986	1987	1988	1989	1990	1991	1992	1993
0-4	80.6	95.0	100.0	94.7	84.2	88.1	93.5	94.3	96.2	94.0	94.2
5-9	93.1	97.6	96.1	92.3	98.1	93.5	100.0	94.0	98.6	96.2	91.7
10-14	95.1	94.8	98.8	93.7	98.5	92.6	96.4	95.2	95.0	92.2	98.9
15-19	93.9	96.2	93.9	94.3	93.1	92.0	98.2	95.3	95.5	92.6	94.2
20-24	93.0	92.5	94.8	91.0	95.5	94.0	96.1	96.4	96.8	95.5	95.2
25-29	88.0	91.3	91.2	91.4	90.2	94.1	91.1	96.9	95.3	93.0	95.2
30-34	87.8	90.2	86.4	92.9	89.7	85.9	88.7	91.0	92.3	92.0	91.1
35-39	86.0	88.2	88.9	85.6	90.1	87.4	94.5	94.5	88.5	93.7	89.8
40-44	78.3	87.4	85.0	84.7	84.0	91.4	86.4	87.1	89.5	89.7	92.1
45-49	84.4	73.2	73.3	83.9	81.7	86.7	87.8	82.9	85.8	87.6	86.5
50-54	75.3	71.4	68.5	79.5	75.9	83.3	82.1	83.3	81.6	85.5	82.7
55-59	75.6	70.3	72.0	73.7	80.6	71.4	77.8	78.2	81.6	86.7	71.7
60-64	66.7	58.3	76.9	81.0	66.7	64.0	67.4	67.6	74.1	82.9	77.4
65+	75.0	66.7	16.7	40.0	46.2	87.5	85.7	69.6	81.8	66.7	62.0
White	87.8	89.4	88.1	89.5	88.6	89.9	90.6	90.3	90.9	91.1	89.5
Black	81.5	81.4	82.7	84.5	85.2	82.9	86.7	91.6	87.8	85.7	87.1
Other	94.3	92.9	89.8	85.0	86.5	88.2	97.1	95.1	94.8	94.8	90.3
Female	88.2	87.3	88.6	89.7	88.0	88.4	90.2	89.6	90.8	90.1	89.1
Male	86.5	89.2	86.7	88.2	88.3	89.3	90.5	91.5	90.5	90.8	89.1
Diabetes	73.1	78.0	75.0	81.0	78.8	81.0	84.1	83.5	79.3	81.3	78.9
Hypertension	84.3	84.7	86.0	86.9	87.0	82.4	86.5	83.9	89.2	90.2	85.9
Glomerulonephritis	92.9	89.9	88.9	91.0	92.4	93.7	91.9	93.5	94.1	93.3	94.1
Cystic Kidney	92.3	93.4	93.4	91.9	87.3	94.9	91.0	94.2	93.9	97.4	90.0
Other Urologic	90.9	88.9	98.1	100.0	90.9	84.6	92.1	94.0	94.6	96.0	88.8
Other Cause	88.7	94.3	91.7	90.6	89.7	91.3	94.4	92.9	95.8	92.6	92.6
Unknown Cause	87.2	91.7	92.4	87.7	91.5	91.9	88.7	94.5	95.3	85.7	91.0
Missing Cause	90.4	91.7	91.8	93.3	91.2	89.0	92.8	93.1	92.0	93.1	92.4
Total Crude Prob.	87.2	88.5	87.5	88.8	88.2	88.9	90.4	90.7	90.6	90.5	89.1

STANDARD ERRORS

0-4	6.5	3.4	0.0	3.6	4.8	4.9	3.6	3.9	2.6	3.3	2.8
5-9	4.6	2.4	2.7	4.2	1.9	4.3	0.0	3.3	1.4	2.6	3.9
10-14	2.4	2.3	1.2	2.7	1.4	2.9	2.0	2.7	2.4	2.8	1.1
15-19	2.0	1.7	2.1	1.9	2.2	2.4	1.2	1.9	1.8	2.4	2.0
20-24	1.7	1.9	1.6	2.0	1.6	1.8	1.4	1.4	1.3	1.5	1.5
25-29	2.0	1.8	1.7	1.7	1.9	1.5	1.8	1.0	1.3	1.5	1.3
30-34	2.0	1.9	2.1	1.6	2.0	2.2	2.1	1.8	1.6	1.6	1.6
35-39	2.6	2.4	2.2	2.4	2.1	2.3	1.5	1.5	2.0	1.4	1.8
40-44	3.7	2.9	3.0	3.1	2.9	2.4	2.7	2.3	2.0	1.9	1.7
45-49	3.7	4.5	4.0	3.5	3.5	3.3	2.8	3.3	2.6	2.3	2.3
50-54	4.9	5.1	4.9	4.5	4.6	3.9	3.9	3.5	3.5	2.8	2.7
55-59	6.6	7.4	6.3	5.8	4.8	6.4	4.9	4.6	3.9	2.9	3.8
60-64	12.9	13.5	11.2	8.3	8.4	9.4	7.0	7.9	5.9	4.3	4.3
65+	18.8	22.2	11.7	18.5	12.7	10.9	12.2	9.3	8.0	6.7	6.8
White	0.9	0.9	0.9	0.8	0.8	0.8	0.8	0.8	0.7	0.7	0.7
Black	2.7	2.8	2.7	2.6	2.6	2.6	2.3	1.9	1.9	2.0	1.7
Other	3.1	3.4	3.9	4.6	3.9	3.7	2.0	2.4	2.3	2.1	2.8
Female	1.3	1.4	1.2	1.2	1.2	1.3	1.1	1.1	1.0	1.0	1.0
Male	1.1	1.0	1.1	1.0	1.0	1.0	0.9	0.9	0.9	0.8	0.8
Diabetes	2.7	2.4	2.4	2.0	2.2	2.2	2.1	2.0	2.0	1.9	1.9
Hypertension	3.1	3.1	3.0	2.8	2.8	3.3	2.7	3.0	2.1	1.8	2.0
Glomerulonephritis	1.2	1.4	1.5	1.4	1.2	1.1	1.2	1.1	1.0	1.0	0.9
Cystic Kidney	3.7	3.1	3.1	3.4	4.2	2.8	3.5	2.5	2.4	1.5	2.7
Other Urologic	4.9	5.9	1.9	0.0	4.9	6.9	4.3	3.3	3.0	2.2	3.5
Other Cause	1.8	1.3	1.5	1.6	1.7	1.6	1.3	1.4	1.0	1.3	1.3
Unknown Cause	3.4	3.0	2.7	4.0	3.1	3.4	3.7	2.4	2.3	3.4	2.6
Missing Cause	2.2	2.3	2.0	1.9	2.0	2.1	1.8	1.8	1.8	1.6	1.7
Total Crude Prob.	0.8	0.8	0.8	0.8	0.8	0.8	0.7	0.7	0.6	0.6	0.6

Table I.41

Ten-year Survival Probabilities*transplant patients (living donor), from day of transplant to ten years, unadjusted*

PROBABILITIES	1978	1979	1980	1981	1982	1983	1984	1985	1986	1987	1988
0-4	50.0	100.0	75.0	77.8	81.3	80.6	90.0	91.7	94.7	82.5	85.7
5-9	81.8	91.7	89.3	100.0	77.5	86.2	95.1	96.1	92.3	90.6	90.3
10-14	79.4	84.3	81.2	83.3	88.5	86.6	88.5	91.6	87.3	92.6	90.1
15-19	82.6	79.1	83.8	91.2	90.8	89.8	89.3	90.2	86.5	85.4	87.2
20-24	80.6	83.9	90.4	85.1	85.8	84.2	87.7	85.3	81.9	89.8	91.1
25-29	77.5	78.0	84.2	80.0	79.2	80.5	80.9	83.8	82.8	83.1	88.2
30-34	62.9	69.5	70.5	67.5	73.9	75.9	78.3	73.2	80.9	75.6	72.7
35-39	59.6	71.7	67.0	67.8	69.7	74.4	74.7	73.9	68.5	75.7	72.8
40-44	55.4	53.1	55.3	53.8	60.3	65.8	66.9	73.6	66.4	64.8	68.3
45-49	39.4	48.2	50.9	50.0	61.3	63.5	54.6	60.0	61.6	68.3	70.5
50-54	36.2	50.0	40.0	49.0	42.6	48.1	62.3	46.1	55.1	49.4	61.1
55-59	56.3	52.9	41.7	45.5	37.5	63.4	40.5	44.0	38.6	52.2	51.0
60-64	0.0	33.3	.	66.7	35.7	33.3	25.0	53.8	47.6	43.3	44.0
65+	25.0	0.0	25.0	66.7	0.0	25.0	66.7	0.0	20.0	23.1	37.5
White	67.4	72.9	75.1	73.5	74.1	77.7	79.2	77.8	75.9	76.8	78.3
Black	69.7	64.9	66.7	59.0	70.1	64.5	62.3	63.9	71.0	69.2	70.2
Other	64.9	78.0	66.7	79.8	71.4	81.1	83.9	79.7	75.0	67.6	81.6
Female	70.6	76.1	74.3	75.5	76.1	76.7	78.7	77.5	76.3	76.3	77.9
Male	65.4	69.8	73.6	70.7	72.0	75.8	76.3	75.2	74.6	75.1	77.1
Diabetes	40.2	51.5	51.1	49.6	53.1	53.8	56.4	55.9	59.3	57.8	60.1
Hypertension	70.1	71.4	62.0	75.6	67.2	70.7	70.2	66.9	70.3	67.8	72.1
Glomerulonephritis	75.6	76.3	81.8	79.0	81.9	83.4	80.9	79.5	81.6	82.8	83.7
Cystic Kidney	62.5	64.0	74.1	62.5	72.1	78.8	86.9	88.5	69.4	69.8	89.8
Other Urologic	100.0	100.0	80.0	84.2	91.9	84.8	85.2	94.2	95.7	81.8	76.9
Other Cause	75.8	73.6	80.9	80.6	83.8	80.1	85.6	83.8	81.6	80.8	83.6
Unknown Cause	68.4	82.4	82.1	65.5	78.6	80.9	84.5	84.8	75.4	82.9	80.6
Missing Cause	63.3	69.9	71.2	73.6	69.2	82.9	85.4	82.5	81.1	83.9	79.3
Total Crude Prob.	67.5	72.2	73.9	72.6	73.5	76.1	77.2	76.1	75.3	75.6	77.4
STANDARD ERRORS											
0-4	18.0	0.0	14.2	9.5	6.8	6.5	4.7	4.5	3.6	5.0	5.3
5-9	8.0	5.5	5.7	0.0	6.5	6.3	3.3	2.7	4.2	4.0	5.2
10-14	4.9	5.0	5.6	5.7	4.0	3.7	3.2	3.0	3.7	3.1	3.3
15-19	3.6	3.9	4.3	2.6	2.4	2.5	2.7	2.6	2.9	3.1	3.0
20-24	3.2	3.1	2.6	2.8	2.4	2.5	2.4	2.6	2.7	2.3	2.2
25-29	3.2	3.2	2.8	2.9	2.5	2.4	2.5	2.2	2.3	2.3	2.0
30-34	4.3	4.0	4.0	3.1	2.8	2.6	2.6	2.8	2.4	2.8	2.8
35-39	5.0	4.5	4.9	4.3	3.5	3.3	3.2	3.1	3.2	3.0	3.1
40-44	5.4	6.2	5.7	5.5	4.3	4.3	4.2	3.7	4.0	3.7	3.9
45-49	5.9	6.6	6.8	5.8	5.0	4.9	5.0	4.4	4.6	4.2	4.4
50-54	6.9	7.4	8.7	6.9	5.9	5.6	5.5	5.2	5.6	5.3	5.1
55-59	11.9	11.6	13.3	10.2	6.9	7.4	7.9	6.9	6.3	6.0	7.0
60-64	0.0	16.4	.	22.2	12.0	12.5	11.3	13.1	10.5	8.8	9.6
65+	16.3	0.0	16.3	22.2	0.0	16.3	22.2	0.0	13.6	10.5	15.3
White	1.6	1.6	1.6	1.4	1.2	1.1	1.1	1.1	1.1	1.1	1.1
Black	4.6	4.5	5.1	4.8	3.4	3.4	3.4	3.4	3.3	3.4	3.2
Other	6.3	5.8	7.4	4.4	6.0	5.3	4.9	5.2	5.5	5.4	4.4
Female	2.3	2.3	2.3	2.0	1.8	1.7	1.7	1.6	1.6	1.6	1.7
Male	2.0	1.9	2.0	1.7	1.5	1.4	1.4	1.4	1.4	1.4	1.3
Diabetes	5.4	6.1	5.3	4.2	3.1	3.0	2.9	2.8	2.6	2.7	2.8
Hypertension	5.5	6.0	6.8	4.6	4.0	3.8	4.0	4.0	3.8	3.9	3.8
Glomerulonephritis	2.7	2.8	2.6	2.3	1.8	1.7	1.9	1.9	1.8	1.8	1.7
Cystic Kidney	9.6	9.4	8.3	8.4	6.7	5.6	4.3	4.0	5.8	5.7	3.9
Other Urologic	0.0	0.0	16.0	8.1	4.4	6.1	6.7	3.2	2.9	6.6	8.1
Other Cause	3.7	3.7	3.4	3.0	2.4	2.3	1.9	2.0	2.1	2.1	2.1
Unknown Cause	4.7	4.0	4.6	5.1	4.0	4.0	3.9	3.7	5.3	4.1	5.0
Missing Cause	2.7	2.6	2.9	2.6	2.9	2.7	2.9	2.8	2.9	2.6	2.7
Total Crude Prob.	1.5	1.5	1.5	1.3	1.1	1.1	1.1	1.1	1.1	1.1	1.1

Table I.42

One-year Survival Probabilities*transplant patients (living donor), from day of transplant to one year, adjusted for age, gender, race, & primary diagnosis*

PROBABILITIES	1987	1988	1989	1990	1991	1992	1993	1994	1995	1996	1997
0-19	98.1	96.0	98.3	97.5	98.3	96.9	98.6	99.2	98.5	98.8	99.4
20-44	96.7	97.8	98.0	98.6	98.7	98.4	98.1	98.7	98.6	99.2	98.8
45 +	94.8	92.2	95.0	93.2	95.7	94.8	95.4	96.2	96.5	96.0	96.1
White	95.9	95.0	96.6	96.4	97.6	96.9	97.2	98.2	97.4	97.8	97.4
Black	94.7	97.5	95.3	97.3	96.3	95.1	95.9	97.2	98.2	97.5	97.9
Other	94.0	98.8	100.0	100.0	100.0	98.4	98.9	96.0	99.2	99.9	100.0
Male	95.4	95.3	96.2	96.7	97.3	96.6	96.9	97.1	97.6	97.9	98.0
Female	95.8	95.8	97.4	96.5	97.8	96.9	97.1	98.8	97.8	97.9	99.1
Diabetes	92.2	92.9	97.5	94.7	95.0	95.0	94.2	96.5	96.3	96.9	95.4
Hypertension	97.9	93.1	95.0	91.6	96.0	96.0	95.8	96.5	97.3	97.5	98.0
Glomerulonephritis	96.2	97.3	95.4	97.9	98.2	97.7	98.5	98.5	98.4	99.0	98.9
Other Cause	96.4	96.0	97.6	97.8	98.3	97.0	97.5	98.4	97.9	97.7	98.2
Total Adjusted Prob.	95.5	95.4	96.7	96.6	97.5	96.7	97.0	97.8	97.6	97.9	97.7
STANDARD ERRORS											
0-19	0.8	1.2	0.8	0.9	0.7	1.0	0.6	0.5	0.6	0.6	0.4
20-44	0.6	0.5	0.4	0.4	0.3	0.3	0.4	0.3	0.3	0.2	0.3
45 +	1.4	1.8	1.3	1.4	1.0	0.9	0.8	0.7	0.6	0.6	0.6
White	0.6	0.7	0.6	0.6	0.4	0.4	0.4	0.3	0.4	0.3	0.3
Black	1.9	1.2	1.7	1.3	1.2	1.3	1.1	0.8	0.6	0.7	0.7
Other	2.9	1.2	0.0	0.0	0.0	1.1	0.9	1.6	0.6	23.4	0.8
Male	0.8	0.8	0.7	0.7	0.5	0.6	0.5	0.5	0.4	0.4	0.3
Female	0.9	1.0	0.7	0.8	0.5	0.6	0.6	0.3	0.4	0.4	114.3
Diabetes	1.8	1.9	1.0	1.5	1.1	1.1	1.2	0.9	0.8	0.7	0.9
Hypertension	1.4	2.5	2.1	2.7	1.6	1.3	1.3	1.1	0.9	0.9	1.0
Glomerulonephritis	1.0	0.9	1.1	0.7	0.6	0.7	0.5	0.5	0.5	0.4	0.4
Other Cause	0.8	0.9	0.7	0.6	0.5	0.6	0.5	0.4	0.4	0.4	0.4
Total Adjusted Prob.	0.6	0.6	0.5	0.5	0.4	0.4	0.4	0.3	0.3	0.3	0.3

Table I.43

Two-year Survival Probabilities*transplant patients (living donor), from day of transplant to two years, adjusted for age, gender, race, & primary diagnosis*

PROBABILITIES	1986	1987	1988	1989	1990	1991	1992	1993	1994	1995	1996
0-19	97.5	96.5	94.6	98.3	97.1	98.3	96.0	97.7	98.9	98.0	98.3
20-44	96.0	94.5	96.4	96.3	97.5	97.3	97.4	97.3	97.8	97.4	97.9
45 +	90.5	87.1	88.7	92.4	88.8	93.3	93.4	93.3	92.3	93.8	93.6
White	94.4	91.9	92.8	94.6	94.5	95.7	95.8	95.9	96.6	95.5	96.2
Black	90.7	91.0	95.1	91.9	94.7	94.7	92.5	93.6	94.1	96.3	94.9
Other	88.6	92.6	96.2	99.8	99.2	100.0	98.4	98.9	94.7	97.6	99.9
Male	92.5	91.3	92.7	93.8	94.9	95.4	95.2	95.8	95.5	95.9	96.3
Female	95.1	91.8	94.0	95.7	94.2	96.1	95.8	95.5	96.4	95.7	96.0
Diabetes	88.8	83.4	87.0	93.0	91.3	90.8	92.1	91.9	93.9	92.5	93.8
Hypertension	89.1	90.5	91.0	95.0	88.9	94.6	94.2	92.9	94.4	95.8	94.3
Glomerulonephritis	95.5	95.1	96.2	93.7	96.4	97.0	96.9	98.1	97.2	97.2	97.9
Other Cause	95.5	93.4	94.3	95.8	96.3	97.1	96.2	96.4	96.6	96.6	96.5
Total Adjusted Prob.	93.4	91.5	93.1	94.6	94.5	95.7	95.4	95.7	95.9	95.8	96.1
STANDARD ERRORS											
0-19	0.9	1.0	1.4	0.8	1.0	0.7	1.1	0.8	0.6	0.8	0.7
20-44	0.6	0.7	0.6	0.6	0.5	0.5	0.5	0.5	0.4	0.4	0.4
45 +	1.9	2.0	2.1	1.5	1.8	1.2	1.0	1.0	1.0	0.8	0.7
White	0.7	0.9	0.9	0.7	0.7	0.5	0.5	0.5	0.4	0.5	0.4
Black	2.5	2.5	1.7	2.2	1.9	1.5	1.6	1.3	1.2	0.9	1.1
Other	4.4	3.2	2.2	0.3	0.8	0.0	1.1	0.9	1.9	1.3	35.2
Male	1.1	1.1	1.0	0.9	0.8	0.7	0.7	0.6	0.6	0.5	0.5
Female	1.0	1.2	1.2	0.9	1.0	0.7	0.7	0.7	0.6	0.6	0.6
Diabetes	2.2	2.6	2.5	1.7	1.9	1.5	1.4	1.4	1.1	1.2	1.0
Hypertension	3.2	3.0	2.9	2.1	3.1	1.9	1.6	1.7	1.4	1.2	1.3
Glomerulonephritis	1.2	1.2	1.0	1.3	0.9	0.8	0.8	0.6	0.7	0.6	0.5
Other Cause	0.9	1.1	1.1	0.9	0.8	0.7	0.7	0.6	0.6	0.6	0.6
Total Adjusted Prob.	0.7	0.8	0.8	0.7	0.6	0.5	0.5	0.5	0.4	0.4	0.4

Table I.44

Five-year Survival Probabilities*transplant patients (living donor), from day of transplant to five years, adjusted for age, gender, race, & primary diagnosis*

PROBABILITIES	1983	1984	1985	1986	1987	1988	1989	1990	1991	1992	1993
0-19	92.3	95.8	96.5	94.3	93.6	91.7	97.3	94.9	96.2	93.5	95.1
20-44	86.5	90.3	89.3	89.7	89.7	89.9	91.9	93.1	93.1	93.2	92.5
45 +	74.2	64.3	62.3	76.1	73.3	76.8	78.7	76.4	81.6	83.6	79.1
White	82.3	84.1	82.2	85.2	84.5	85.7	87.5	87.2	89.3	89.7	87.6
Black	74.6	71.4	74.5	78.7	79.4	78.3	81.0	87.9	85.3	83.6	85.9
Other	93.3	90.1	87.8	82.8	85.2	88.0	99.4	96.7	94.8	95.0	93.1
Male	80.3	82.6	79.8	81.5	83.1	84.6	87.4	87.9	88.1	88.5	87.0
Female	82.4	81.5	82.5	87.4	84.0	84.4	87.2	86.6	89.8	89.8	87.9
Diabetes	63.4	63.5	64.1	72.6	70.1	71.1	79.7	78.3	76.7	80.0	75.4
Hypertension	76.8	71.1	77.5	81.4	79.6	75.1	82.5	77.7	84.6	88.0	82.0
Glomerulonephritis	89.5	85.3	85.0	86.9	90.4	92.1	89.2	91.7	92.5	92.1	93.0
Other Cause	85.4	89.4	88.6	88.3	86.7	87.3	90.6	91.5	92.7	91.3	90.6
Total Adjusted Prob.	81.1	82.0	80.8	83.7	83.5	84.3	87.3	87.1	88.7	88.9	87.4
STANDARD ERRORS											
0-19	1.6	1.1	1.0	1.3	1.4	1.7	1.0	1.3	1.0	1.4	1.2
20-44	1.2	1.0	1.0	1.0	1.0	1.0	0.9	0.8	0.8	0.8	0.8
45 +	3.3	4.0	3.4	2.9	2.7	2.9	2.4	2.4	1.9	1.6	1.6
White	1.4	1.4	1.3	1.2	1.2	1.2	1.1	1.0	0.9	0.8	0.8
Black	3.8	4.1	3.7	3.6	3.6	3.5	3.2	2.9	2.4	2.3	2.0
Other	3.7	5.0	5.2	5.2	4.5	4.3	0.6	1.9	2.5	2.2	2.8
Male	1.7	1.7	1.7	1.6	1.5	1.5	1.3	1.3	1.1	1.0	1.0
Female	2.0	2.2	2.0	1.6	1.7	1.9	1.5	1.5	1.2	1.1	1.2
Diabetes	4.4	4.4	4.2	3.5	3.4	3.6	2.8	3.1	2.4	2.2	2.2
Hypertension	4.5	5.5	4.8	4.2	4.2	4.5	3.9	4.3	3.1	2.2	2.6
Glomerulonephritis	1.8	2.0	2.0	2.0	1.7	1.5	1.6	1.4	1.3	1.2	1.2
Other Cause	1.7	1.5	1.5	1.5	1.5	1.6	1.3	1.1	1.0	1.0	1.0
Total Adjusted Prob.	1.3	1.3	1.3	1.2	1.1	1.2	1.0	1.0	0.8	0.8	0.8

Table I.45

Ten-year Survival Probabilities*transplant patients (living donor), from day of transplant to ten years, adjusted for age, gender, race, & primary diagnosis*

PROBABILITIES	1978	1979	1980	1981	1982	1983	1984	1985	1986	1987	1988
0-19	79.0	84.6	85.6	89.4	87.1	87.1	90.1	92.1	89.2	87.5	88.1
20-44	61.9	66.4	69.3	66.0	71.5	75.1	78.1	77.6	76.6	77.0	77.4
45 +	31.0	36.8	35.3	42.3	38.0	47.0	42.4	38.5	45.3	47.9	52.3
White	47.1	55.6	57.0	56.4	58.6	67.7	68.5	66.8	65.6	67.9	69.2
Black	61.7	55.9	42.0	45.9	61.8	52.5	43.3	46.0	61.0	57.0	61.9
Other	49.3	.	46.1	60.5	63.3	77.3	77.6	74.7	70.5	64.7	80.7
Male	47.0	50.7	55.0	51.9	55.7	64.6	61.2	62.1	59.9	63.8	66.9
Female	49.7	61.7	54.2	59.3	63.2	65.2	68.8	65.2	71.1	68.2	70.1
Diabetes	23.8	25.2	25.8	33.5	34.5	40.2	33.8	40.1	44.6	43.1	42.5
Hypertension	44.0	48.2	28.6	53.9	48.5	57.3	44.6	48.0	58.9	48.7	59.5
Glomerulonephritis	61.4	62.7	70.6	67.8	70.9	75.5	71.8	71.0	72.8	78.2	79.5
Other Cause	56.0	64.3	63.5	60.0	66.6	74.2	77.7	76.9	72.2	74.4	76.0
Total Adjusted Prob.	48.2	54.4	54.1	55.0	58.5	64.9	64.0	63.2	64.1	65.7	67.8
STANDARD ERRORS											
0-19	3.1	2.8	2.9	2.2	2.1	2.0	1.7	1.6	1.9	1.9	1.9
20-44	2.3	2.4	2.3	2.0	1.6	1.6	1.5	1.5	1.5	1.5	1.5
45 +	5.2	6.5	7.5	5.0	4.3	3.9	4.4	3.6	3.6	3.2	3.6
White	2.6	2.7	2.8	2.3	2.0	1.8	1.9	1.7	1.7	1.6	1.7
Black	6.9	6.5	9.7	6.0	5.0	4.6	4.7	4.5	4.5	4.6	4.4
Other	9.2	.	11.0	8.4	7.9	6.4	7.4	7.7	6.4	6.4	5.6
Male	3.0	3.3	3.5	2.7	2.3	2.2	2.3	2.1	2.2	2.1	2.1
Female	3.8	4.2	4.4	3.4	3.1	2.8	2.8	2.7	2.4	2.3	2.6
Diabetes	7.2	8.5	9.9	6.7	5.0	5.0	4.8	4.7	4.6	4.1	4.4
Hypertension	11.2	11.3	11.3	7.8	6.2	5.4	6.4	6.3	5.7	5.7	5.2
Glomerulonephritis	4.2	4.5	4.3	3.6	2.9	2.7	2.7	2.6	2.7	2.6	2.3
Other Cause	2.7	2.6	2.9	2.7	2.4	2.2	2.2	2.0	2.1	2.0	2.2
Total Adjusted Prob.	2.4	2.6	2.7	2.1	1.8	1.7	1.8	1.7	1.6	1.5	1.6