

## **Table of Contents**

### **Section D: Treatment Modalities**

- D.1 Counts of Reported ESRD Patients: by Treatment Modality  
*Incident & Point Prevalent Patients*
- D.2 Incident ESRD Patients: by Dialysis/Functioning Graft Status  
*Incident Patients by Demographic Characteristics & Primary Diagnosis*
- D.3 Prevalent ESRD Patients, by Dialysis/functioning graft Status  
*Point Prevalent Patients on December 31, by Demographic Characteristics & Primary Diagnosis*
- D.11 Incident ESRD Patients for Selected Years: by Diabetic Status & State/Territory
- D.12 Prevalent ESRD Patients for Selected Years: by Diabetic Status & State/Territory  
*Point Prevalent Patients on December 31*

**Table D.1**  
**Counts of Reported ESRD Patients: by Treatment Modality**  
*incident & point prevalent patients*

<b>INCIDENT COUNTS</b>	<b>1991</b>	<b>1992</b>	<b>1993</b>	<b>1994</b>	<b>1995</b>	<b>1996</b>	<b>1997</b>	<b>1998</b>	<b>1999</b>	<b>2000</b>
<b>Modality</b>										
Dialysis	50,878	55,414	57,060	62,351	63,888	67,210	70,362	74,750	77,492	78,427
Functioning graft	998	1,006	1,021	1,023	1,031	897	835	832	867	1,017
<b>Dialysis type</b>										
Center hemodialysis	40,563	44,577	45,245	49,231	51,186	54,501	59,458	65,527	68,976	69,303
Center self hemodialysis	66	63	49	39	67	64	54	47	46	29
Home hemodialysis	30	31	29	39	289	858	136	144	179	226
CAPD	5,418	5,631	5,729	5,476	5,735	5,294	4,736	4,356	4,169	3,885
CCPD	346	389	460	691	1,661	1,839	1,738	1,821	1,825	1,878
Other PD	673	712	664	772	602	487	473	414	310	256
Uncertain dialysis	3,782	4,011	4,884	6,103	4,348	4,167	3,767	2,441	1,987	2,850
All	51,876	56,420	58,081	63,374	64,919	68,107	71,197	75,582	78,359	79,444
<b>PERCENTAGES</b>										
<b>Modality</b>										
Dialysis	98.1	98.2	98.2	98.4	98.4	98.7	98.8	98.9	98.9	98.7
Functioning graft	1.9	1.8	1.8	1.6	1.6	1.3	1.2	1.1	1.1	1.3
<b>Dialysis type</b>										
Center hemodialysis	79.7	80.4	79.3	79.0	80.1	81.1	84.5	87.7	89.0	88.4
Center self hemodialysis	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.0
Home hemo	0.1	0.1	0.1	0.1	0.5	1.3	0.2	0.2	0.2	0.3
CAPD	10.6	10.2	10.0	8.8	9.0	7.9	6.7	5.8	5.4	5.0
CCPD	0.7	0.7	0.8	1.1	2.6	2.7	2.5	2.4	2.4	2.4
Other PD	1.3	1.3	1.2	1.2	0.9	0.7	0.7	0.6	0.4	0.3
Uncertain dialysis	7.4	7.2	8.6	9.8	6.8	6.2	5.4	3.3	2.6	3.6
All	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0
<b>POINT PREVALENT COUNTS</b>										
<b>Modality</b>										
Dialysis	141,866	156,299	168,547	183,272	196,864	208,475	219,609	230,463	241,161	248,066
Functioning graft	39,142	42,944	47,304	49,882	53,450	57,905	62,511	67,203	71,091	74,725
<b>Dialysis type</b>										
Center hemodialysis	115,499	127,322	138,804	150,044	160,439	173,472	187,402	201,197	212,596	220,705
Center self hemodialysis	746	811	152	225	531	552	486	364	336	282
Home hemodialysis	1,607	1,638	343	606	1,768	1,971	1,525	1,204	1,143	1,310
CAPD	17,838	19,691	21,081	21,686	20,646	18,384	16,041	13,930	12,065	10,773
CCPD	2,232	2,901	3,646	4,586	6,983	8,742	9,342	9,333	10,087	10,173
Other PD	675	659	612	741	697	616	537	494	377	421
Uncertain dialysis	3,269	3,277	3,909	5,384	5,800	4,738	4,276	3,941	4,557	4,402
All	181,008	199,243	215,851	233,154	250,314	266,380	282,120	297,666	312,252	322,791
<b>PERCENTAGES</b>										
<b>Modality</b>										
Dialysis	78.4	78.4	78.1	78.6	78.6	78.3	77.8	77.4	77.2	76.9
Functioning graft	21.6	21.6	21.9	21.4	21.4	21.7	22.2	22.6	22.8	23.1
<b>Dialysis type</b>										
Center hemodialysis	81.4	81.5	82.4	81.9	81.5	83.2	85.3	87.3	88.2	89.0
Center self hemodialysis	0.5	0.5	0.1	0.1	0.3	0.3	0.2	0.2	0.1	0.1
Home hemodialysis	1.1	1.0	0.2	0.3	0.9	0.9	0.7	0.5	0.5	0.5
CAPD	12.6	12.6	12.5	11.8	10.5	8.8	7.3	6.0	5.0	4.3
CCPD	1.6	1.9	2.2	2.5	3.5	4.2	4.3	4.0	4.2	4.1
Other PD	0.5	0.4	0.4	0.4	0.4	0.3	0.2	0.2	0.2	0.2
Uncertain dialysis	2.3	2.1	2.3	2.9	2.9	2.3	1.9	1.7	1.9	1.8
All	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0

The incident cohorts and associated modalities are determined at the time of ESRD initiation without applying the 60-day stable modality rule.

The point prevalent cohorts and associated modalities are determined on December 31 of each calendar year without applying the 60-day stable modality rule.

**Table D.2**  
**Incident ESRD Patients: by Dialysis/Functioning Graft Status**

*incident patients by demographic characteristics & primary diagnosis*

ALL DIALYSIS	1991	1992	1993	1994	1995	1996	1997	1998	1999	2000
0-4	114	98	98	82	109	110	96	91	81	115
5-9	75	61	80	91	76	71	58	65	76	63
10-14	163	157	156	162	176	182	157	151	141	145
15-19	358	399	361	351	389	376	311	346	347	312
20-24	767	841	783	769	781	767	680	703	639	635
25-29	1,333	1,332	1,355	1,323	1,251	1,243	1,192	1,204	1,150	1,154
30-34	1,843	1,914	1,917	1,915	1,921	1,812	1,694	1,740	1,708	1,641
35-39	2,300	2,511	2,412	2,556	2,553	2,524	2,404	2,339	2,472	2,281
40-44	2,843	2,985	2,958	3,008	3,299	3,304	3,161	3,375	3,257	3,261
45-49	3,047	3,349	3,536	3,822	4,064	4,158	4,060	4,431	4,498	4,477
50-54	3,428	3,892	3,992	4,579	4,676	4,647	5,014	5,355	5,747	5,784
55-59	4,417	4,781	4,854	5,470	5,514	5,581	5,913	6,411	6,704	6,727
60-64	5,927	6,203	6,317	6,717	7,068	7,059	7,256	7,690	7,755	7,909
65-69	7,878	8,450	8,741	9,483	9,323	9,873	10,309	10,611	10,605	10,609
70-74	7,210	8,021	8,423	9,380	9,453	10,327	10,867	11,335	11,521	11,587
75-79	5,336	5,947	6,225	7,089	7,254	8,180	9,074	9,859	10,760	10,981
80-84	2,734	3,199	3,386	3,948	4,184	4,813	5,591	5,978	6,565	7,036
85+	1,100	1,262	1,465	1,603	1,794	2,182	2,523	3,058	3,461	3,700
Missing/unknown	*	12	*	*	*	*	*	*	*	10
0-19	710	715	695	686	750	739	622	653	645	635
20-44	9,086	9,583	9,425	9,571	9,805	9,650	9,131	9,361	9,226	8,972
45-64	16,819	18,225	18,699	20,588	21,322	21,445	22,243	23,887	24,704	24,897
65-74	15,088	16,471	17,164	18,863	18,776	20,200	21,176	21,946	22,126	22,196
75+	9,170	10,408	11,076	12,640	13,232	15,175	17,188	18,895	20,786	21,717
Unknown	*	12	*	*	*	*	*	*	*	10
Male	27,265	29,454	30,525	33,124	33,936	36,202	37,645	40,008	41,782	42,268
Female	23,613	25,960	26,535	29,218	29,949	31,001	32,715	34,736	35,706	36,156
Unknown	.	.	.	*	*	*	*	*	*	*
White	33,195	35,971	36,319	40,046	40,288	42,176	45,715	48,059	50,782	51,725
Black	15,626	17,065	18,113	19,138	19,449	20,425	20,592	22,100	22,191	21,961
Native American	685	839	781	863	1,030	1,142	815	1,223	916	950
Asian	1,267	1,425	1,664	1,911	2,065	2,236	2,111	2,439	2,592	2,490
Other/unknown	105	114	183	393	1,056	1,231	1,129	929	1,011	1,301
†Hispanic					6,068	7,452	7,399	8,191	8,304	8,423
†Non-Hispanic					57,820	59,758	62,963	66,559	69,188	70,004
Diabetes	18,524	20,554	20,371	23,712	26,114	28,991	31,075	33,428	34,666	35,125
Hypertension	15,452	17,220	16,407	18,784	16,811	17,902	19,208	20,458	21,206	20,968
Glomerulonephritis	5,952	6,171	6,017	6,500	6,568	6,945	6,727	6,790	6,616	6,017
Cystic kidney	1,398	1,486	1,431	1,548	1,586	1,482	1,536	1,601	1,638	1,491
Other urologic	1,065	1,103	959	1,022	1,154	1,353	1,262	1,303	1,392	1,302
Other cause	4,722	5,002	4,905	5,429	6,565	6,906	7,158	7,557	7,762	7,822
Unknown cause	2,207	2,412	2,119	2,196	2,228	2,518	2,662	2,921	2,921	3,036
Missing cause	1,558	1,466	4,851	3,160	2,862	1,113	734	692	1,291	2,666
All	50,878	55,414	57,060	62,351	63,888	67,210	70,362	74,750	77,492	78,427

The incident cohorts and associated modalities are determined at the time of ESRD initiation without applying the 60-day stable modality rule.

The age of the point prevalent patient is determined as of December 31 of each calendar year.

†CMS has only started collecting Hispanic ethnicity data as of April, 1995, so Hispanic data in years prior to 1995 has been left blank.

\* Values for cells with fewer than 10 patients are suppressed.

." Zero values in this cell.

**Table D.2 (continued)**  
**Incident ESRD Patients: by Dialysis/Functioning Graft Status**

*incident patients by demographic characteristics & primary diagnosis*

<b>HEMODIALYSIS</b>	<b>1991</b>	<b>1992</b>	<b>1993</b>	<b>1994</b>	<b>1995</b>	<b>1996</b>	<b>1997</b>	<b>1998</b>	<b>1999</b>	<b>2000</b>
0-4	24	16	22	17	19	17	28	16	16	34
5-9	24	20	30	33	30	27	19	25	27	27
10-14	69	81	65	79	83	79	76	78	68	79
15-19	269	281	252	247	266	266	223	270	271	227
20-24	620	667	610	585	636	623	574	611	553	495
25-29	1,052	1,057	1,024	1,011	967	984	975	1,013	970	909
30-34	1,413	1,453	1,425	1,422	1,457	1,425	1,388	1,439	1,432	1,310
35-39	1,755	1,963	1,841	1,914	1,924	2,029	1,986	1,972	2,112	1,824
40-44	2,142	2,325	2,233	2,224	2,567	2,615	2,609	2,919	2,795	2,674
45-49	2,392	2,642	2,707	2,912	3,081	3,346	3,426	3,821	3,915	3,714
50-54	2,733	3,109	3,149	3,498	3,685	3,756	4,195	4,590	5,023	4,886
55-59	3,645	3,900	3,896	4,304	4,413	4,586	5,006	5,607	5,907	5,827
60-64	4,868	5,094	5,150	5,337	5,741	5,915	6,264	6,802	6,890	6,977
65-69	6,286	6,898	6,968	7,637	7,588	8,172	8,739	9,350	9,555	9,627
70-74	5,846	6,534	6,879	7,614	7,872	8,617	9,295	10,099	10,431	10,635
75-79	4,350	4,894	5,119	5,840	6,119	6,968	7,778	8,871	9,890	10,172
80-84	2,257	2,683	2,750	3,310	3,559	4,131	4,892	5,416	6,086	6,604
85+	911	1,047	1,202	1,322	1,533	1,866	2,173	2,811	3,255	3,527
Missing/unknown	*	*	*	*	*	*	*	*	*	10
0-19	386	398	369	376	398	389	346	389	382	367
20-44	6,982	7,465	7,133	7,156	7,551	7,676	7,532	7,954	7,862	7,212
45-64	13,638	14,745	14,902	16,051	16,920	17,603	18,891	20,820	21,735	21,404
65-74	12,132	13,432	13,847	15,251	15,460	16,789	18,034	19,449	19,986	20,262
75+	7,518	8,624	9,071	10,472	11,211	12,965	14,843	17,098	19,231	20,303
Unknown	*	*	*	*	*	*	*	*	*	10
Male	21,737	23,742	24,366	26,254	27,428	29,942	32,063	35,269	37,361	37,384
Female	18,922	20,929	20,957	23,054	24,114	25,479	27,583	30,443	31,836	32,171
Unknown	.	.	.	*	.	*	*	*	*	*
White	25,776	28,181	28,158	31,136	31,759	33,942	37,847	41,514	44,702	45,546
Black	13,173	14,522	15,128	15,751	16,441	17,690	18,375	20,205	20,476	19,879
Native American	563	702	629	689	848	970	712	1,082	840	824
Asian	1,068	1,191	1,284	1,479	1,634	1,832	1,778	2,116	2,298	2,197
Other/unknown	79	75	124	254	860	989	936	801	885	1,112
†Hispanic					4,927	6,201	6,211	7,177	7,557	7,744
†Non-Hispanic					46,615	49,222	53,437	58,541	61,644	61,814
Diabetes	14,717	16,397	16,155	18,907	21,166	23,796	26,347	29,442	31,058	31,826
Hypertension	12,703	14,396	13,539	15,573	14,147	15,330	16,701	18,418	19,462	19,403
Glomerulonephritis	4,654	4,794	4,623	4,992	5,089	5,436	5,483	5,667	5,680	5,135
Cystic kidney	1,033	1,128	1,051	1,127	1,173	1,121	1,211	1,300	1,316	1,221
Other urologic	904	895	787	818	966	1,137	1,078	1,153	1,253	1,181
Other cause	3,671	4,006	3,888	4,320	5,369	5,764	6,060	6,667	6,965	7,138
Unknown cause	1,773	1,997	1,698	1,797	1,870	2,140	2,275	2,630	2,639	2,800
Missing cause	1,204	1,058	3,582	1,775	1,762	699	493	441	828	854
All	40,659	44,671	45,323	49,309	51,542	55,423	59,648	65,718	69,201	69,558

The incident cohorts and associated modalities are determined at the time of ESRD initiation without applying the 60-day stable modality rule.

The age of the incident patient is determined as of the date of ESRD initiation.

†CMS has only started collecting Hispanic ethnicity data as of April, 1995, so Hispanic data in years prior to 1995 has been left blank.

\* Values for cells with fewer than 10 patients are suppressed.

." Zero values in this cell.

**Table D.2 (continued)**  
**Incident ESRD Patients: by Dialysis/Functioning Graft Status**

*incident patients by demographic characteristics & primary diagnosis*

<b>PERITONEAL DIALYSIS</b>	<b>1991</b>	<b>1992</b>	<b>1993</b>	<b>1994</b>	<b>1995</b>	<b>1996</b>	<b>1997</b>	<b>1998</b>	<b>1999</b>	<b>2000</b>
0-4	52	39	37	34	71	79	65	72	60	78
5-9	37	15	28	34	41	39	36	38	45	36
10-14	64	37	50	42	71	89	75	65	69	63
15-19	63	79	67	61	107	94	81	71	68	64
20-24	107	127	120	114	115	115	101	83	71	69
25-29	225	197	227	200	227	223	194	173	157	139
30-34	343	363	336	310	348	333	269	268	248	199
35-39	419	417	407	400	483	414	359	319	315	288
40-44	549	492	495	432	557	562	467	399	396	380
45-49	488	516	558	540	753	653	529	531	492	464
50-54	507	546	575	608	726	688	668	643	591	551
55-59	536	622	603	613	770	730	698	678	651	565
60-64	706	752	672	712	887	807	739	657	674	656
65-69	653	653	772	721	853	778	731	743	687	712
70-74	515	563	603	668	659	767	671	658	660	648
75-79	328	383	388	404	437	455	475	456	477	517
80-84	133	158	184	205	219	217	221	230	229	238
85+	39	59	67	69	72	90	95	93	104	96
Missing/unknown	.	*	.	.	.	.	.	.	.	.
0-19	216	170	182	171	290	301	257	246	242	241
20-44	1,643	1,596	1,585	1,456	1,730	1,647	1,390	1,242	1,187	1,075
45-64	2,237	2,436	2,408	2,473	3,136	2,878	2,634	2,509	2,408	2,236
65-74	1,168	1,216	1,375	1,389	1,512	1,545	1,402	1,401	1,347	1,360
75+	500	600	639	678	728	762	791	779	810	851
Unknown	.	*	.	.	.	.	.	.	.	.
Male	3,177	3,220	3,251	3,286	3,944	3,821	3,418	3,229	3,163	3,127
Female	2,587	2,800	2,938	2,881	3,452	3,312	3,056	2,948	2,831	2,636
Unknown	.	.	.	.	.	.	.	.	.	.
White	4,188	4,373	4,420	4,409	5,211	5,041	4,742	4,468	4,463	4,272
Black	1,381	1,425	1,518	1,446	1,689	1,532	1,345	1,296	1,222	1,126
Native American	75	73	68	80	116	120	70	100	51	77
Asian	114	143	176	209	272	283	201	220	180	185
Other/unknown	*	*	*	23	108	157	116	93	78	103
†Hispanic					605	697	606	611	587	582
†Non-Hispanic					6,791	6,436	5,868	5,566	5,407	5,181
Diabetes	2,250	2,420	2,312	2,456	3,130	3,238	2,912	2,767	2,732	2,644
Hypertension	1,304	1,374	1,377	1,409	1,431	1,301	1,285	1,256	1,189	1,173
Glomerulonephritis	891	935	893	903	1,155	1,135	986	931	802	778
Cystic kidney	277	265	286	286	338	300	275	261	289	245
Other urologic	70	84	90	85	109	123	101	97	105	95
Other cause	618	570	573	551	752	731	661	638	617	563
Unknown cause	215	216	227	185	214	243	216	193	189	178
Missing cause	139	156	431	292	267	62	38	34	71	87
All	5,764	6,020	6,189	6,167	7,396	7,133	6,474	6,177	5,994	5,763

The incident cohorts and associated modalities are determined at the time of ESRD initiation without applying the 60-day stable modality rule.

The age of the incident patient is determined as of the date of ESRD initiation.

†CMS has only started collecting Hispanic ethnicity data as of April, 1995, so Hispanic data in years prior to 1995 has been left blank.

\* Values for cells with fewer than 10 patients are suppressed.

." Zero values in this cell.

**Table D.2 (continued)**  
**Incident ESRD Patients: by Dialysis/Functioning Graft Status**  
*incident patients by demographic characteristics & primary diagnosis*

TRANSPLANT	1991	1992	1993	1994	1995	1996	1997	1998	1999	2000
0-4	17	19	22	13	15	18	12	*	12	14
5-9	33	30	17	31	26	23	21	19	17	17
10-14	48	36	38	40	35	26	29	30	34	29
15-19	44	39	64	60	64	48	36	26	38	39
20-24	51	51	44	46	43	43	40	41	29	29
25-29	97	92	88	81	80	52	56	50	52	57
30-34	151	135	133	135	114	85	69	77	59	73
35-39	125	136	133	127	156	105	85	119	87	94
40-44	159	138	125	132	120	116	104	91	111	119
45-49	99	108	121	135	125	118	101	93	115	122
50-54	58	87	94	83	95	104	86	100	93	121
55-59	58	67	58	76	66	61	86	75	79	109
60-64	35	40	47	40	52	62	58	45	58	82
65-69	17	21	24	19	28	29	44	47	55	74
70-74	*	*	*	*	11	*	*	*	26	36
75-79	*	.	*	*	*	*	*	*	*	*
80-84	.	*	.	.	.	.	.	*	*	.
85+	.	.	.	.	.	.	*	*	.	.
Missing/unknown	.	.	*	.	.	.	.	.	.	.
0-19	142	124	141	144	140	115	98	82	101	99
20-44	583	552	523	521	513	401	354	378	338	372
45-64	250	302	320	334	338	345	331	313	345	434
65-74	22	27	33	22	39	34	48	54	81	110
75+	*	*	*	*	*	*	*	*	*	*
Unknown	.	.	*	.	.	.	.	.	.	.
Male	587	566	605	570	579	519	490	461	492	558
Female	411	440	416	453	452	378	345	371	375	459
Unknown	.	.	.	.	.	.	.	.	.	.
White	878	865	898	885	877	776	739	735	767	897
Black	80	112	88	99	110	78	69	64	69	79
Native American	15	*	*	10	13	*	*	*	*	*
Asian	23	20	19	22	25	24	10	20	19	20
Other/unknown	*	*	10	*	*	11	11	*	*	18
†Hispanic					33	47	36	48	46	59
†Non-Hispanic					998	850	799	784	821	958
Diabetes	222	221	223	217	210	228	198	187	186	185
Hypertension	46	65	58	62	51	46	56	45	54	70
Glomerulonephritis	151	181	168	161	197	168	156	161	154	170
Cystic kidney	93	93	88	68	105	108	84	100	100	113
Other urologic	36	28	40	29	31	32	34	33	20	37
Other cause	137	140	143	137	166	155	161	129	140	150
Unknown cause	35	37	48	35	29	35	32	26	14	45
Missing cause	278	241	253	314	242	125	114	151	199	247
All	998	1,006	1,021	1,023	1,031	897	835	832	867	1,017

The incident cohorts and associated modalities are determined at the time of ESRD initiation without applying the 60-day stable modality rule.

The age of the incident patient is determined as of the date of ESRD initiation.

†CMS has only started collecting Hispanic ethnicity data as of April, 1995, so Hispanic data in years prior to 1995 has been left blank.

\* Values for cells with fewer than 10 patients are suppressed.

." Zero values in this cell.

**Table D.3**  
**Prevalent ESRD Patients, by Dialysis/Functioning Graft Status**

*point prevalent patients on December 31, by demographic characteristics & primary diagnosis*

<b>ALL DIALYSIS</b>	<b>1991</b>	<b>1992</b>	<b>1993</b>	<b>1994</b>	<b>1995</b>	<b>1996</b>	<b>1997</b>	<b>1998</b>	<b>1999</b>	<b>2000</b>
0-4	144	159	156	157	177	174	149	147	149	168
5-9	145	143	145	165	166	151	153	130	142	153
10-14	338	336	344	362	378	361	336	347	313	312
15-19	891	927	944	881	944	953	932	945	904	837
20-24	2,202	2,357	2,410	2,471	2,514	2,470	2,414	2,438	2,470	2,449
25-29	3,914	4,100	4,137	4,239	4,451	4,534	4,577	4,711	4,754	4,584
30-34	6,021	6,325	6,649	6,886	7,156	7,089	6,959	6,874	7,064	7,064
35-39	7,727	8,406	8,801	9,348	9,827	9,920	10,019	10,133	10,520	10,466
40-44	9,487	10,196	10,804	11,325	12,352	12,898	13,430	13,804	14,244	14,254
45-49	10,100	11,178	12,257	13,574	15,086	16,091	16,524	17,297	18,171	18,660
50-54	11,095	12,454	13,558	15,045	16,145	17,293	18,382	19,721	21,518	22,803
55-59	13,419	14,537	15,615	16,899	18,185	19,270	20,633	22,033	23,245	24,039
60-64	16,607	17,742	18,728	19,932	21,401	22,415	23,525	24,668	25,680	26,369
65-69	20,152	22,199	23,878	25,597	26,755	28,040	29,019	29,879	30,802	31,243
70-74	18,050	20,382	22,404	24,906	26,434	27,989	29,688	30,812	31,395	31,750
75-79	12,523	14,139	15,456	17,550	19,365	21,508	23,603	25,348	26,910	27,928
80-84	6,457	7,641	8,648	9,823	10,864	11,938	13,192	14,249	15,159	16,703
85+	2,581	3,063	3,604	4,109	4,662	5,379	6,069	6,914	7,704	8,263
Missing/unknown	13	15	*	*	*	*	*	13	17	21
0-19	1,518	1,565	1,589	1,565	1,665	1,639	1,570	1,569	1,508	1,470
20-44	29,351	31,384	32,801	34,269	36,300	36,911	37,399	37,960	39,052	38,817
45-64	51,221	55,911	60,158	65,450	70,817	75,069	79,064	83,719	88,614	91,871
65-74	38,202	42,581	46,282	50,503	53,189	56,029	58,707	60,691	62,197	62,993
75+	21,561	24,843	27,708	31,482	34,891	38,825	42,864	46,511	49,773	52,894
Unknown	13	15	*	*	*	*	*	13	17	21
Male	73,856	81,298	87,869	95,552	102,665	109,249	115,527	121,758	128,229	132,326
Female	68,010	75,000	80,677	87,711	94,187	99,208	104,070	108,687	112,915	115,721
Unknown	.	*	*	*	12	18	12	18	17	19
White	81,924	89,621	95,240	103,214	109,594	114,876	121,362	126,980	133,663	138,033
Black	53,548	59,406	64,976	70,459	75,643	80,180	83,959	88,113	91,378	93,217
Native American	1,992	2,337	2,567	2,831	3,215	3,611	3,647	4,088	4,129	4,102
Asian	4,046	4,547	5,288	6,015	6,820	7,403	7,790	8,331	8,902	9,270
Other/unknown	356	388	476	753	1,592	2,405	2,851	2,951	3,089	3,444
†Hispanic					13,605	17,679	20,645	23,370	25,692	27,420
†Non-Hispanic					183,259	190,796	198,964	207,093	215,469	220,646
Diabetes	41,354	47,893	52,328	59,098	66,629	74,385	82,171	89,152	95,114	99,705
Hypertension	40,835	45,912	49,082	53,611	55,496	57,991	60,617	63,539	66,062	67,447
Glomerulonephritis	21,322	22,742	23,930	25,410	26,999	28,150	28,945	29,644	30,381	29,892
Cystic kidney	5,743	6,004	6,187	6,499	6,774	6,822	6,838	7,006	7,238	7,157
Other urologic	2,760	3,098	3,209	3,391	3,653	3,998	4,266	4,437	4,683	4,714
Other cause	15,603	16,311	16,686	17,528	19,199	20,276	21,359	22,341	23,297	23,638
Unknown cause	6,656	7,084	7,174	7,471	7,618	7,927	8,180	8,503	8,798	8,941
Missing cause	7,593	7,255	9,951	10,264	10,496	8,926	7,233	5,841	5,588	6,572
All	141,866	156,299	168,547	183,272	196,864	208,475	219,609	230,463	241,161	248,066

The point prevalent cohorts and associated modalities are determined on December 31 of each calendar year without applying the 60-day stable modality rule.

The age of the point prevalent patient is determined as of December 31 of each calendar year.

†CMS has only started collecting Hispanic ethnicity data as of April, 1995, so Hispanic data in years prior to 1995 has been left blank.

\* Values for cells with fewer than 10 patients are suppressed.

." Zero values in this cell.

**Table D.3 (continued)**  
**Prevalent ESRD Patients, by Dialysis/Functioning Graft Status**

*point prevalent patients on December 31, by demographic characteristics & primary diagnosis*

HEMODIALYSIS	1991	1992	1993	1994	1995	1996	1997	1998	1999	2000
0-4	24	24	32	38	45	39	30	30	33	49
5-9	49	43	48	51	55	61	63	58	59	68
10-14	127	137	128	152	152	154	139	156	155	149
15-19	537	555	566	519	578	585	597	626	595	544
20-24	1,686	1,763	1,765	1,830	1,845	1,865	1,869	1,939	1,941	1,949
25-29	2,988	3,105	3,103	3,169	3,293	3,481	3,605	3,795	3,899	3,720
30-34	4,556	4,777	4,906	5,067	5,313	5,448	5,545	5,551	5,745	5,803
35-39	5,934	6,455	6,703	7,080	7,415	7,797	8,081	8,415	8,750	8,734
40-44	7,376	7,973	8,379	8,687	9,452	10,243	10,963	11,647	12,078	12,132
45-49	8,023	8,839	9,617	10,580	11,768	13,053	13,756	14,665	15,625	16,124
50-54	9,038	10,048	10,917	11,911	12,881	14,145	15,503	17,007	18,610	19,900
55-59	11,209	12,147	12,923	13,826	14,866	16,153	17,586	19,166	20,389	21,194
60-64	14,065	14,850	15,746	16,578	17,859	19,087	20,419	21,858	22,916	23,787
65-69	17,152	18,980	20,220	21,689	22,761	24,124	25,541	26,755	27,816	28,604
70-74	15,777	17,744	19,338	21,485	23,009	24,575	26,481	27,885	28,758	29,398
75-79	11,115	12,545	13,740	15,554	17,264	19,310	21,432	23,325	25,071	26,246
80-84	5,826	6,949	7,842	8,864	9,861	10,891	12,159	13,328	14,283	15,912
85+	2,358	2,824	3,319	3,794	4,319	4,982	5,639	6,546	7,335	7,963
Missing/unknown	12	13	*	*	*	*	*	13	17	21
0-19	737	759	774	760	830	839	829	870	842	810
20-44	22,540	24,073	24,856	25,833	27,318	28,834	30,063	31,347	32,413	32,338
45-64	42,335	45,884	49,203	52,895	57,374	62,438	67,264	72,696	77,540	81,005
65-74	32,929	36,724	39,558	43,174	45,770	48,699	52,022	54,640	56,574	58,002
75+	19,299	22,318	24,901	28,212	31,444	35,183	39,230	43,199	46,689	50,121
Unknown	12	13	*	*	*	*	*	13	17	21
Male	61,101	67,296	72,423	78,477	84,720	92,403	100,050	107,564	114,149	118,820
Female	56,751	62,475	66,876	72,398	78,015	83,588	89,358	95,188	99,909	103,458
Unknown	.	.	.	.	*	*	*	13	17	19
White	65,156	71,142	75,349	81,575	87,058	93,407	101,300	108,762	115,751	121,237
Black	47,237	52,519	57,042	61,416	66,193	71,464	76,041	80,716	84,273	86,259
Native American	1,668	1,966	2,164	2,400	2,736	3,135	3,202	3,663	3,729	3,722
Asian	3,481	3,834	4,374	4,925	5,506	6,079	6,534	7,121	7,675	8,072
Other/unknown	310	310	370	559	1,245	1,910	2,336	2,503	2,647	3,007
†Hispanic					11,460	15,008	17,790	20,522	22,889	24,845
†Non-Hispanic					151,278	160,987	171,623	182,243	191,186	197,452
Diabetes	34,269	39,759	43,522	49,289	55,991	63,625	71,774	79,512	85,815	91,269
Hypertension	35,114	39,625	42,166	46,023	47,934	50,871	53,996	57,480	60,233	62,092
Glomerulonephritis	16,827	17,862	18,624	19,659	20,855	22,222	23,404	24,510	25,409	25,381
Cystic kidney	4,588	4,715	4,814	5,053	5,233	5,404	5,573	5,862	6,041	6,115
Other urologic	2,383	2,621	2,693	2,844	3,098	3,412	3,690	3,900	4,144	4,237
Other cause	12,577	13,071	13,224	13,909	15,305	16,571	17,901	19,095	20,038	20,739
Unknown cause	5,586	5,960	5,981	6,223	6,371	6,738	7,037	7,470	7,821	8,052
Missing cause	6,508	6,158	8,275	7,875	7,951	7,152	6,038	4,936	4,574	4,412
All	117,852	129,771	139,299	150,875	162,738	175,995	189,413	202,765	214,075	222,297

The point prevalent cohorts and associated modalities are determined on December 31 of each calendar year without applying the 60-day stable modality rule.

The age of the point prevalent patient is determined as of December 31 of each calendar year.

†CMS has only started collecting Hispanic ethnicity data as of April, 1995, so Hispanic data in years prior to 1995 has been left blank.

\* Values for cells with fewer than 10 patients are suppressed.

." Zero values in this cell.



**Table D.3 (continued)**  
**Prevalent ESRD Patients, by Dialysis/Functioning Graft Status**

*point prevalent patients on December 31, by demographic characteristics & primary diagnosis*

PERITONEAL DIALYSIS	1991	1992	1993	1994	1995	1996	1997	1998	1999	2000
0-4	83	86	77	93	106	117	102	102	95	108
5-9	71	66	72	77	85	82	84	60	64	73
10-14	153	151	159	149	158	165	172	166	127	136
15-19	277	282	296	277	281	307	287	261	225	223
20-24	446	515	561	558	547	528	479	436	429	379
25-29	796	848	896	916	949	922	877	800	702	672
30-34	1,264	1,365	1,511	1,523	1,518	1,436	1,260	1,180	1,106	1,000
35-39	1,519	1,705	1,832	1,911	1,939	1,825	1,693	1,503	1,480	1,374
40-44	1,804	1,947	2,089	2,175	2,370	2,313	2,181	1,908	1,811	1,688
45-49	1,799	2,038	2,268	2,459	2,667	2,649	2,454	2,312	2,105	2,040
50-54	1,758	2,067	2,274	2,575	2,710	2,784	2,532	2,362	2,404	2,328
55-59	1,880	2,085	2,308	2,469	2,719	2,721	2,657	2,467	2,389	2,255
60-64	2,185	2,497	2,533	2,672	2,934	2,844	2,643	2,325	2,215	2,122
65-69	2,473	2,748	3,096	3,188	3,256	3,139	2,847	2,579	2,486	2,276
70-74	1,867	2,200	2,547	2,757	2,753	2,681	2,499	2,383	2,199	2,022
75-79	1,103	1,292	1,407	1,545	1,646	1,635	1,642	1,519	1,445	1,395
80-84	452	522	599	719	760	727	715	656	623	635
85+	139	176	200	208	231	251	259	244	247	220
Missing/unknown	*	*	*	*	.	.	.	.	.	.
0-19	584	585	604	596	630	671	645	589	511	540
20-44	5,829	6,380	6,889	7,083	7,323	7,024	6,490	5,827	5,528	5,113
45-64	7,622	8,687	9,383	10,175	11,030	10,998	10,286	9,466	9,113	8,745
65-74	4,340	4,948	5,643	5,945	6,009	5,820	5,346	4,962	4,685	4,298
75+	1,694	1,990	2,206	2,472	2,637	2,613	2,616	2,419	2,315	2,250
Unknown	*	*	*	*	.	.	.	.	.	.
Male	10,631	11,917	13,041	13,721	14,390	14,043	13,034	11,872	11,292	10,767
Female	9,439	10,674	11,685	12,550	13,239	13,083	12,349	11,391	10,860	10,179
Unknown	.	*	*	*	.	.	.	.	.	.
White	14,152	15,863	17,003	17,869	18,589	17,977	16,826	15,373	14,634	13,818
Black	5,164	5,795	6,611	7,101	7,353	7,210	6,680	6,124	5,833	5,456
Native American	279	322	336	364	404	415	388	379	343	314
Asian	445	573	732	876	1,069	1,151	1,076	1,022	1,001	1,017
Other/unknown	30	39	45	62	214	373	413	365	341	341
†Hispanic					1,779	2,226	2,308	2,284	2,325	2,276
†Non-Hispanic					25,850	24,900	23,075	20,979	19,827	18,670
Diabetes	5,853	6,913	7,352	8,034	8,849	9,091	8,686	8,004	7,707	7,336
Hypertension	4,787	5,443	5,965	6,426	6,416	6,037	5,581	5,133	4,956	4,703
Glomerulonephritis	3,909	4,311	4,670	4,942	5,202	5,223	4,964	4,519	4,206	3,937
Cystic kidney	1,014	1,122	1,195	1,224	1,270	1,251	1,124	1,032	996	914
Other urologic	297	375	428	455	458	477	482	450	440	422
Other cause	2,551	2,738	2,975	3,012	3,227	3,131	2,942	2,761	2,637	2,463
Unknown cause	882	941	1,031	1,045	1,031	1,019	976	905	800	748
Missing cause	777	749	1,111	1,134	1,176	897	628	459	410	423
All	20,070	22,592	24,727	26,272	27,629	27,126	25,383	23,263	22,152	20,946

The point prevalent cohorts and associated modalities are determined on December 31 of each calendar year without applying the 60-day stable modality rule.

†CMS has only started collecting Hispanic ethnicity data as of April, 1995, so Hispanic data in years prior to 1995 has been left blank.

\* Values for cells with fewer than 10 patients are suppressed.

." Zero values in this cell.

**Table D.3 (continued)**  
**Prevalent ESRD Patients, by Dialysis/Functioning Graft Status**

*point prevalent patients on December 31, by demographic characteristics & primary diagnosis*

TRANSPLANT	1991	1992	1993	1994	1995	1996	1997	1998	1999	2000
0-4	130	140	154	147	136	143	140	138	142	136
5-9	348	345	354	341	365	380	408	420	429	364
10-14	476	508	532	521	553	618	629	674	708	713
15-19	851	840	892	974	1,053	1,051	1,091	1,094	1,156	1,133
20-24	1,687	1,760	1,772	1,703	1,663	1,695	1,693	1,735	1,782	1,793
25-29	3,071	3,094	3,196	3,128	3,200	3,277	3,366	3,372	3,333	3,221
30-34	4,594	4,839	5,173	5,186	5,303	5,349	5,374	5,503	5,450	5,469
35-39	5,405	5,816	6,221	6,454	6,715	7,100	7,346	7,639	7,720	7,769
40-44	5,545	6,020	6,623	6,906	7,303	7,736	8,226	8,694	9,115	9,339
45-49	4,669	5,353	6,070	6,613	7,228	7,935	8,375	8,912	9,250	9,848
50-54	3,974	4,473	5,074	5,426	5,957	6,700	7,697	8,481	9,332	10,044
55-59	3,356	3,803	4,240	4,556	4,905	5,421	6,187	7,003	7,637	8,152
60-64	2,700	3,000	3,343	3,551	3,945	4,514	5,017	5,563	6,089	6,585
65-69	1,658	2,076	2,539	2,956	3,366	3,859	4,360	4,837	5,320	5,859
70-74	540	726	922	1,160	1,392	1,643	1,993	2,356	2,649	3,129
75-79	110	129	170	221	321	411	523	663	812	964
80-84	19	18	25	36	38	62	72	96	137	173
85+	*	*	*	*	*	11	14	23	30	34
Missing/unknown	*	.	*	.	.	.	.	.	.	.
0-19	1,805	1,833	1,932	1,983	2,107	2,192	2,268	2,326	2,435	2,346
20-44	20,302	21,529	22,985	23,377	24,184	25,157	26,005	26,943	27,400	27,591
45-64	14,699	16,629	18,727	20,146	22,035	24,570	27,276	29,959	32,308	34,629
65-74	2,198	2,802	3,461	4,116	4,758	5,502	6,353	7,193	7,969	8,988
75+	137	151	198	260	366	484	609	782	979	1,171
Unknown	*	.	*	.	.	.	.	.	.	.
Male	23,930	26,129	28,713	30,400	32,610	35,156	37,911	40,563	42,727	44,896
Female	15,212	16,815	18,591	19,482	20,840	22,749	24,600	26,640	28,364	29,829
Unknown	.	.	.	.	.	.	.	.	.	.
White	29,995	32,813	36,083	37,833	40,564	43,680	46,771	49,972	52,654	55,172
Black	7,121	7,840	8,686	9,342	9,962	10,953	12,109	13,160	13,988	14,747
Native American	479	541	597	649	693	751	804	863	928	972
Asian	1,343	1,537	1,704	1,814	1,963	2,189	2,411	2,662	2,856	3,052
Other/unknown	204	213	234	244	268	332	416	546	665	782
†Hispanic					3,177	3,650	4,246	4,994	5,645	6,301
†Non-Hispanic					50,273	54,255	58,265	62,209	65,446	68,424
Diabetes	8,193	9,045	10,311	11,248	12,123	13,315	14,511	15,749	16,730	17,726
Hypertension	5,345	6,004	6,716	7,318	7,923	8,678	9,389	10,081	10,628	11,108
Glomerulonephritis	10,616	11,610	12,576	13,030	13,866	14,904	16,042	17,203	18,183	19,037
Cystic kidney	2,953	3,351	3,683	3,877	4,207	4,598	5,036	5,465	5,873	6,217
Other urologic	630	728	885	938	1,042	1,142	1,263	1,409	1,521	1,586
Other cause	6,054	6,619	7,200	7,352	7,792	8,406	8,994	9,677	10,224	10,673
Unknown cause	1,742	1,824	1,975	2,045	2,148	2,293	2,500	2,728	2,889	3,078
Missing cause	3,609	3,763	3,958	4,074	4,349	4,569	4,776	4,891	5,043	5,300
All	39,142	42,944	47,304	49,882	53,450	57,905	62,511	67,203	71,091	74,725

The point prevalent cohorts and associated modalities are determined on December 31 of each calendar year without applying the 60-day stable modality rule.

The age of the point prevalent patient is determined as of December 31 of each calendar year.

†CMS has only started collecting Hispanic ethnicity data as of April, 1995, so Hispanic data in years prior to 1995 has been left blank.

\* Values for cells with fewer than 10 patients are suppressed.

." Zero values in this cell.

**Table D.11  
Incident ESRD Patients for Selected Years: by Diabetic Status & State/Territory**

1992 ALL	Center hemo	Center self hemo	Home hemo	CAPD	CCPD	Other PD	Uncertain dialysis	Transplant	All
Alabama	942	*	.	109	*	20	*	32	1,120
Alaska	33	.	.	*	*	*	.	*	42
Arizona	820	.	.	104	*	*	26	*	961
Arkansas	383	.	.	85	*	*	27	*	512
California	4,228	*	*	420	14	*	1,049	90	5,814
Colorado	405	*	*	53	*	*	18	11	497
Connecticut	445	.	*	142	21	20	19	15	663
Delaware	158	.	.	30	.	.	*	*	195
District of Columbia	286	.	.	10	.	*	*	*	309
Florida	3,112	*	.	243	45	30	121	50	3,602
Georgia	1,576	.	.	182	12	25	47	29	1,871
Hawaii	259	.	.	*	.	.	*	*	281
Idaho	81	.	.	22	*	.	10	*	122
Illinois	2,234	*	*	234	14	28	90	42	2,647
Indiana	851	.	.	197	*	*	71	21	1,148
Iowa	362	*	.	69	*	*	48	20	512
Kansas	329	.	.	71	*	*	16	12	433
Kentucky	606	*	.	89	*	16	25	14	756
Louisiana	1,151	.	*	97	*	*	30	14	1,296
Maine	126	.	.	14	*	.	*	*	148
Maryland	1,056	.	*	126	*	*	14	19	1,228
Massachusetts	979	.	.	102	*	15	20	15	1,137
Michigan	1,606	*	*	321	13	17	27	54	2,042
Minnesota	576	*	.	34	*	14	43	24	695
Mississippi	634	.	*	50	*	*	27	*	727
Missouri	863	*	.	127	*	*	107	20	1,138
Montana	94	.	.	11	*	*	*	*	120
Nebraska	160	.	*	76	*	*	19	19	284
Nevada	186	*	.	26	*	*	12	*	236
New Hampshire	121	.	.	15	.	*	*	*	149
New Jersey	1,389	*	*	265	22	34	267	13	1,995
New Mexico	253	.	.	38	*	*	21	*	321
New York	3,195	*	.	500	18	72	64	39	3,892
North Carolina	1,269	*	.	226	70	56	136	13	1,773
North Dakota	102	.	.	10	.	.	*	*	122
Ohio	1,833	*	.	251	13	86	98	65	2,347
Oklahoma	472	.	*	74	.	55	26	15	644
Oregon	276	.	.	65	*	17	95	14	470
Pennsylvania	2,379	*	.	219	11	37	140	51	2,841
Rhode Island	184	.	.	10	*	.	.	*	199
South Carolina	760	*	*	97	*	11	123	16	1,022
South Dakota	82	.	*	*	.	.	17	*	110
Tennessee	825	.	*	107	*	*	224	27	1,197
Texas	3,005	*	.	218	26	24	747	51	4,076
Utah	158	.	.	13	.	*	*	17	192
Vermont	50	.	.	*	.	.	*	*	61
Virginia	1,322	.	.	156	*	15	14	29	1,542
Washington	621	18	*	48	*	11	20	16	741
West Virginia	296	.	.	65	*	*	.	*	374
Wisconsin	671	*	.	105	*	11	97	44	936
Wyoming	26	*	.	10	.	*	.	*	41
American Samoa	12	.	.	.	.	.	*	.	13
Guam	37	.	.	.	.	.	*	.	38
Puerto Rico	631	.	.	67	.	*	*	*	710
Saipan	*	.	.	.	.	.	.	.	*
Virgin Islands	21	.	.	*	.	.	.	.	23
Foreign	*	.	.	*	.	.	.	*	11
Unknown	29	.	.	*	.	.	*	.	35
All	44,577	63	31	5,631	389	712	4,011	1,006	56,420

The incident cohorts and associated modalities are determined at the time of ESRD initiation without applying the 60-day stable modality rule.

\* Values for cells with fewer than 10 patients are suppressed.

." Zero values in this cell.

**Table D.11 (continued)**  
**Incident ESRD Patients for Selected Years: by Diabetic Status & State/Territory**

1992 DIABETIC	Center hemo	Center self hemo	Home hemo	CAPD	CCPD	Other PD	Uncertain		All
							dialysis	Transplant	
Alabama	316	.	.	46	*	10	*	*	380
Alaska	13	.	.	*	.	.	.	.	14
Arizona	405	.	.	44	*	*	12	*	464
Arkansas	123	.	.	38	.	*	*	*	170
California	1,653	*	*	159	*	*	437	12	2,271
Colorado	164	*	*	24	*	.	*	*	204
Connecticut	151	.	*	54	*	*	*	*	232
Delaware	48	.	.	*	.	.	.	.	55
District of Columbia	99	.	.	*	.	*	*	.	106
Florida	952	*	.	94	13	*	41	12	1,119
Georgia	532	.	.	73	*	10	16	*	638
Hawaii	137	.	.	*	.	.	*	.	146
Idaho	37	.	.	12	*	.	*	*	59
Illinois	674	.	.	83	*	11	22	13	808
Indiana	309	.	.	68	*	*	16	*	403
Iowa	128	.	.	32	.	*	13	12	186
Kansas	112	.	.	30	*	.	*	.	148
Kentucky	195	.	.	26	*	*	*	*	240
Louisiana	436	.	*	37	*	.	15	*	494
Maine	44	.	.	*	*	.	*	.	50
Maryland	397	.	.	47	*	*	*	*	458
Massachusetts	328	.	.	54	*	*	*	*	397
Michigan	624	.	.	148	*	*	*	19	808
Minnesota	206	*	.	16	.	*	17	*	253
Mississippi	225	.	.	23	*	*	*	*	258
Missouri	318	*	.	48	*	*	35	*	417
Montana	49	.	.	*	*	*	.	*	63
Nebraska	58	.	.	31	*	*	*	13	113
Nevada	57	.	.	12	*	.	*	*	77
New Hampshire	35	.	.	*	.	*	.	*	42
New Jersey	453	.	.	91	*	12	73	*	640
New Mexico	146	.	.	26	*	*	*	.	185
New York	1,061	*	.	196	*	25	18	*	1,311
North Carolina	447	*	.	92	27	18	39	.	624
North Dakota	52	.	.	*	.	.	*	.	60
Ohio	691	.	.	108	*	41	30	10	883
Oklahoma	227	.	*	32	.	31	15	*	308
Oregon	104	.	.	30	*	*	33	.	171
Pennsylvania	863	*	.	97	*	17	36	*	1,027
Rhode Island	61	.	.	*	.	.	.	*	65
South Carolina	265	*	*	36	*	*	50	*	365
South Dakota	46	.	*	*	.	.	*	*	56
Tennessee	309	.	.	34	*	*	62	*	418
Texas	1,259	*	.	92	*	13	317	*	1,699
Utah	68	.	.	*	.	.	*	*	82
Vermont	12	.	.	*	.	.	*	.	16
Virginia	428	.	.	59	*	*	*	*	505
Washington	254	*	.	18	*	*	10	10	305
West Virginia	119	.	.	32	*	*	.	.	153
Wisconsin	268	.	.	47	*	*	35	16	373
Wyoming	*	.	.	*	.	*	.	.	*
American Samoa	*	.	.	.	.	.	*	.	*
Guam	24	.	.	.	.	.	.	.	24
Puerto Rico	351	.	.	33	.	*	*	*	390
Saipan	*	.	.	.	.	.	.	.	*
Virgin Islands	*	.	.	*	.	.	.	.	*
Foreign	*	.	.	*	.	.	.	.	*
Unknown	13	.	.	*	.	.	.	.	14
All	16,368	19	10	2,272	148	285	1,452	221	20,775

The incident cohorts and associated modalities are determined at the time of ESRD initiation without applying the 60-day stable modality rule.  
 \* Values for cells with fewer than 10 patients are suppressed.  
 ." Zero values in this cell.

**Table D.11 (continued)**  
**Incident ESRD Patients for Selected Years: by Diabetic Status & State/Territory**

1992 NON-DIABETIC	Center hemo	Center self hemo	Home hemo	CAPD	CCPD	Other PD	Uncertain dialysis	Transplant	All
Alabama	626	*	.	63	*	10	*	29	740
Alaska	20	.	.	*	*	*	.	*	28
Arizona	415	.	.	60	.	*	14	*	497
Arkansas	260	.	.	47	*	*	23	*	342
California	2,575	*	*	261	11	*	612	78	3,543
Colorado	241	.	.	29	*	*	*	*	293
Connecticut	294	.	.	88	15	11	12	11	431
Delaware	110	.	.	23	.	.	*	*	140
District of Columbia	187	.	.	*	.	.	*	*	203
Florida	2,160	.	.	149	32	24	80	38	2,483
Georgia	1,044	.	.	109	*	15	31	26	1,233
Hawaii	122	.	.	*	.	.	*	*	135
Idaho	44	.	.	10	*	.	*	*	63
Illinois	1,560	*	*	151	*	17	68	29	1,839
Indiana	542	.	.	129	*	*	55	16	745
Iowa	234	*	.	37	*	*	35	*	326
Kansas	217	.	.	41	.	*	12	12	285
Kentucky	411	*	.	63	*	10	20	10	516
Louisiana	715	.	.	60	*	*	15	10	802
Maine	82	.	.	11	.	.	*	*	98
Maryland	659	.	*	79	*	*	10	13	770
Massachusetts	651	.	.	48	*	12	11	13	740
Michigan	982	*	*	173	*	13	22	35	1,234
Minnesota	370	*	.	18	*	*	26	17	442
Mississippi	409	.	*	27	*	*	20	*	469
Missouri	545	*	.	79	*	*	72	15	721
Montana	45	.	.	*	.	*	*	*	57
Nebraska	102	.	*	45	*	*	12	*	171
Nevada	129	*	.	14	*	*	*	*	159
New Hampshire	86	.	.	12	.	*	*	*	107
New Jersey	936	*	*	174	13	22	194	11	1,355
New Mexico	107	.	.	12	.	.	12	*	136
New York	2,134	*	.	304	12	47	46	35	2,581
North Carolina	822	*	.	134	43	38	97	13	1,149
North Dakota	50	.	.	*	.	.	*	*	62
Ohio	1,142	*	.	143	10	45	68	55	1,464
Oklahoma	245	.	*	42	.	24	11	13	336
Oregon	172	.	.	35	*	15	62	14	299
Pennsylvania	1,516	*	.	122	*	20	104	43	1,814
Rhode Island	123	.	.	*	*	.	.	*	134
South Carolina	495	.	*	61	*	*	73	13	657
South Dakota	36	.	*	*	.	.	12	*	54
Tennessee	516	.	*	73	*	*	162	22	779
Texas	1,746	*	.	126	17	11	430	43	2,377
Utah	90	.	.	*	.	*	*	10	110
Vermont	38	.	.	*	.	.	*	*	45
Virginia	894	.	.	97	*	*	10	25	1,037
Washington	367	12	*	30	*	*	10	*	436
West Virginia	177	.	.	33	*	*	.	*	221
Wisconsin	403	*	.	58	*	*	62	28	563
Wyoming	20	*	.	*	.	*	.	*	32
American Samoa	*	.	.	.	.	.	.	.	*
Guam	13	.	.	.	.	.	*	.	14
Puerto Rico	280	.	.	34	.	*	*	*	320
Saipan	*	.	.	.	.	.	.	.	*
Virgin Islands	16	.	.	*	.	.	.	.	17
Foreign	*	.	.	.	.	.	.	*	*
Unknown	16	.	.	*	.	.	*	.	21
All	28,209	44	21	3,359	241	427	2,559	785	35,645

The incident cohorts and associated modalities are determined at the time of ESRD initiation without applying the 60-day stable modality rule.

\* Values for cells with fewer than 10 patients are suppressed.

." Zero values in this cell.

**Table D.11 (continued)**  
**Incident ESRD Patients for Selected Years: by Diabetic Status & State/Territory**

1996	Center	Center	Home				Uncertain		
ALL	hemo	self hemo	hemo	CAPD	CCPD	Other PD	dialysis	Transplant	All
Alabama	1,157	.	*	123	33	*	15	21	1,361
Alaska	50	.	.	*	.	.	.	*	57
Arizona	909	.	*	95	12	.	57	11	1,093
Arkansas	543	*	*	79	19	*	19	*	672
California	5,353	*	*	559	108	22	892	68	7,010
Colorado	456	*	*	57	12	*	10	14	554
Connecticut	620	.	*	137	20	*	49	11	841
Delaware	177	.	*	15	*	.	*	*	204
District of Columbia	291	.	*	*	*	.	*	.	309
Florida	3,288	.	502	240	130	35	263	45	4,503
Georgia	1,602	.	94	200	65	13	207	22	2,203
Hawaii	302	.	*	32	*	.	10	*	353
Idaho	143	.	.	29	*	.	19	10	203
Illinois	2,646	.	14	229	72	13	112	44	3,130
Indiana	1,075	*	*	199	*	*	91	18	1,398
Iowa	406	*	*	63	15	.	47	19	552
Kansas	401	*	.	82	16	.	*	*	517
Kentucky	689	.	*	69	38	13	62	14	886
Louisiana	1,351	.	*	59	22	.	15	12	1,462
Maine	179	.	*	13	*	.	*	*	201
Maryland	1,366	.	15	105	41	*	60	26	1,615
Massachusetts	1,156	.	*	73	32	*	34	15	1,317
Michigan	2,051	.	*	291	18	*	63	38	2,476
Minnesota	635	*	*	30	15	*	32	31	752
Mississippi	794	.	*	79	16	*	14	*	919
Missouri	1,171	.	*	149	57	*	*	14	1,406
Montana	112	*	.	14	*	*	*	*	137
Nebraska	219	.	.	75	10	*	14	13	333
Nevada	261	.	*	33	*	.	21	*	334
New Hampshire	127	.	.	20	*	.	*	*	158
New Jersey	1,723	*	*	166	158	14	240	23	2,328
New Mexico	315	.	.	56	20	*	25	*	427
New York	4,058	15	19	340	103	49	71	40	4,695
North Carolina	1,497	.	58	154	137	52	259	13	2,170
North Dakota	97	.	.	*	*	.	*	12	124
Ohio	2,323	.	*	294	118	72	139	57	3,012
Oklahoma	633	*	*	52	20	28	26	11	772
Oregon	376	.	*	72	*	*	68	*	534
Pennsylvania	2,989	.	*	167	79	28	170	42	3,478
Rhode Island	222	.	.	10	*	.	12	*	250
South Carolina	882	.	51	68	41	*	164	21	1,229
South Dakota	121	.	.	11	14	.	30	*	178
Tennessee	1,221	*	.	82	49	10	33	21	1,417
Texas	3,968	*	*	226	109	13	550	28	4,909
Utah	200	.	.	18	*	*	*	12	241
Vermont	74	.	*	13	*	.	10	*	100
Virginia	1,570	.	12	110	42	30	58	30	1,852
Washington	703	24	*	73	16	*	13	27	865
West Virginia	418	.	*	30	28	*	14	*	503
Wisconsin	761	.	*	80	43	*	114	29	1,039
Wyoming	46	.	.	*	*	.	*	*	61
American Samoa	10	.	.	.	.	.	.	.	10
Guam	32	.	.	*	.	*	*	*	45
Puerto Rico	684	.	*	91	52	14	11	*	855
Saipan	12	.	.	.	.	.	.	.	12
Virgin Islands	26	.	.	*	.	.	*	.	28
Foreign	*	*	*	*	.	.	.	.	13
Unknown	*	.	.	.	.	.	*	.	*
All	54,501	64	858	5,294	1,839	487	4,167	897	68,107

The incident cohorts and associated modalities are determined at the time of ESRD initiation without applying the 60-day stable modality rule.

\* Values for cells with fewer than 10 patients are suppressed.

." Zero values in this cell.

**Table D.11 (continued)**  
**Incident ESRD Patients for Selected Years: by Diabetic Status & State/Territory**

<b>1996 DIABETIC</b>	<b>Center hemo</b>	<b>Center self hemo</b>	<b>Home hemo</b>	<b>CAPD</b>	<b>CCPD</b>	<b>Other PD</b>	<b>Uncertain dialysis</b>	<b>Transplant</b>	<b>All</b>
Alabama	458	.	*	53	11	*	*	*	539
Alaska	25	.	.	*	.	.	.	.	28
Arizona	438	.	*	47	*	.	25	*	525
Arkansas	220	.	*	44	*	.	11	*	285
California	2,389	*	.	261	44	*	396	17	3,117
Colorado	233	*	*	28	*	*	*	*	277
Connecticut	234	.	.	54	*	*	23	*	324
Delaware	61	.	*	*	*	.	.	*	71
District of Columbia	125	.	.	*	*	.	.	.	128
Florida	1,216	.	184	104	56	*	72	*	1,646
Georgia	675	.	39	91	29	*	105	*	950
Hawaii	188	.	.	18	*	.	*	.	213
Idaho	60	.	.	13	.	.	*	*	84
Illinois	1,037	.	*	107	28	*	44	12	1,242
Indiana	458	*	*	86	*	*	39	*	593
Iowa	174	*	*	26	*	.	23	*	234
Kansas	169	*	.	34	*	.	*	*	214
Kentucky	285	.	*	30	18	*	32	*	373
Louisiana	592	.	*	37	*	.	*	*	643
Maine	65	.	.	*	*	.	*	.	70
Maryland	551	.	*	51	18	*	10	11	650
Massachusetts	456	.	*	31	18	*	16	*	528
Michigan	924	.	.	134	14	*	28	12	1,118
Minnesota	251	*	*	10	*	*	*	*	289
Mississippi	357	.	*	32	*	*	*	.	401
Missouri	506	.	*	71	33	*	.	*	615
Montana	50	.	.	*	*	*	*	*	60
Nebraska	83	.	.	42	*	*	*	*	144
Nevada	85	.	*	10	*	.	12	*	118
New Hampshire	48	.	.	*	*	.	*	.	57
New Jersey	720	*	*	83	74	*	112	*	998
New Mexico	193	.	.	32	12	*	16	*	256
New York	1,708	*	*	123	53	15	23	*	1,939
North Carolina	639	.	30	67	52	22	131	*	946
North Dakota	36	.	.	.	.	.	*	*	42
Ohio	1,011	.	*	156	57	27	53	12	1,318
Oklahoma	307	.	.	21	13	11	*	*	362
Oregon	144	.	.	28	*	*	23	*	201
Pennsylvania	1,211	.	*	67	36	13	61	*	1,395
Rhode Island	85	.	.	*	*	.	*	.	97
South Carolina	371	.	22	31	17	*	76	*	527
South Dakota	64	.	.	*	*	.	*	.	82
Tennessee	482	.	.	35	25	*	16	*	568
Texas	2,029	*	*	111	58	10	256	*	2,476
Utah	100	.	.	11	*	.	*	*	116
Vermont	30	.	*	*	*	.	*	*	40
Virginia	628	.	*	50	12	16	13	*	730
Washington	288	11	.	42	*	*	*	11	364
West Virginia	183	.	.	17	14	*	*	*	222
Wisconsin	312	.	*	33	22	*	41	*	419
Wyoming	23	.	.	*	.	.	.	*	28
American Samoa	*	.	.	.	.	.	.	.	*
Guam	22	.	.	*	.	.	*	.	32
Puerto Rico	405	.	.	47	25	*	*	*	491
Saipan	*	.	.	.	.	.	.	.	*
Virgin Islands	15	.	.	*	.	.	.	.	16
Foreign	*	.	.	.	.	.	.	.	*
Unknown	*	*	.	*	.	.	.	.	*
All	23,435	27	334	2,416	822	206	1,751	228	29,219

The incident cohorts and associated modalities are determined at the time of ESRD initiation without applying the 60-day stable modality rule.

\* Values for cells with fewer than 10 patients are suppressed.

." Zero values in this cell.

**Table D.11 (continued)**  
**Incident ESRD Patients for Selected Years: by Diabetic Status & State/Territory**

1996 NON-DIABETIC	Center hemo	Center self hemo	Home hemo	CAPD	CCPD	Other PD	Uncertain dialysis	Transplant	All
Alabama	699	.	*	70	22	*	11	17	822
Alaska	25	.	.	*	.	.	.	*	29
Arizona	471	.	*	48	*	.	32	*	568
Arkansas	323	*	.	35	12	*	*	*	387
California	2,964	*	*	298	64	13	496	51	3,893
Colorado	223	*	.	29	*	*	*	*	277
Connecticut	386	.	*	83	11	*	26	*	517
Delaware	116	.	*	*	.	.	*	*	133
District of Columbia	166	.	*	*	*	.	*	.	181
Florida	2,072	.	318	136	74	26	191	40	2,857
Georgia	927	.	55	109	36	*	102	17	1,253
Hawaii	114	.	*	14	*	.	*	*	140
Idaho	83	.	.	16	*	.	10	*	119
Illinois	1,609	.	*	122	44	*	68	32	1,888
Indiana	617	.	*	113	*	*	52	14	805
Iowa	232	.	.	37	10	.	24	15	318
Kansas	232	.	.	48	10	.	*	*	303
Kentucky	404	.	.	39	20	*	30	11	513
Louisiana	759	.	*	22	16	.	13	*	819
Maine	114	.	*	11	*	.	*	*	131
Maryland	815	.	*	54	23	*	50	15	965
Massachusetts	700	.	*	42	14	*	18	11	789
Michigan	1,127	.	*	157	*	*	35	26	1,358
Minnesota	384	.	*	20	10	*	24	22	463
Mississippi	437	.	*	47	13	*	12	*	518
Missouri	665	.	*	78	24	*	*	11	791
Montana	62	*	.	10	*	.	*	*	77
Nebraska	136	.	.	33	*	.	*	*	189
Nevada	176	.	.	23	*	.	*	*	216
New Hampshire	79	.	.	14	*	.	*	*	101
New Jersey	1,003	.	*	83	84	12	128	18	1,330
New Mexico	122	.	.	24	*	*	*	*	171
New York	2,350	11	14	217	50	34	48	32	2,756
North Carolina	858	.	28	87	85	30	128	*	1,224
North Dakota	61	.	.	*	*	.	*	*	82
Ohio	1,312	.	*	138	61	45	86	45	1,694
Oklahoma	326	*	*	31	*	17	19	*	410
Oregon	232	.	*	44	*	.	45	*	333
Pennsylvania	1,778	.	*	100	43	15	109	37	2,083
Rhode Island	137	.	.	*	*	.	*	*	153
South Carolina	511	.	29	37	24	.	88	13	702
South Dakota	57	.	.	*	*	.	22	*	96
Tennessee	739	*	.	47	24	*	17	15	849
Texas	1,939	*	*	115	51	*	294	22	2,433
Utah	100	.	.	*	.	*	*	10	125
Vermont	44	.	.	*	.	.	*	.	60
Virginia	942	.	*	60	30	14	45	23	1,122
Washington	415	13	*	31	12	*	*	16	501
West Virginia	235	.	*	13	14	*	*	*	281
Wisconsin	449	.	*	47	21	*	73	21	620
Wyoming	23	.	.	*	*	.	*	*	33
American Samoa	*	.	.	.	.	.	.	.	*
Guam	10	.	.	*	.	*	.	.	13
Puerto Rico	279	.	*	44	27	*	*	*	364
Saipan	*	.	.	.	.	.	.	.	*
Virgin Islands	11	.	.	.	.	.	*	.	12
Foreign	*	.	*	*	.	.	.	.	*
Unknown	*	.	.	.	.	.	*	.	*
All	31,066	37	524	2,878	1,017	281	2,416	669	38,888

The incident cohorts and associated modalities are determined at the time of ESRD initiation without applying the 60-day stable modality rule.

\* Values for cells with fewer than 10 patients are suppressed.

." Zero values in this cell.



**Table D.11 (continued)**  
**Incident ESRD Patients for Selected Years: by Diabetic Status & State/Territory**

2000	Center	Center	Home				Uncertain		
ALL	hemo	self hemo	hemo	CAPD	CCPD	Other PD	dialysis	Transplant	All
Alabama	1,358	.	.	86	42	.	67	21	1,574
Alaska	52	.	.	15	.	.	*	*	73
Arizona	1,207	.	12	70	11	*	35	17	1,354
Arkansas	664	.	.	66	20	*	33	13	797
California	7,633	*	*	426	110	10	284	84	8,554
Colorado	579	.	*	30	*	.	27	10	652
Connecticut	632	.	.	102	34	*	62	19	851
Delaware	210	.	.	*	*	.	*	*	222
District of Columbia	308	.	*	11	.	*	*	.	327
Florida	4,492	.	51	188	117	14	202	50	5,114
Georgia	2,491	.	*	123	62	*	87	18	2,796
Hawaii	395	.	.	29	*	.	29	*	459
Idaho	188	.	.	34	*	.	*	*	241
Illinois	2,973	.	68	168	85	*	105	52	3,455
Indiana	1,442	.	*	122	28	10	89	16	1,709
Iowa	517	.	*	45	18	.	27	21	631
Kansas	497	*	*	71	17	.	23	*	620
Kentucky	911	.	.	61	18	*	64	12	1,068
Louisiana	1,631	.	*	79	25	*	40	12	1,793
Maine	246	.	.	11	11	.	*	*	276
Maryland	1,489	.	*	79	39	*	58	41	1,717
Massachusetts	1,204	.	*	50	42	*	70	*	1,385
Michigan	2,405	.	*	189	24	*	77	44	2,750
Minnesota	831	.	.	24	13	*	23	35	927
Mississippi	864	.	*	73	44	*	25	*	1,017
Missouri	1,469	.	*	76	39	.	34	19	1,645
Montana	115	*	*	12	*	.	10	*	149
Nebraska	347	.	*	83	*	.	12	10	457
Nevada	388	.	*	22	12	*	20	*	449
New Hampshire	177	.	.	23	*	*	14	*	225
New Jersey	2,297	*	*	76	70	12	95	44	2,599
New Mexico	413	.	.	23	26	.	26	*	493
New York	4,651	*	12	159	74	54	124	57	5,136
North Carolina	2,427	.	*	112	93	44	96	19	2,794
North Dakota	122	.	.	11	*	.	*	*	150
Ohio	3,115	*	*	259	88	*	132	69	3,679
Oklahoma	817	.	*	60	40	11	37	*	974
Oregon	549	.	.	52	14	.	18	10	643
Pennsylvania	3,276	*	*	109	66	*	140	41	3,640
Rhode Island	244	.	.	13	*	.	23	*	284
South Carolina	1,243	.	*	50	69	*	56	16	1,439
South Dakota	183	.	.	10	*	.	11	*	210
Tennessee	1,447	.	.	55	68	*	66	23	1,666
Texas	5,298	.	*	202	188	*	215	49	5,969
Utah	236	.	.	24	*	.	17	13	295
Vermont	87	.	.	*	*	.	18	*	113
Virginia	1,694	*	*	98	28	*	80	29	1,939
Washington	949	12	*	49	22	*	29	16	1,082
West Virginia	532	.	.	39	51	*	24	11	660
Wisconsin	1,046	.	*	69	27	*	49	42	1,238
Wyoming	65	.	.	*	*	.	*	*	73
American Samoa	19	.	.	.	.	.	*	.	20
Guam	51	.	.	*	*	.	.	*	57
Puerto Rico	766	.	*	32	93	*	33	*	931
Saipan	19	.	.	.	.	.	*	.	20
Virgin Islands	22	.	.	*	*	.	*	.	26
Foreign	13	.	.	*	.	.	*	*	19
Unknown	*	.	.	.	.	.	*	.	*
All	69,303	29	226	3,885	1,878	256	2,850	1,017	79,444

The incident cohorts and associated modalities are determined at the time of ESRD initiation without applying the 60-day stable modality rule.  
 \* Values for cells with fewer than 10 patients are suppressed.  
 ." Zero values in this cell.

**Table D.11 (continued)**  
**Incident ESRD Patients for Selected Years: by Diabetic Status & State/Territory**

2000 DIABETIC	Center hemo	Center self hemo	Home hemo	CAPD	CCPD	Other PD	Uncertain		All
							dialysis	Transplant	
Alabama	613	.	.	35	21	.	11	*	683
Alaska	26	.	.	*	.	.	*	.	31
Arizona	644	.	*	32	*	*	*	*	697
Arkansas	280	.	.	36	*	.	*	.	328
California	3,779	*	*	204	32	*	25	13	4,062
Colorado	299	.	.	18	*	.	.	*	320
Connecticut	268	.	.	52	16	.	24	*	364
Delaware	97	.	.	*	*	.	.	.	100
District of Columbia	119	.	*	*	.	.	*	.	123
Florida	1,889	.	23	71	38	*	39	*	2,071
Georgia	1,095	.	*	57	28	*	*	*	1,196
Hawaii	243	.	.	16	.	.	15	*	275
Idaho	84	.	.	15	*	.	*	*	112
Illinois	1,218	.	35	71	43	.	18	10	1,395
Indiana	594	.	*	55	16	*	21	*	695
Iowa	212	.	.	29	*	.	*	*	258
Kansas	239	.	*	41	*	.	*	.	295
Kentucky	400	.	.	34	*	.	23	*	465
Louisiana	713	.	.	33	*	*	.	*	758
Maine	106	.	.	*	*	.	*	*	118
Maryland	598	.	*	29	15	*	*	*	659
Massachusetts	451	.	*	22	18	*	25	*	521
Michigan	1,100	.	*	87	*	*	13	12	1,226
Minnesota	326	.	.	11	*	*	*	11	355
Mississippi	372	.	*	34	15	*	*	*	427
Missouri	682	.	*	31	15	.	*	*	737
Montana	56	*	.	*	*	.	*	*	70
Nebraska	142	.	.	47	*	.	*	*	198
Nevada	187	.	*	12	*	.	.	.	209
New Hampshire	61	.	.	*	*	.	*	.	77
New Jersey	965	.	*	34	36	*	*	*	1,049
New Mexico	261	.	.	13	16	.	10	.	300
New York	1,970	*	*	70	30	26	18	*	2,123
North Carolina	1,104	.	*	39	32	28	15	*	1,228
North Dakota	50	.	.	*	.	.	*	*	60
Ohio	1,494	*	*	139	44	*	39	12	1,733
Oklahoma	406	.	.	41	16	*	*	*	479
Oregon	231	.	.	25	*	.	*	.	269
Pennsylvania	1,414	.	*	45	34	*	48	*	1,550
Rhode Island	73	.	.	*	.	.	*	.	86
South Carolina	579	.	*	21	32	*	*	*	641
South Dakota	81	.	.	*	.	.	*	.	86
Tennessee	658	.	.	23	28	*	13	*	730
Texas	2,883	.	*	109	86	*	36	*	3,128
Utah	119	.	.	18	*	.	*	*	149
Vermont	28	.	.	.	*	.	10	*	40
Virginia	771	.	.	50	15	*	12	*	858
Washington	433	*	.	28	*	*	*	*	481
West Virginia	290	.	.	18	29	*	*	*	349
Wisconsin	430	.	.	24	10	*	11	10	486
Wyoming	36	.	.	*	.	.	*	.	38
American Samoa	16	.	.	.	.	.	.	.	16
Guam	27	.	.	*	.	.	.	.	28
Puerto Rico	475	.	*	22	48	*	.	.	548
Saipan	14	.	.	.	.	.	.	.	14
Virgin Islands	12	.	.	*	.	.	.	.	13
Foreign	*	.	.	.	.	.	.	.	*
Unknown	.	.	.	.	.	.	.	.	.
All	31,716	*	101	1,835	809	119	536	185	35,310

The incident cohorts and associated modalities are determined at the time of ESRD initiation without applying the 60-day stable modality rule.

\* Values for cells with fewer than 10 patients are suppressed.

." Zero values in this cell.

**Table D.11 (continued)**  
**Incident ESRD Patients for Selected Years: by Diabetic Status & State/Territory**

2000 NON-DIABETIC	Center hemo	Center self hemo	Home hemo	CAPD	CCPD	Other PD	Uncertain dialysis	Transplant	All
Alabama	745	.	.	51	21	.	56	18	891
Alaska	26	.	.	11	.	.	*	*	42
Arizona	563	.	*	38	*	.	32	14	657
Arkansas	384	.	.	30	14	*	27	13	469
California	3,854	*	*	222	78	*	259	71	4,492
Colorado	280	.	*	12	*	.	27	*	332
Connecticut	364	.	.	50	18	*	38	15	487
Delaware	113	.	.	*	*	.	*	*	122
District of Columbia	189	.	.	*	.	*	*	.	204
Florida	2,603	.	28	117	79	10	163	43	3,043
Georgia	1,396	.	*	66	34	*	80	16	1,600
Hawaii	152	.	.	13	*	.	14	*	184
Idaho	104	.	.	19	*	.	*	*	129
Illinois	1,755	.	33	97	42	*	87	42	2,060
Indiana	848	.	.	67	12	*	68	14	1,014
Iowa	305	.	*	16	12	.	20	17	373
Kansas	258	*	*	30	*	.	18	*	325
Kentucky	511	.	.	27	12	*	41	10	603
Louisiana	918	.	*	46	17	*	40	*	1,035
Maine	140	.	.	*	*	.	*	.	158
Maryland	891	.	*	50	24	*	50	35	1,058
Massachusetts	753	.	*	28	24	*	45	*	864
Michigan	1,305	.	*	102	16	*	64	32	1,524
Minnesota	505	.	.	13	*	.	21	24	572
Mississippi	492	.	*	39	29	*	23	*	590
Missouri	787	.	*	45	24	.	31	15	908
Montana	59	*	*	*	.	.	*	*	79
Nebraska	205	.	*	36	*	.	*	*	259
Nevada	201	.	*	10	*	*	20	*	240
New Hampshire	116	.	.	14	*	*	*	*	148
New Jersey	1,332	*	*	42	34	10	92	36	1,550
New Mexico	152	.	.	10	10	.	16	*	193
New York	2,681	*	*	89	44	28	106	54	3,013
North Carolina	1,323	.	*	73	61	16	81	11	1,566
North Dakota	72	.	.	*	*	.	*	*	90
Ohio	1,621	*	*	120	44	*	93	57	1,946
Oklahoma	411	.	*	19	24	*	30	*	495
Oregon	318	.	.	27	*	.	13	10	374
Pennsylvania	1,862	*	*	64	32	*	92	36	2,090
Rhode Island	171	.	.	*	*	.	16	*	198
South Carolina	664	.	.	29	37	*	52	14	798
South Dakota	102	.	.	*	*	.	*	*	124
Tennessee	789	.	.	32	40	*	53	21	936
Texas	2,415	.	*	93	102	*	179	43	2,841
Utah	117	.	.	*	*	.	14	*	146
Vermont	59	.	.	*	*	.	*	*	73
Virginia	923	*	*	48	13	*	68	21	1,081
Washington	516	*	*	21	17	*	25	13	601
West Virginia	242	.	.	21	22	*	16	*	311
Wisconsin	616	.	*	45	17	*	38	32	752
Wyoming	29	.	.	*	*	.	*	*	35
American Samoa	*	.	.	.	.	.	*	.	*
Guam	24	.	.	*	*	.	.	*	29
Puerto Rico	291	.	.	10	45	*	33	*	383
Saipan	*	.	.	.	.	.	*	.	*
Virgin Islands	10	.	.	.	*	.	*	.	13
Foreign	10	.	.	*	.	.	*	*	16
Unknown	*	.	.	.	.	.	*	.	*
All	37,587	20	125	2,050	1,069	137	2,314	832	44,134

The incident cohorts and associated modalities are determined at the time of ESRD initiation without applying the 60-day stable modality rule.

\* Values for cells with fewer than 10 patients are suppressed.

." Zero values in this cell.

**Table D.12**  
**Prevalent ESRD Pts for Selected Yrs: by Diabetic Status & State/Territory**  
*point prevalent patients on December 31*

1992	Center	Center	Home				Uncertain		
ALL	hemo	self hemo	hemo	CAPD	CCPD	Other PD	dialysis	Transplant	All
Alabama	2,759	*	*	344	85	*	19	834	4,056
Alaska	70	*	*	21	.	*	*	72	170
Arizona	2,073	21	15	267	86	*	28	588	3,080
Arkansas	1,069	*	*	219	32	*	22	416	1,771
California	13,742	74	37	1,644	269	*	875	4,803	21,452
Colorado	1,082	*	15	178	46	*	35	498	1,858
Connecticut	1,319	*	33	405	80	14	24	461	2,338
Delaware	454	*	*	85	*	.	*	119	667
District of Columbia	925	*	*	30	*	*	*	110	1,077
Florida	8,205	20	24	845	193	33	131	2,409	11,860
Georgia	4,770	*	15	506	144	27	64	1,116	6,646
Hawaii	791	15	*	49	*	.	*	179	1,046
Idaho	240	.	10	79	*	*	12	191	541
Illinois	5,920	57	40	836	96	27	94	2,026	9,096
Indiana	2,321	*	24	719	33	*	40	857	4,005
Iowa	910	*	66	227	35	*	23	520	1,785
Kansas	918	.	25	267	31	*	*	372	1,621
Kentucky	1,452	*	12	339	17	20	23	708	2,573
Louisiana	3,203	*	26	320	72	*	29	667	4,322
Maine	306	.	*	49	16	.	*	212	592
Maryland	2,977	43	23	428	34	12	32	842	4,391
Massachusetts	2,542	15	38	319	46	27	34	969	3,990
Michigan	4,079	*	54	1,139	66	22	42	1,788	7,193
Minnesota	1,567	*	26	149	13	*	25	888	2,674
Mississippi	2,030	*	120	165	*	*	22	362	2,709
Missouri	2,406	*	27	465	91	*	39	950	3,989
Montana	238	.	11	56	*	*	*	153	465
Nebraska	412	*	*	245	26	*	*	292	997
Nevada	434	*	17	74	21	*	11	259	819
New Hampshire	295	.	*	55	26	*	*	165	554
New Jersey	4,191	35	104	793	141	50	233	1,051	6,598
New Mexico	762	*	*	185	15	*	10	261	1,240
New York	9,240	224	139	1,643	154	41	115	2,213	13,769
North Carolina	4,082	17	44	792	257	64	93	984	6,333
North Dakota	204	.	*	28	*	.	*	155	399
Ohio	4,826	*	16	913	82	61	113	2,085	8,099
Oklahoma	1,231	*	16	290	*	18	14	506	2,084
Oregon	844	*	37	290	35	*	78	429	1,717
Pennsylvania	6,400	67	48	813	87	57	168	2,249	9,889
Rhode Island	502	.	.	33	10	.	*	163	712
South Carolina	2,578	61	22	295	59	*	101	653	3,776
South Dakota	205	.	49	19	*	.	*	123	405
Tennessee	2,792	*	57	379	48	*	62	971	4,318
Texas	9,783	55	62	943	257	42	420	2,956	14,518
Utah	384	*	*	49	.	*	10	353	805
Vermont	126	.	12	39	*	.	*	79	258
Virginia	3,700	*	65	551	56	16	36	1,051	5,478
Washington	1,562	17	191	207	28	31	30	782	2,848
West Virginia	641	*	*	171	25	*	*	359	1,218
Wisconsin	1,606	25	30	408	35	*	68	1,220	3,395
Wyoming	63	*	12	39	*	.	*	71	193
American Samoa	30	.	.	.	.	.	*	.	31
Guam	73	*	.	.	.	.	*	13	95
Puerto Rico	1,778	.	16	263	13	*	12	335	2,419
Saipan	15	.	.	.	.	.	*	*	21
Virgin Islands	58	.	.	*	.	.	.	*	71
Foreign	42	.	*	*	*	.	*	26	80
Unknown	95	.	.	*	*	.	10	24	137
All	127,322	811	1,638	19,691	2,901	659	3,277	42,944	199,243

The point prevalent cohorts and associated modalities are determined on December 31 of each calendar year without applying the 60-day stable modality rule.

\* Values for cells with fewer than 10 patients are suppressed.

." Zero values in this cell.

**Table D.12 (continued)**  
**Prevalent ESRD Pts for Selected Yrs: by Diabetic Status & State/Territory**  
*point prevalent patients on December 31*

1992 DIABETIC	Center hemo	Center self hemo	Home hemo	CAPD	CCPD	Other PD	Uncertain		All
							dialysis	Transplant	
Alabama	699	*	*	92	16	*	*	95	909
Alaska	19	*	.	*	.	*	.	13	41
Arizona	937	*	*	90	23	*	*	144	1,213
Arkansas	285	.	.	77	*	.	*	70	438
California	4,695	20	*	507	62	*	338	870	6,503
Colorado	391	.	*	73	12	.	12	162	655
Connecticut	386	*	*	113	27	*	*	92	632
Delaware	139	*	.	17	.	.	.	21	178
District of Columbia	259	.	.	*	.	*	*	16	285
Florida	2,194	*	*	230	49	13	28	472	2,994
Georgia	1,333	*	*	139	37	*	15	213	1,747
Hawaii	363	*	.	23	.	.	*	19	413
Idaho	74	.	*	29	*	.	*	54	165
Illinois	1,517	*	*	240	24	*	20	445	2,263
Indiana	693	.	*	218	*	*	*	165	1,102
Iowa	249	.	10	71	13	*	*	146	496
Kansas	266	.	*	109	*	*	*	77	464
Kentucky	436	.	*	95	*	*	*	111	665
Louisiana	977	.	*	90	24	*	11	119	1,229
Maine	89	.	.	13	*	.	*	33	140
Maryland	899	*	*	128	10	*	*	152	1,210
Massachusetts	674	*	.	97	11	*	*	195	994
Michigan	1,272	.	17	386	14	10	*	419	2,126
Minnesota	500	*	12	47	*	*	10	321	900
Mississippi	547	.	22	54	*	.	*	45	676
Missouri	711	.	*	145	21	*	11	186	1,082
Montana	90	.	*	25	*	*	.	51	175
Nebraska	127	*	*	78	*	.	*	115	336
Nevada	122	.	*	22	*	.	*	62	220
New Hampshire	71	.	*	12	*	*	.	44	136
New Jersey	1,160	*	19	223	51	18	71	175	1,719
New Mexico	381	.	.	91	*	*	*	52	533
New York	2,582	32	12	468	48	11	19	353	3,525
North Carolina	1,197	*	*	232	83	13	26	185	1,748
North Dakota	90	.	.	13	.	.	*	53	160
Ohio	1,481	*	*	297	27	24	33	508	2,375
Oklahoma	491	*	*	117	*	11	*	117	749
Oregon	241	.	*	98	*	*	21	107	485
Pennsylvania	1,901	23	*	265	33	18	34	472	2,752
Rhode Island	129	.	.	*	*	.	*	30	167
South Carolina	789	15	*	78	10	*	33	102	1,034
South Dakota	84	.	32	*	.	.	*	47	170
Tennessee	732	*	*	90	16	.	21	180	1,043
Texas	3,680	12	*	324	73	14	151	616	4,876
Utah	148	.	*	15	.	.	*	134	302
Vermont	39	.	*	14	.	.	.	19	73
Virginia	1,035	.	13	153	16	10	*	179	1,414
Washington	503	*	39	66	11	*	*	239	880
West Virginia	214	.	.	55	*	*	*	87	366
Wisconsin	516	*	*	159	*	*	24	407	1,119
Wyoming	18	.	*	10	.	.	*	20	54
American Samoa	12	.	.	.	.	.	*	.	13
Guam	34	*	.	.	.	.	.	*	39
Puerto Rico	750	.	*	91	*	*	*	31	879
Saipan	*	.	.	.	.	.	*	.	*
Virgin Islands	*	.	.	*	.	.	.	.	*
Foreign	*	.	.	*	.	.	.	*	13
Unknown	42	.	.	*	*	.	.	*	47
All	39,281	158	320	6,112	801	216	1,005	9,045	56,938

The point prevalent cohorts and associated modalities are determined on December 31 of each calendar year without applying the 60-day stable modality rule.

\* Values for cells with fewer than 10 patients are suppressed.

." Zero values in this cell.

**Table D.12 (continued)**  
**Prevalent ESRD Pts for Selected Yrs: by Diabetic Status & State/Territory**  
*point prevalent patients on December 31*

1992 NON-DIABETIC	Center hemo	Center self hemo	Home hemo	CAPD	CCPD	Other PD	Uncertain dialysis	Transplant	All
Alabama	2,060	.	*	252	69	*	16	739	3,147
Alaska	51	.	*	14	.	*	*	59	129
Arizona	1,136	14	*	177	63	*	23	444	1,867
Arkansas	784	*	*	142	29	*	19	346	1,333
California	9,047	54	28	1,137	207	*	537	3,933	14,949
Colorado	691	*	10	105	34	*	23	336	1,203
Connecticut	933	*	29	292	53	11	18	369	1,706
Delaware	315	*	*	68	*	.	*	98	489
District of Columbia	666	*	*	24	*	*	*	94	792
Florida	6,011	15	21	615	144	20	103	1,937	8,866
Georgia	3,437	*	14	367	107	20	49	903	4,899
Hawaii	428	14	*	26	*	.	*	160	633
Idaho	166	.	*	50	*	*	*	137	376
Illinois	4,403	51	34	596	72	22	74	1,581	6,833
Indiana	1,628	*	20	501	24	*	31	692	2,903
Iowa	661	*	56	156	22	*	17	374	1,289
Kansas	652	.	21	158	26	.	*	295	1,157
Kentucky	1,016	*	*	244	10	11	20	597	1,908
Louisiana	2,226	*	20	230	48	*	18	548	3,093
Maine	217	.	*	36	12	.	*	179	452
Maryland	2,078	38	18	300	24	10	23	690	3,181
Massachusetts	1,868	14	38	222	35	19	26	774	2,996
Michigan	2,807	*	37	753	52	12	34	1,369	5,067
Minnesota	1,067	*	14	102	*	.	15	567	1,774
Mississippi	1,483	*	98	111	*	*	17	317	2,033
Missouri	1,695	*	22	320	70	*	28	764	2,907
Montana	148	.	*	31	*	*	*	102	290
Nebraska	285	*	*	167	18	*	*	177	661
Nevada	312	*	*	52	17	*	*	197	599
New Hampshire	224	.	*	43	19	*	*	121	418
New Jersey	3,031	33	85	570	90	32	162	876	4,879
New Mexico	381	*	*	94	10	*	*	209	707
New York	6,658	192	127	1,175	106	30	96	1,860	10,244
North Carolina	2,885	13	36	560	174	51	67	799	4,585
North Dakota	114	.	*	15	*	.	*	102	239
Ohio	3,345	*	12	616	55	37	80	1,577	5,724
Oklahoma	740	*	10	173	*	*	*	389	1,335
Oregon	603	*	28	192	27	*	57	322	1,232
Pennsylvania	4,499	44	42	548	54	39	134	1,777	7,137
Rhode Island	373	.	.	27	*	.	*	133	545
South Carolina	1,789	46	17	217	49	*	68	551	2,742
South Dakota	121	.	17	13	*	.	*	76	235
Tennessee	2,060	*	54	289	32	*	41	791	3,275
Texas	6,103	43	56	619	184	28	269	2,340	9,642
Utah	236	*	*	34	.	*	*	219	503
Vermont	87	.	11	25	*	.	*	60	185
Virginia	2,665	*	52	398	40	*	28	872	4,064
Washington	1,059	10	152	141	17	24	22	543	1,968
West Virginia	427	*	*	116	19	*	*	272	852
Wisconsin	1,090	22	26	249	31	*	44	813	2,276
Wyoming	45	*	*	29	*	.	*	51	139
American Samoa	18	.	.	.	.	.	.	.	18
Guam	39	*	.	.	.	.	*	12	56
Puerto Rico	1,028	.	15	172	10	*	10	304	1,540
Saipan	11	.	.	.	.	.	*	*	14
Virgin Islands	52	.	.	*	.	.	.	*	62
Foreign	34	.	*	*	*	.	*	25	67
Unknown	53	.	.	*	.	.	10	21	90
All	88,041	653	1,318	13,579	2,100	443	2,272	33,899	142,305

The point prevalent cohorts and associated modalities are determined on December 31 of each calendar year without applying the 60-day stable modality rule.

\* Values for cells with fewer than 10 patients are suppressed.

." Zero values in this cell.

**Table D.12 (continued)**  
**Prevalent ESRD Pts for Selected Yrs: by Diabetic Status & State/Territory**  
*point prevalent patients on December 31*

1996	Center	Center	Home				Uncertain		
ALL	hemo	self hemo	hemo	CAPD	CCPD	Other PD	dialysis	Transplant	All
Alabama	3,809	*	14	362	203	17	28	1,172	5,606
Alaska	118	*	*	27	*	.	*	94	249
Arizona	2,839	20	27	207	185	*	65	853	4,200
Arkansas	1,670	.	13	210	53	*	29	638	2,617
California	17,963	46	35	1,815	901	46	922	6,509	28,237
Colorado	1,449	*	12	160	86	*	25	735	2,470
Connecticut	1,848	.	*	297	235	*	80	694	3,166
Delaware	571	*	.	76	25	*	17	184	875
District of Columbia	1,101	*	*	35	10	*	16	115	1,294
Florida	10,569	23	392	787	508	23	323	3,234	15,859
Georgia	6,730	*	87	510	339	14	72	1,512	9,265
Hawaii	1,121	*	*	93	14	.	17	205	1,454
Idaho	345	.	*	89	30	.	16	259	747
Illinois	7,730	*	121	809	324	12	208	2,855	12,061
Indiana	3,254	*	22	678	145	12	84	1,097	5,294
Iowa	1,169	.	31	183	74	*	53	722	2,235
Kansas	1,208	*	16	257	76	*	24	445	2,030
Kentucky	1,927	.	*	220	122	17	75	946	3,316
Louisiana	4,347	.	20	238	191	*	28	944	5,777
Maine	517	.	*	41	32	.	*	265	869
Maryland	4,059	21	48	410	191	10	163	952	5,854
Massachusetts	3,270	*	46	230	189	14	51	1,350	5,156
Michigan	5,909	*	59	1,121	205	11	82	2,386	9,774
Minnesota	1,982	.	*	120	62	*	41	1,248	3,463
Mississippi	2,960	*	68	202	60	*	35	549	3,884
Missouri	3,316	*	67	431	224	*	33	1,284	5,361
Montana	330	.	*	38	10	.	*	235	624
Nebraska	612	.	*	267	33	.	15	367	1,298
Nevada	738	.	10	92	36	*	31	342	1,251
New Hampshire	357	.	*	74	38	.	14	216	702
New Jersey	5,581	77	46	594	361	67	280	1,556	8,562
New Mexico	1,060	.	*	162	88	21	17	393	1,744
New York	12,402	211	103	1,366	479	53	159	3,149	17,922
North Carolina	6,187	*	42	701	427	75	98	1,435	8,968
North Dakota	237	.	.	36	16	.	*	187	484
Ohio	6,529	*	75	1,028	384	51	164	2,673	10,905
Oklahoma	1,852	*	11	217	80	11	24	706	2,904
Oregon	1,192	*	47	250	75	*	44	608	2,219
Pennsylvania	8,458	36	63	620	321	29	334	2,960	12,821
Rhode Island	636	.	.	51	28	.	17	226	958
South Carolina	3,688	*	23	297	197	*	74	945	5,233
South Dakota	387	.	*	39	22	.	*	152	610
Tennessee	3,911	18	42	297	191	*	72	1,292	5,830
Texas	13,865	33	80	947	655	13	441	3,979	20,013
Utah	546	.	*	60	20	*	12	494	1,139
Vermont	186	.	*	46	12	.	*	120	375
Virginia	5,040	*	66	446	236	12	163	1,087	7,053
Washington	2,121	18	144	267	81	10	31	1,019	3,691
West Virginia	912	*	*	115	86	*	49	342	1,522
Wisconsin	2,231	*	39	310	168	*	71	1,625	4,454
Wyoming	126	.	*	23	12	.	*	89	263
American Samoa	35	.	.	.	.	.	.	.	35
Guam	107	*	.	25	*	.	*	15	151
Puerto Rico	2,112	.	10	388	196	11	25	384	3,126
Saipan	39	.	.	.	.	.	.	*	43
Virgin Islands	85	.	.	*	*	.	*	*	100
Foreign	51	*	*	*	.	.	*	30	88
Unknown	78	.	*	11	*	.	66	21	179
All	173,472	552	1,971	18,384	8,742	616	4,738	57,905	266,380

The point prevalent cohorts and associated modalities are determined on December 31 of each calendar year without applying the 60-day stable modality rule.

\* Values for cells with fewer than 10 patients are suppressed.

." Zero values in this cell.

**Table D.12 (continued)**

**Prevalent ESRD Patients for Selected Years: by Diabetic Status & State/Territory**

*point prevalent patients on December 31*

1996 DIABETIC	Center hemo	Center self hemo	Home hemo	CAPD	CCPD	Other PD	Uncertain dialysis	Transplant	All
Alabama	1,256	.	*	108	55	*	*	193	1,627
Alaska	54	.	*	13	*	.	.	23	93
Arizona	1,407	*	*	87	68	*	26	246	1,850
Arkansas	541	.	*	95	15	*	10	141	806
California	7,120	11	*	637	258	13	355	1,324	9,722
Colorado	612	.	*	59	18	*	*	240	938
Connecticut	625	.	*	101	77	*	30	146	983
Delaware	178	.	.	18	*	*	*	49	254
District of Columbia	338	*	*	*	*	*	*	14	368
Florida	3,269	*	138	230	170	*	69	671	4,562
Georgia	2,301	*	23	172	105	*	11	310	2,930
Hawaii	582	*	.	45	*	.	*	28	670
Idaho	129	.	*	40	*	.	*	76	263
Illinois	2,491	.	49	263	97	*	47	654	3,604
Indiana	1,084	.	*	217	46	*	25	254	1,634
Iowa	415	.	*	71	27	*	20	205	746
Kansas	430	*	*	97	26	*	*	111	674
Kentucky	664	.	*	64	41	*	24	171	970
Louisiana	1,539	.	*	64	49	*	*	210	1,881
Maine	158	.	.	11	*	.	*	65	243
Maryland	1,357	.	17	102	56	.	37	198	1,767
Massachusetts	1,060	*	11	63	61	*	15	282	1,498
Michigan	2,229	*	15	402	72	*	28	615	3,366
Minnesota	660	.	.	37	27	*	*	454	1,188
Mississippi	1,005	*	*	66	18	*	*	81	1,183
Missouri	1,163	.	25	160	81	*	*	299	1,735
Montana	145	.	*	18	*	.	*	83	254
Nebraska	231	.	*	109	12	.	*	144	502
Nevada	247	.	*	37	18	*	*	72	388
New Hampshire	108	.	.	24	10	.	*	45	191
New Jersey	1,935	22	*	196	118	24	105	313	2,721
New Mexico	616	.	.	75	41	12	10	113	867
New York	4,143	34	18	411	151	23	43	576	5,399
North Carolina	2,224	*	*	218	136	32	21	316	2,955
North Dakota	95	.	.	19	.	.	*	62	180
Ohio	2,333	.	19	394	148	19	48	661	3,622
Oklahoma	819	.	*	90	31	*	*	182	1,136
Oregon	408	.	*	83	24	*	13	177	715
Pennsylvania	2,911	*	13	200	103	13	84	651	3,981
Rhode Island	197	.	.	10	10	.	*	43	267
South Carolina	1,282	.	*	89	61	*	20	185	1,644
South Dakota	187	.	.	12	*	.	*	53	263
Tennessee	1,339	*	*	88	70	*	17	253	1,780
Texas	6,100	15	14	366	225	*	196	899	7,821
Utah	250	.	*	28	*	.	*	186	474
Vermont	67	.	*	19	*	.	*	42	134
Virginia	1,648	*	18	130	56	*	24	214	2,095
Washington	772	*	29	105	17	*	*	312	1,245
West Virginia	350	*	*	35	32	*	14	72	508
Wisconsin	781	.	*	101	53	*	29	535	1,509
Wyoming	60	.	*	12	*	.	.	30	108
American Samoa	17	.	.	.	.	.	.	.	17
Guam	59	*	.	13	.	.	.	*	74
Puerto Rico	918	.	*	155	88	*	*	29	1,200
Saipan	22	.	.	.	.	.	.	.	22
Virgin Islands	33	.	.	.	.	.	.	.	33
Foreign	*	*	.	*	.	.	.	*	13
Unknown	21	.	.	*	.	.	*	*	27
All	62,994	121	510	6,266	2,825	239	1,430	13,315	87,700

The point prevalent cohorts and associated modalities are determined on December 31 of each calendar year without applying the 60-day stable modality rule.

\* Values for cells with fewer than 10 patients are suppressed.

." Zero values in this cell.



**Table D.12 (continued)**

**Prevalent ESRD Patients for Selected Years: by Diabetic Status & State/Territory**

*point prevalent patients on December 31*

1996 NON-DIABETIC	Center hemo	Center self hemo	Home hemo	CAPD	CCPD	Other PD	Uncertain dialysis	Transplant	All
Alabama	2,553	*	11	254	148	*	24	979	3,979
Alaska	64	*	*	14	*	.	*	71	156
Arizona	1,432	16	18	120	117	*	39	607	2,350
Arkansas	1,129	.	10	115	38	*	19	497	1,811
California	10,843	35	31	1,178	643	33	567	5,185	18,515
Colorado	837	*	*	101	68	*	20	495	1,532
Connecticut	1,223	.	*	196	158	*	50	548	2,183
Delaware	393	*	.	58	19	.	15	135	621
District of Columbia	763	*	*	30	*	*	13	101	926
Florida	7,300	17	254	557	338	14	254	2,563	11,297
Georgia	4,429	.	64	338	234	*	61	1,202	6,335
Hawaii	539	.	*	48	*	.	*	177	784
Idaho	216	.	*	49	23	.	*	183	484
Illinois	5,239	*	72	546	227	*	161	2,201	8,457
Indiana	2,170	*	16	461	99	10	59	843	3,660
Iowa	754	.	25	112	47	*	33	517	1,489
Kansas	778	.	13	160	50	*	20	334	1,356
Kentucky	1,263	.	*	156	81	13	51	775	2,346
Louisiana	2,808	.	14	174	142	*	20	734	3,896
Maine	359	.	*	30	26	.	*	200	626
Maryland	2,702	21	31	308	135	10	126	754	4,087
Massachusetts	2,210	*	35	167	128	*	36	1,068	3,658
Michigan	3,680	.	44	719	133	*	54	1,771	6,408
Minnesota	1,322	.	*	83	35	*	33	794	2,275
Mississippi	1,955	*	62	136	42	*	32	468	2,701
Missouri	2,153	*	42	271	143	*	29	985	3,626
Montana	185	.	*	20	*	.	*	152	370
Nebraska	381	.	*	158	21	.	10	223	796
Nevada	491	.	*	55	18	*	23	270	863
New Hampshire	249	.	*	50	28	*	10	171	511
New Jersey	3,646	55	38	398	243	43	175	1,243	5,841
New Mexico	444	.	*	87	47	*	*	280	877
New York	8,259	177	85	955	328	30	116	2,573	12,523
North Carolina	3,963	*	35	483	291	43	77	1,119	6,013
North Dakota	142	.	.	17	16	.	*	125	304
Ohio	4,196	*	56	634	236	32	116	2,012	7,283
Oklahoma	1,033	*	*	127	49	*	19	524	1,768
Oregon	784	*	38	167	51	*	31	431	1,504
Pennsylvania	5,547	30	50	420	218	16	250	2,309	8,840
Rhode Island	439	.	.	41	18	.	10	183	691
South Carolina	2,406	*	19	208	136	*	54	760	3,589
South Dakota	200	.	*	27	14	.	*	99	347
Tennessee	2,572	14	38	209	121	*	55	1,039	4,050
Texas	7,765	18	66	581	430	*	245	3,080	12,192
Utah	296	.	*	32	12	*	11	308	665
Vermont	119	.	*	27	*	.	*	78	241
Virginia	3,392	*	48	316	180	*	139	873	4,958
Washington	1,349	12	115	162	64	*	28	707	2,446
West Virginia	562	*	*	80	54	*	35	270	1,014
Wisconsin	1,450	*	31	209	115	*	42	1,090	2,945
Wyoming	66	.	*	11	*	.	*	59	155
American Samoa	18	.	.	.	.	.	.	.	18
Guam	48	*	.	12	*	.	*	14	77
Puerto Rico	1,194	.	*	233	108	*	21	355	1,926
Saipan	17	.	.	.	.	.	.	*	21
Virgin Islands	52	.	.	*	*	.	*	*	67
Foreign	42	.	*	*	.	.	*	28	75
Unknown	57	.	*	10	*	.	65	17	152
All	110,478	431	1,461	12,118	5,917	377	3,308	44,590	178,680

The point prevalent cohorts and associated modalities are determined on December 31 of each calendar year without applying the 60-day stable modality rule.

\* Values for cells with fewer than 10 patients are suppressed.

." Zero values in this cell.

**Table D.12 (continued)**

**Prevalent ESRD Patients for Selected Years: by Diabetic Status & State/Territory**

*point prevalent patients on December 31*

2000 All	Center hemo	Center self hemo	Home hemo	CAPD	CCPD	Other PD	Uncertain		All
							dialysis	Transplant	
Alabama	4,827	.	22	220	244	*	80	1,502	6,903
Alaska	155	.	*	41	*	.	*	110	318
Arizona	3,706	10	23	150	142	*	66	1,127	5,227
Arkansas	2,019	.	*	178	75	*	28	800	3,107
California	22,590	20	39	1,173	966	18	557	8,173	33,536
Colorado	1,887	.	18	78	87	.	48	923	3,041
Connecticut	2,280	*	10	216	223	*	51	826	3,612
Delaware	679	.	.	13	36	.	28	256	1,012
District of Columbia	1,232	.	*	22	19	*	15	151	1,442
Florida	12,731	*	159	397	646	16	233	4,259	18,448
Georgia	8,659	*	127	346	330	15	199	1,915	11,592
Hawaii	1,388	*	*	76	31	.	26	290	1,817
Idaho	478	.	*	86	44	.	*	366	989
Illinois	9,415	*	63	488	408	*	214	3,443	14,043
Indiana	4,493	*	26	373	229	*	91	1,439	6,659
Iowa	1,557	*	22	116	87	.	36	878	2,697
Kansas	1,620	.	16	175	77	*	31	561	2,482
Kentucky	2,630	.	*	115	119	12	73	1,113	4,068
Louisiana	5,694	*	29	150	188	*	55	1,172	7,297
Maine	672	.	*	23	36	*	10	328	1,074
Maryland	4,799	*	16	266	196	*	98	1,720	7,104
Massachusetts	3,564	.	19	146	155	13	92	1,629	5,618
Michigan	7,634	.	27	589	270	*	148	2,919	11,588
Minnesota	2,334	.	*	65	72	*	41	1,582	4,098
Mississippi	3,649	*	52	142	166	*	41	732	4,787
Missouri	4,284	*	43	232	234	*	69	1,540	6,405
Montana	428	.	*	33	20	*	*	293	787
Nebraska	924	*	*	185	50	.	21	472	1,657
Nevada	1,009	.	*	77	51	*	37	450	1,632
New Hampshire	478	.	*	47	24	.	22	259	834
New Jersey	7,288	17	33	227	335	14	168	2,189	10,271
New Mexico	1,537	.	*	92	114	*	32	482	2,264
New York	15,588	199	72	631	542	45	257	4,336	21,670
North Carolina	8,267	.	39	378	442	126	130	1,954	11,336
North Dakota	286	.	.	34	*	.	12	234	574
Ohio	8,846	.	22	637	463	*	167	3,282	13,423
Oklahoma	2,434	.	11	169	153	23	44	840	3,674
Oregon	1,500	.	19	126	104	*	28	785	2,563
Pennsylvania	9,978	*	33	315	325	*	248	3,644	14,553
Rhode Island	702	.	.	28	15	.	22	296	1,063
South Carolina	4,678	*	14	161	305	*	87	1,226	6,476
South Dakota	512	.	*	25	32	.	12	198	780
Tennessee	4,958	.	51	187	247	10	108	1,657	7,218
Texas	18,857	.	73	569	819	*	292	5,184	25,803
Utah	707	*	*	63	14	.	25	650	1,464
Vermont	306	.	*	*	16	.	12	150	495
Virginia	6,313	*	47	305	286	11	126	1,704	8,794
Washington	2,663	*	84	153	119	*	51	1,398	4,484
West Virginia	1,267	*	*	87	112	*	27	561	2,060
Wisconsin	2,963	*	15	192	144	*	74	1,958	5,351
Wyoming	201	.	*	13	16	.	*	123	360
American Samoa	52	.	.	.	.	.	*	*	54
Guam	198	.	.	10	*	.	*	13	225
Puerto Rico	2,501	.	15	137	324	*	37	567	3,586
Saipan	77	.	.	.	.	.	.	*	86
Virgin Islands	112	.	.	*	*	.	*	11	130
Foreign	48	.	.	*	.	.	*	25	81
Unknown	51	.	.	*	*	.	*	20	79
All	220,705	282	1,310	10,773	10,173	421	4,402	74,725	322,791

The point prevalent cohorts and associated modalities are determined on December 31 of each calendar year without applying the 60-day stable modality rule.

\* Values for cells with fewer than 10 patients are suppressed.

." Zero values in this cell.

**Table D.12 (continued)**

**Prevalent ESRD Patients for Selected Years: by Diabetic Status & State/Territory**

*point prevalent patients on December 31*

<b>2000 DIABETIC</b>	<b>Center hemo</b>	<b>Center self hemo</b>	<b>Home hemo</b>	<b>CAPD</b>	<b>CCPD</b>	<b>Other PD</b>	<b>Uncertain</b>		<b>All</b>
							<b>dialysis</b>	<b>Transplant</b>	
Alabama	1,947	.	*	54	77	*	17	279	2,384
Alaska	71	.	*	17	.	.	.	27	116
Arizona	1,972	.	11	55	56	*	17	326	2,439
Arkansas	740	.	*	68	20	.	*	189	1,024
California	10,003	*	*	446	286	*	124	1,756	12,633
Colorado	879	.	*	31	23	.	*	299	1,243
Connecticut	868	.	*	93	80	*	11	158	1,216
Delaware	278	.	.	*	10	.	11	69	369
District of Columbia	384	.	.	*	*	.	*	33	427
Florida	4,641	*	50	124	192	*	39	889	5,945
Georgia	3,364	.	41	101	95	*	42	416	4,067
Hawaii	766	.	*	38	16	.	10	54	885
Idaho	208	.	*	41	23	.	*	104	380
Illinois	3,485	*	25	172	134	*	47	844	4,714
Indiana	1,752	.	12	138	83	*	11	321	2,322
Iowa	603	.	*	53	29	.	*	262	959
Kansas	700	.	*	66	27	*	*	138	942
Kentucky	1,066	.	*	46	35	*	12	220	1,387
Louisiana	2,234	.	*	49	54	*	*	275	2,625
Maine	252	.	.	*	15	*	*	77	350
Maryland	1,722	.	*	70	63	*	18	404	2,283
Massachusetts	1,266	.	*	50	50	*	27	336	1,736
Michigan	3,192	.	*	222	86	.	31	780	4,318
Minnesota	878	.	.	19	17	.	10	531	1,455
Mississippi	1,371	*	*	50	57	.	*	116	1,610
Missouri	1,767	.	10	80	83	.	21	367	2,328
Montana	191	.	.	17	13	.	*	110	333
Nebraska	388	*	*	69	20	.	*	180	667
Nevada	416	.	*	35	23	*	*	103	585
New Hampshire	159	.	*	14	*	.	*	64	250
New Jersey	2,837	*	*	67	118	*	31	469	3,532
New Mexico	952	.	*	48	59	*	11	158	1,230
New York	5,853	49	11	212	163	21	46	869	7,224
North Carolina	3,383	.	*	101	111	47	25	470	4,142
North Dakota	123	.	.	12	*	.	*	75	217
Ohio	3,741	.	*	283	181	*	39	854	5,104
Oklahoma	1,151	.	*	90	66	12	*	228	1,558
Oregon	565	.	*	50	41	*	*	216	886
Pennsylvania	3,843	.	*	108	105	*	69	826	4,962
Rhode Island	252	.	.	*	*	.	*	62	333
South Carolina	1,905	.	*	41	100	*	15	241	2,307
South Dakota	251	.	.	11	*	.	*	68	342
Tennessee	1,895	.	*	56	84	*	27	333	2,406
Texas	9,237	.	23	226	322	*	55	1,243	11,108
Utah	330	*	*	34	*	.	*	229	608
Vermont	125	.	*	*	*	.	*	38	178
Virginia	2,462	.	10	117	82	*	20	349	3,044
Washington	1,086	*	22	54	40	*	13	420	1,639
West Virginia	614	*	.	35	54	*	*	128	839
Wisconsin	1,104	.	*	60	44	*	18	619	1,848
Wyoming	97	.	*	*	*	.	*	40	150
American Samoa	32	.	.	.	.	.	.	.	32
Guam	111	.	.	*	*	.	.	.	116
Puerto Rico	1,245	.	*	61	136	*	*	59	1,511
Saipan	53	.	.	.	.	.	.	.	53
Virgin Islands	46	.	.	*	*	.	.	*	50
Foreign	14	.	.	*	.	.	.	*	16
Unknown	*	.	.	.	.	.	.	*	*
All	90,871	67	331	3,914	3,422	171	929	17,726	117,431

The point prevalent cohorts and associated modalities are determined on December 31 of each calendar year without applying the 60-day stable modality rule.

\* Values for cells with fewer than 10 patients are suppressed.

." Zero values in this cell.

**Table D.12 (continued)**  
**Prevalent ESRD Patients for Selected Years: by Diabetic Status & State/Territory**  
*point prevalent patients on December 31*

2000 NON-DIABETIC	Center hemo	Center self hemo	Home hemo	CAPD	CCPD	Other PD	Uncertain dialysis	Transplant	All
Alabama	2,880	.	16	166	167	*	63	1,223	4,519
Alaska	84	.	*	24	*	.	*	83	202
Arizona	1,734	10	12	95	86	*	49	801	2,788
Arkansas	1,279	.	*	110	55	*	22	611	2,083
California	12,587	12	32	727	680	15	433	6,417	20,903
Colorado	1,008	.	11	47	64	.	44	624	1,798
Connecticut	1,412	*	*	123	143	*	40	668	2,396
Delaware	401	.	.	12	26	.	17	187	643
District of Columbia	848	.	*	16	17	*	13	118	1,015
Florida	8,090	*	109	273	454	*	194	3,370	12,503
Georgia	5,295	*	86	245	235	*	157	1,499	7,525
Hawaii	622	*	*	38	15	.	16	236	932
Idaho	270	.	*	45	21	.	*	262	609
Illinois	5,930	*	38	316	274	*	167	2,599	9,329
Indiana	2,741	*	14	235	146	*	80	1,118	4,337
Iowa	954	*	17	63	58	.	29	616	1,738
Kansas	920	.	14	109	50	*	23	423	1,540
Kentucky	1,564	.	*	69	84	*	61	893	2,681
Louisiana	3,460	*	21	101	134	*	52	897	4,672
Maine	420	.	*	19	21	.	*	251	724
Maryland	3,077	*	12	196	133	*	80	1,316	4,821
Massachusetts	2,298	.	17	96	105	*	65	1,293	3,882
Michigan	4,442	.	20	367	184	*	117	2,139	7,270
Minnesota	1,456	.	*	46	55	*	31	1,051	2,643
Mississippi	2,278	*	44	92	109	*	34	616	3,177
Missouri	2,517	*	33	152	151	*	48	1,173	4,077
Montana	237	.	*	16	*	*	*	183	454
Nebraska	536	.	*	116	30	.	14	292	990
Nevada	593	.	*	42	28	*	32	347	1,047
New Hampshire	319	.	*	33	15	.	19	195	584
New Jersey	4,451	15	27	160	217	12	137	1,720	6,739
New Mexico	585	.	*	44	55	*	21	324	1,034
New York	9,735	150	61	419	379	24	211	3,467	14,446
North Carolina	4,884	.	34	277	331	79	105	1,484	7,194
North Dakota	163	.	.	22	*	.	*	159	357
Ohio	5,105	.	18	354	282	*	128	2,428	8,319
Oklahoma	1,283	.	*	79	87	11	36	612	2,116
Oregon	935	.	15	76	63	.	19	569	1,677
Pennsylvania	6,135	*	26	207	220	*	179	2,818	9,591
Rhode Island	450	.	.	20	11	.	15	234	730
South Carolina	2,773	*	12	120	205	*	72	985	4,169
South Dakota	261	.	*	14	24	.	*	130	438
Tennessee	3,063	.	44	131	163	*	81	1,324	4,812
Texas	9,620	.	50	343	497	*	237	3,941	14,695
Utah	377	.	*	29	*	.	19	421	856
Vermont	181	.	*	*	*	.	*	112	317
Virginia	3,851	*	37	188	204	*	106	1,355	5,750
Washington	1,577	*	62	99	79	*	38	978	2,845
West Virginia	653	.	*	52	58	*	22	433	1,221
Wisconsin	1,859	*	13	132	100	*	56	1,339	3,503
Wyoming	104	.	*	10	*	.	*	83	210
American Samoa	20	.	.	.	.	.	*	*	22
Guam	87	.	.	*	*	.	*	13	109
Puerto Rico	1,256	.	11	76	188	*	34	508	2,075
Saipan	24	.	.	.	.	.	.	*	33
Virgin Islands	66	.	.	*	*	.	*	10	80
Foreign	34	.	.	*	.	.	*	24	65
Unknown	50	.	.	*	*	.	*	17	75
All	129,834	215	979	6,859	6,751	250	3,473	56,999	205,360

The point prevalent cohorts and associated modalities are determined on December 31 of each calendar year without applying the 60-day stable modality rule.

\* Values for cells with fewer than 10 patients are suppressed.

." Zero values in this cell.