

Reference Section D
treatment modalities



table d.1 · 341

counts of reported esrd patients: by treatment modality
incident & point prevalent patients

table d.2 · 342

incident esrd patients: by dialysis/functioning graft status
by demographic characteristics & primary diagnosis

table d.3 · 346

prevalent esrd patients: by dialysis/functioning graft status
point prevalent patients on december 31, by demographic characteristics & primary diagnosis

table d.4 · 350

incident esrd patients for selected years: by age & modality

table d.5 · 352

prevalent esrd patients for selected years: by age & modality
point prevalent patients on december 31

table d.6 · 354

incident esrd patients for selected years: by gender, race, ethnicity, & primary diagnosis; grouped by modality
incident patients

table d.7 · 356

prevalent esrd patients for selected years: by gender, race, ethnicity, & primary diagnosis; grouped by modality
point prevalent patients on december 31

table d.8 · 358

treatment status two years after first esrd service
incident patients, 1996-1998 combined, by gender, age at first service, & modality at 90 days

table d.9 · 361

prevalent esrd patients prevalent for selected years: by patient vintage
point prevalent patients on december 31, by year-end treatment modality & number of years patient has had esrd (vintage)

table d.10 · 363

prevalent esrd patients for selected years: by number of years on modality
point prevalent patients on december 31, by year-end treatment modality

table d.11 · 365

incident esrd patients for selected years: by diabetic status & state/territory

table d.12 · 374

prevalent esrd patients for selected years: by diabetic status & state/territory
point prevalent patients on december 31

table d.13 · 383

prevalent dialysis patients: by age & state/territory
point prevalent dialysis patients on december 31, 1998-2000 combined, by state & age; grouped by modality

Table D.1

Counts of Reported ESRD Patients: by Treatment Modality*incident & point prevalent patients*

INCIDENT COUNTS	1991	1992	1993	1994	1995	1996	1997	1998	1999	2000
Modality										
Dialysis	52,345	57,387	59,701	71,733	71,262	75,975	80,313	85,346	89,525	94,022
Functioning graft	1,777	1,740	1,926	1,747	1,612	1,556	1,598	1,716	1,899	2,170
Dialysis type										
Center hemodialysis	41,413	45,623	46,348	53,689	57,223	61,530	68,056	74,726	79,625	83,294
Center self hemodialysis	66	63	49	39	68	64	54	47	46	29
Home hemodialysis	31	31	29	41	349	1,010	174	182	239	312
CAPD	5,528	5,722	5,820	5,806	6,302	5,932	5,485	5,077	4,921	4,724
CCPD	348	393	472	763	1,858	2,103	2,067	2,187	2,198	2,377
Other PD	680	725	677	877	656	512	492	434	328	271
Uncertain dialysis	4,279	4,830	6,306	10,518	4,806	4,824	3,985	2,693	2,168	3,015
All	54,122	59,127	61,627	73,480	72,874	77,531	81,911	87,062	91,424	96,192
PERCENTAGES										
Modality										
Dialysis	96.7	97.1	96.9	97.6	97.8	98.0	98.0	98.0	97.9	97.7
Functioning graft	3.3	2.9	3.1	2.4	2.2	2.0	2.0	2.0	2.1	2.3
Dialysis type										
Center hemodialysis	79.1	79.5	77.6	74.8	80.3	81.0	84.7	87.6	88.9	88.6
Center self hemodialysis	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.1	.
Home hemo	0.1	0.1	.	0.1	0.5	1.3	0.2	0.2	0.3	0.3
CAPD	10.6	10.0	9.7	8.1	8.8	7.8	6.8	5.9	5.5	5.0
CCPD	0.7	0.7	0.8	1.1	2.6	2.8	2.6	2.6	2.5	2.5
Other PD	1.3	1.3	1.1	1.2	0.9	0.7	0.6	0.5	0.4	0.3
Uncertain dialysis	8.2	8.4	10.6	14.7	6.7	6.3	5.0	3.2	2.4	3.2
All	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0
POINT PREVALENT COUNTS										
Modality										
Dialysis	144,558	159,850	173,401	195,544	212,425	226,949	238,381	251,505	264,042	275,053
Functioning graft	55,393	60,584	66,232	71,344	75,711	81,125	86,961	93,098	97,879	103,809
Dialysis type										
Center hemodialysis	116,652	128,720	140,379	154,796	168,903	185,612	202,326	218,520	232,120	244,410
Center self hemodialysis	747	814	153	225	533	552	486	364	337	282
Home hemodialysis	1,613	1,645	351	611	1,829	2,124	1,676	1,327	1,235	1,429
CAPD	17,969	19,833	21,220	22,073	21,371	19,414	17,324	15,322	13,473	12,307
CCPD	2,239	2,909	3,666	4,668	7,192	9,093	9,844	9,954	10,745	10,980
Other PD	685	678	636	858	825	724	600	542	418	458
Uncertain dialysis	4,653	5,251	6,996	12,313	11,772	9,430	6,125	5,476	5,714	5,187
All	199,951	220,434	239,633	266,888	288,136	308,074	325,342	344,603	361,921	378,862
PERCENTAGES										
Modality										
Dialysis	72.3	72.5	72.4	73.3	73.7	73.7	73.3	73.0	73.0	72.6
Functioning graft	27.7	27.5	27.6	26.7	26.3	26.3	26.7	27.0	27.0	27.4
Dialysis type										
Center hemodialysis	80.7	80.5	81.0	79.2	79.5	81.8	84.9	86.9	87.9	88.9
Center self hemodialysis	0.5	0.5	0.1	0.1	0.3	0.2	0.2	0.1	0.1	0.1
Home hemodialysis	1.1	1.0	0.2	0.3	0.9	0.9	0.7	0.5	0.5	0.5
CAPD	12.4	12.4	12.2	11.3	10.1	8.6	7.3	6.1	5.1	4.5
CCPD	1.5	1.8	2.1	2.4	3.4	4.0	4.1	4.0	4.1	4.0
Other PD	0.5	0.4	0.4	0.4	0.4	0.3	0.3	0.2	0.2	0.2
Uncertain dialysis	3.2	3.3	4.0	6.3	5.5	4.2	2.6	2.2	2.2	1.9
All	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0

The incident cohorts and associated modalities are determined at the time of ESRD initiation without applying the 60-day stable modality rule.

The point prevalent cohorts and associated modalities are determined on December 31 of each calendar year without applying the 60-day stable modality rule.

." Zero values in this cell.

Table D.2
Incident ESRD Patients: by Dialysis/Functioning Graft Status
by demographic characteristics & primary diagnosis

ALL DIALYSIS	1991	1992	1993	1994	1995	1996	1997	1998	1999	2000
0-4	123	102	117	129	155	155	156	164	164	220
5-9	78	63	88	124	110	111	97	105	127	118
10-14	170	169	174	210	220	230	236	218	236	239
15-19	380	428	404	477	489	511	475	485	523	518
20-24	813	910	868	1,030	1,009	981	905	965	907	1,000
25-29	1,404	1,425	1,498	1,807	1,574	1,614	1,615	1,655	1,580	1,723
30-34	1,934	2,034	2,086	2,586	2,416	2,358	2,309	2,353	2,367	2,459
35-39	2,411	2,674	2,628	3,463	3,201	3,224	3,205	3,216	3,328	3,324
40-44	2,974	3,188	3,209	3,929	4,038	4,175	4,115	4,442	4,426	4,694
45-49	3,195	3,522	3,793	4,817	4,901	5,223	5,281	5,714	5,938	6,267
50-54	3,573	4,101	4,265	5,657	5,529	5,697	6,262	6,838	7,453	7,913
55-59	4,622	5,056	5,211	6,641	6,566	6,810	7,275	7,945	8,480	9,008
60-64	6,284	6,603	6,816	8,143	8,274	8,456	8,874	9,454	9,642	10,135
65-69	7,929	8,546	8,849	9,914	9,614	10,240	10,703	10,948	11,073	11,359
70-74	7,245	8,081	8,509	9,735	9,647	10,623	11,199	11,575	11,892	12,250
75-79	5,354	5,982	6,282	7,295	7,393	8,401	9,282	10,064	11,079	11,495
80-84	2,743	3,218	3,418	4,076	4,281	4,931	5,716	6,083	6,738	7,382
85+	1,104	1,271	1,479	1,671	1,838	2,232	2,602	3,114	3,560	3,887
Missing/unknown	*	14	*	29	*	*	*	*	12	31
0-19	751	762	783	940	974	1,007	964	972	1,050	1,095
20-44	9,536	10,231	10,289	12,815	12,238	12,352	12,149	12,631	12,608	13,200
45-64	17,674	19,282	20,085	25,258	25,270	26,186	27,692	29,951	31,513	33,323
65-74	15,174	16,627	17,358	19,649	19,261	20,863	21,902	22,523	22,965	23,609
75+	9,201	10,471	11,179	13,042	13,512	15,564	17,600	19,261	21,377	22,764
Unknown	*	14	*	29	*	*	*	*	12	31
Male	28,020	30,400	31,781	37,523	37,511	40,483	42,606	45,327	47,795	50,341
Female	24,322	26,985	27,917	34,147	33,733	35,470	37,702	40,013	41,723	43,673
Unknown	*	*	*	63	18	22	*	*	*	*
White	33,895	36,800	37,347	43,600	43,539	46,157	50,573	53,275	56,919	59,760
Black	16,017	17,588	18,739	21,786	22,359	23,820	24,281	25,995	26,566	27,332
Native American	706	863	820	994	1,172	1,337	977	1,409	1,133	1,210
Asian	1,325	1,511	1,774	2,279	2,486	2,710	2,655	3,036	3,185	3,206
Other/unknown	402	625	1,021	3,074	1,706	1,951	1,827	1,631	1,722	2,514
†Hispanic					7,119	8,877	9,008	9,938	10,283	11,154
†Non-Hispanic					64,143	67,098	71,305	75,408	79,242	82,868
Diabetes	18,964	21,136	21,114	25,845	28,791	32,345	35,056	37,551	39,398	41,494
Hypertension	15,754	17,628	16,907	20,362	18,331	19,663	21,278	22,576	23,786	24,457
Glomerulonephritis	6,126	6,377	6,290	7,268	7,467	7,982	7,964	8,121	8,037	7,793
Cystic kidney	1,430	1,527	1,487	1,713	1,762	1,731	1,833	1,946	1,992	1,939
Other urologic	1,095	1,146	1,012	1,145	1,302	1,509	1,447	1,504	1,628	1,567
Other cause	4,894	5,207	5,144	6,322	7,828	8,347	8,703	9,324	9,770	10,192
Unknown cause	2,260	2,513	2,226	2,534	2,550	2,939	3,099	3,391	3,460	3,756
Missing cause	1,822	1,853	5,521	6,544	3,231	1,459	933	933	1,454	2,824
All	52,345	57,387	59,701	71,733	71,262	75,975	80,313	85,346	89,525	94,022

The incident cohorts and associated modalities are determined at the time of ESRD initiation without applying the 60-day stable modality rule.

The age of the point prevalent patient is determined as of December 31 of each calendar year.

†CMS has only started collecting Hispanic ethnicity data as of April, 1995, so Hispanic data in years prior to 1995 has been left blank.

* Values for cells with fewer than 10 patients are suppressed.

Table D.2 (continued)

Incident ESRD Patients: by Dialysis/Functioning Graft Status

by demographic characteristics & primary diagnosis

HEMODIALYSIS	1991	1992	1993	1994	1995	1996	1997	1998	1999	2000
0-4	24	17	26	28	38	30	46	37	45	80
5-9	26	21	32	46	49	40	36	42	52	55
10-14	73	85	71	94	107	98	117	112	128	134
15-19	283	293	268	302	339	361	343	374	414	390
20-24	641	700	639	706	824	796	765	843	795	827
25-29	1,094	1,107	1,070	1,246	1,229	1,283	1,344	1,393	1,355	1,411
30-34	1,461	1,517	1,497	1,720	1,861	1,871	1,927	1,966	2,020	2,034
35-39	1,810	2,039	1,943	2,336	2,438	2,590	2,690	2,728	2,863	2,769
40-44	2,220	2,433	2,339	2,642	3,175	3,334	3,419	3,856	3,830	3,962
45-49	2,483	2,736	2,822	3,408	3,784	4,215	4,491	4,932	5,168	5,323
50-54	2,811	3,219	3,276	4,022	4,384	4,632	5,264	5,889	6,554	6,803
55-59	3,761	4,046	4,045	4,907	5,325	5,630	6,204	6,980	7,504	7,901
60-64	5,117	5,352	5,386	6,121	6,783	7,116	7,758	8,397	8,615	9,035
65-69	6,309	6,936	6,995	7,791	7,824	8,474	9,089	9,654	9,991	10,326
70-74	5,862	6,553	6,904	7,760	8,035	8,855	9,586	10,310	10,780	11,261
75-79	4,358	4,911	5,142	5,914	6,229	7,139	7,957	9,054	10,189	10,662
80-84	2,261	2,694	2,759	3,357	3,644	4,231	5,000	5,518	6,249	6,935
85+	913	1,051	1,207	1,345	1,567	1,907	2,244	2,862	3,347	3,697
Missing/unknown	*	*	*	24	*	*	*	*	11	30
0-19	406	416	397	470	533	529	542	565	639	659
20-44	7,226	7,796	7,488	8,650	9,527	9,874	10,145	10,786	10,863	11,003
45-64	14,172	15,353	15,529	18,458	20,276	21,593	23,717	26,198	27,841	29,062
65-74	12,171	13,489	13,899	15,551	15,859	17,329	18,675	19,964	20,771	21,587
75+	7,532	8,656	9,108	10,616	11,440	13,277	15,201	17,434	19,785	21,294
Unknown	*	*	*	24	*	*	*	*	11	30
Male	22,210	24,293	24,924	28,488	30,387	33,487	36,392	39,919	42,715	44,680
Female	19,300	21,424	21,502	25,280	27,252	29,114	31,890	35,030	37,190	38,947
Unknown	.	.	.	*	*	*	*	*	*	*
White	26,254	28,738	28,675	33,189	34,452	37,167	41,908	45,899	49,986	52,670
Black	13,481	14,925	15,591	17,580	18,991	20,555	21,735	23,745	24,528	24,914
Native American	581	720	655	789	971	1,138	858	1,254	1,040	1,065
Asian	1,105	1,237	1,344	1,739	1,986	2,226	2,243	2,642	2,822	2,831
Other/unknown	89	97	161	472	1,240	1,518	1,540	1,415	1,534	2,155
†Hispanic					5,846	7,460	7,675	8,770	9,373	10,278
†Non-Hispanic					51,794	55,144	60,609	66,185	70,537	73,357
Diabetes	15,021	16,774	16,568	20,543	23,508	26,675	29,910	33,128	35,363	37,714
Hypertension	12,927	14,648	13,829	16,741	15,503	16,887	18,619	20,383	21,869	22,658
Glomerulonephritis	4,758	4,925	4,751	5,493	5,838	6,254	6,512	6,786	6,876	6,651
Cystic kidney	1,054	1,155	1,073	1,244	1,307	1,298	1,427	1,574	1,595	1,583
Other urologic	923	923	817	910	1,090	1,273	1,234	1,332	1,471	1,413
Other cause	3,810	4,163	4,047	5,025	6,480	7,014	7,436	8,251	8,778	9,300
Unknown cause	1,809	2,068	1,752	2,035	2,151	2,502	2,653	3,060	3,129	3,462
Missing cause	1,208	1,061	3,589	1,778	1,763	701	493	441	829	854
All	41,510	45,717	46,426	53,769	57,640	62,604	68,284	74,955	79,910	83,635

The incident cohorts and associated modalities are determined at the time of ESRD initiation without applying the 60-day stable modality rule.

The age of the incident patient is determined as of the date of ESRD initiation.

†CMS has only started collecting Hispanic ethnicity data as of April, 1995, so Hispanic data in years prior to 1995 has been left blank.

* Values for cells with fewer than 10 patients are suppressed.

." Zero values in this cell.

Table D.2 (continued)
Incident ESRD Patients: by Dialysis/Functioning Graft Status
by demographic characteristics & primary diagnosis

PERITONEAL DIALYSIS	1991	1992	1993	1994	1995	1996	1997	1998	1999	2000
0-4	55	40	42	46	93	108	104	120	112	135
5-9	37	15	29	40	53	65	56	57	67	63
10-14	64	37	53	53	86	109	103	95	104	98
15-19	67	81	71	73	130	123	116	102	98	102
20-24	111	128	121	129	140	137	131	105	92	100
25-29	228	199	233	226	267	268	238	233	199	198
30-34	357	371	345	330	398	395	334	339	311	285
35-39	434	426	417	435	559	508	441	416	403	376
40-44	558	505	507	470	645	657	590	503	513	507
45-49	498	523	566	583	843	773	664	679	647	624
50-54	522	560	582	649	832	783	823	798	732	741
55-59	546	634	613	659	861	832	845	807	800	751
60-64	724	771	692	780	982	907	845	783	811	803
65-69	655	656	773	729	868	806	752	761	715	756
70-74	517	566	606	678	666	782	694	672	681	672
75-79	330	383	388	406	443	468	487	466	489	534
80-84	133	159	186	210	221	222	228	231	237	248
85+	39	59	67	71	73	92	100	97	108	107
Missing/unknown	*	*	*	*	.	.	*	.	.	*
0-19	223	173	195	212	362	405	379	374	381	398
20-44	1,688	1,629	1,623	1,590	2,009	1,965	1,734	1,596	1,518	1,466
45-64	2,290	2,488	2,453	2,671	3,518	3,295	3,177	3,067	2,990	2,919
65-74	1,172	1,222	1,379	1,407	1,534	1,588	1,446	1,433	1,396	1,428
75+	502	601	641	687	737	782	815	794	834	889
Unknown	*	*	*	*	.	.	*	.	.	*
Male	3,239	3,276	3,301	3,485	4,328	4,262	3,952	3,784	3,717	3,803
Female	2,636	2,838	2,991	3,084	3,832	3,773	3,599	3,480	3,402	3,298
Unknown	*	*	*	.	.	.
White	4,257	4,440	4,488	4,630	5,621	5,556	5,433	5,176	5,208	5,144
Black	1,417	1,448	1,545	1,571	1,938	1,786	1,594	1,541	1,476	1,418
Native American	76	73	68	88	130	135	81	111	67	94
Asian	120	147	180	244	325	344	270	281	241	262
Other/unknown	*	*	11	36	146	214	174	155	127	183
†Hispanic					715	848	742	755	748	773
†Non-Hispanic					7,445	7,187	6,810	6,509	6,371	6,328
Diabetes	2,289	2,449	2,349	2,593	3,430	3,584	3,320	3,195	3,144	3,115
Hypertension	1,322	1,401	1,400	1,487	1,557	1,416	1,432	1,404	1,362	1,402
Glomerulonephritis	923	950	907	984	1,287	1,314	1,186	1,136	1,020	1,036
Cystic kidney	281	271	293	301	376	358	355	331	363	330
Other urologic	71	84	91	90	128	135	128	119	121	127
Other cause	633	583	590	613	872	885	824	815	805	768
Unknown cause	218	220	230	207	243	281	268	230	233	236
Missing cause	139	157	432	294	267	62	39	34	71	87
All	5,876	6,115	6,292	6,569	8,160	8,035	7,552	7,264	7,119	7,101

The incident cohorts and associated modalities are determined at the time of ESRD initiation without applying the 60-day stable modality rule.

The age of the incident patient is determined as of the date of ESRD initiation.

†CMS has only started collecting Hispanic ethnicity data as of April, 1995, so Hispanic data in years prior to 1995 has been left blank.

* Values for cells with fewer than 10 patients are suppressed.

." Zero values in this cell.

Table D.2 (continued)

Incident ESRD Patients: by Dialysis/Functioning Graft Status

by demographic characteristics & primary diagnosis

TRANSPLANT	1991	1992	1993	1994	1995	1996	1997	1998	1999	2000
0-4	44	46	55	36	29	49	46	29	45	34
5-9	65	59	56	48	47	43	45	37	49	45
10-14	80	76	74	62	75	59	58	67	74	82
15-19	94	75	121	105	96	93	83	64	92	102
20-24	122	109	116	94	71	78	73	73	75	75
25-29	181	173	168	151	129	99	109	107	129	121
30-34	242	212	251	210	180	146	136	170	141	188
35-39	212	217	227	219	219	189	187	210	199	210
40-44	256	237	219	230	183	211	193	225	244	273
45-49	179	184	206	219	194	196	188	212	245	279
50-54	106	132	177	134	154	156	161	208	228	263
55-59	112	107	113	127	106	105	155	150	171	216
60-64	55	71	90	76	78	87	101	94	112	142
65-69	23	30	37	28	37	36	52	56	62	93
70-74	*	*	12	*	12	*	*	*	31	42
75-79	*	*	*	*	*	*	*	*	*	*
80-84	.	*	*	*	*
85+	.	*	*	*	.	.
Missing/unknown	.	.	*	.	.	*	.	.	.	*
0-19	283	256	306	251	247	244	232	197	260	263
20-44	1,013	948	981	904	782	723	698	785	788	867
45-64	452	494	586	556	532	544	605	664	756	900
65-74	28	38	49	32	49	42	59	63	93	135
75+	*	*	*	*	*	*	*	*	*	*
Unknown	.	.	*	.	.	*	.	.	.	*
Male	1,035	961	1,085	993	906	932	950	972	1,091	1,232
Female	739	775	834	754	706	624	648	744	808	938
Unknown	*	*	*
White	1,387	1,315	1,468	1,139	1,096	1,124	1,305	1,428	1,589	1,788
Black	228	252	273	453	400	337	192	178	198	208
Native American	26	14	16	16	16	11	*	14	11	12
Asian	67	81	80	88	54	50	46	53	44	78
Other/unknown	69	78	89	51	46	34	47	43	57	84
†Hispanic					63	86	72	91	84	112
†Non-Hispanic					1,549	1,470	1,526	1,625	1,815	2,058
Diabetes	258	254	265	278	271	316	302	280	320	278
Hypertension	70	94	78	98	66	68	81	83	85	109
Glomerulonephritis	184	215	214	214	253	253	255	270	279	309
Cystic kidney	104	103	97	86	130	146	133	192	170	205
Other urologic	41	30	42	36	41	51	52	53	40	63
Other cause	167	175	197	181	236	228	259	220	234	252
Unknown cause	48	54	66	49	52	57	60	42	26	85
Missing cause	905	815	967	805	563	437	456	576	745	869
All	1,777	1,740	1,926	1,747	1,612	1,556	1,598	1,716	1,899	2,170

The incident cohorts and associated modalities are determined at the time of ESRD initiation without applying the 60-day stable modality rule.

The age of the incident patient is determined as of the date of ESRD initiation.

†CMS has only started collecting Hispanic ethnicity data as of April, 1995, so Hispanic data in years prior to 1995 has been left blank.

* Values for cells with fewer than 10 patients are suppressed.

." Zero values in this cell.

Table D.3

Prevalent ESRD Patients: by Dialysis/Functioning Graft Status

point prevalent patients on December 31, by demographic characteristics & primary diagnosis

ALL DIALYSIS	1991	1992	1993	1994	1995	1996	1997	1998	1999	2000
0-4	162	170	187	235	275	277	253	266	289	364
5-9	160	162	170	216	225	225	225	213	231	240
10-14	369	375	385	433	464	461	453	478	458	481
15-19	958	1,001	1,038	1,055	1,132	1,187	1,194	1,218	1,212	1,165
20-24	2,295	2,493	2,571	2,834	2,988	2,968	2,880	2,980	3,026	3,109
25-29	4,074	4,294	4,399	4,887	5,194	5,383	5,411	5,601	5,667	5,613
30-34	6,216	6,560	6,972	7,772	8,198	8,225	8,101	8,072	8,301	8,516
35-39	7,943	8,698	9,199	10,477	11,218	11,511	11,553	11,838	12,267	12,367
40-44	9,690	10,530	11,289	12,576	13,954	14,757	15,218	15,843	16,477	16,785
45-49	10,364	11,502	12,738	14,833	16,795	18,203	18,767	19,820	20,851	21,734
50-54	11,343	12,786	14,023	16,383	17,861	19,416	20,677	22,418	24,567	26,480
55-59	13,745	14,995	16,225	18,375	20,128	21,580	23,053	24,789	26,350	27,823
60-64	17,176	18,393	19,561	21,750	23,725	25,140	26,353	27,904	29,160	30,228
65-69	20,279	22,391	24,134	26,237	27,594	29,086	30,031	30,963	31,977	32,694
70-74	18,128	20,509	22,573	25,382	26,994	28,692	30,386	31,511	32,189	32,847
75-79	12,565	14,207	15,573	17,852	19,755	22,021	24,097	25,919	27,553	28,782
80-84	6,480	7,680	8,710	10,011	11,094	12,244	13,469	14,552	15,508	17,221
85+	2,591	3,081	3,636	4,207	4,801	5,542	6,246	7,103	7,938	8,575
Missing/unknown	20	23	18	29	30	31	14	17	21	29
0-19	1,649	1,708	1,780	1,939	2,096	2,150	2,125	2,175	2,190	2,250
20-44	30,218	32,575	34,430	38,546	41,552	42,844	43,163	44,334	45,738	46,390
45-64	52,628	57,676	62,547	71,341	78,509	84,339	88,850	94,931	100,928	106,265
65-74	38,407	42,900	46,707	51,619	54,588	57,778	60,417	62,474	64,166	65,541
75+	21,636	24,968	27,919	32,070	35,650	39,807	43,812	47,574	50,999	54,578
Unknown	20	23	18	29	30	31	14	17	21	29
Male	75,190	82,950	90,065	101,128	109,869	117,881	124,401	131,932	139,441	145,879
Female	69,363	76,894	83,325	94,350	102,480	108,975	113,936	119,540	124,580	129,145
Unknown	*	*	11	66	76	93	44	33	21	29
White	83,097	91,030	97,028	107,609	115,395	122,146	129,515	136,274	144,164	150,913
Black	54,273	60,302	66,048	73,562	80,332	86,475	91,062	96,206	100,148	103,221
Native American	2,030	2,383	2,633	2,994	3,457	3,950	3,999	4,475	4,555	4,605
Asian	4,176	4,716	5,506	6,524	7,578	8,396	8,920	9,603	10,215	10,699
Other/unknown	982	1,419	2,186	4,855	5,663	5,982	4,885	4,947	4,960	5,615
†Hispanic					14,683	19,781	23,768	27,041	29,757	32,436
†Non-Hispanic					197,742	207,168	214,613	224,464	234,285	242,617
Diabetes	41,919	48,727	53,459	61,725	70,704	79,915	88,916	96,868	103,591	110,041
Hypertension	41,298	46,531	49,905	55,602	58,312	61,591	64,625	68,041	71,163	73,631
Glomerulonephritis	21,666	23,158	24,465	26,519	28,621	30,262	31,351	32,389	33,246	33,182
Cystic kidney	5,805	6,079	6,286	6,726	7,089	7,265	7,328	7,591	7,882	7,928
Other urologic	2,812	3,163	3,295	3,561	3,909	4,323	4,628	4,843	5,114	5,202
Other cause	15,913	16,656	17,082	18,529	20,893	22,538	23,982	25,362	26,800	27,688
Unknown cause	6,789	7,250	7,406	7,956	8,271	8,765	9,064	9,493	9,883	10,248
Missing cause	8,356	8,286	11,503	14,926	14,626	12,290	8,487	6,918	6,363	7,133
All	144,558	159,850	173,401	195,544	212,425	226,949	238,381	251,505	264,042	275,053

The point prevalent cohorts and associated modalities are determined on December 31 of each calendar year without applying the 60-day stable modality rule. The age of the point prevalent patient is determined as of December 31 of each calendar year.

†CMS has only started collecting Hispanic ethnicity data as of April, 1995, so Hispanic data in years prior to 1995 has been left blank.

* Values for cells with fewer than 10 patients are suppressed.

Table D.3 (continued)

Prevalent ESRD Patients: by Dialysis/Functioning Graft Status

point prevalent patients on December 31, by demographic characteristics & primary diagnosis

HEMODIALYSIS	1991	1992	1993	1994	1995	1996	1997	1998	1999	2000
0-4	24	24	38	60	80	72	67	67	86	142
5-9	51	46	51	64	73	83	86	83	95	102
10-14	131	142	134	165	179	189	183	216	230	239
15-19	557	578	597	578	668	720	758	793	804	775
20-24	1,716	1,806	1,811	1,962	2,091	2,196	2,227	2,371	2,399	2,525
25-29	3,051	3,177	3,175	3,414	3,701	4,032	4,262	4,511	4,684	4,624
30-34	4,635	4,862	5,007	5,396	5,867	6,198	6,469	6,557	6,829	7,073
35-39	6,010	6,560	6,847	7,518	8,141	8,833	9,323	9,823	10,243	10,439
40-44	7,465	8,104	8,543	9,161	10,305	11,460	12,377	13,355	13,974	14,349
45-49	8,139	8,970	9,787	11,106	12,778	14,500	15,577	16,776	17,905	18,830
50-54	9,138	10,174	11,080	12,448	13,847	15,629	17,354	19,248	21,237	23,147
55-59	11,335	12,324	13,111	14,427	16,001	17,788	19,576	21,522	23,099	24,561
60-64	14,406	15,201	16,060	17,378	19,263	21,023	22,832	24,669	25,993	27,271
65-69	17,198	19,040	20,279	21,880	23,162	24,768	26,346	27,669	28,860	29,926
70-74	15,813	17,783	19,380	21,647	23,279	25,008	27,053	28,464	29,465	30,408
75-79	11,129	12,573	13,780	15,662	17,461	19,624	21,828	23,790	25,641	27,046
80-84	5,837	6,967	7,861	8,917	9,968	11,072	12,374	13,577	14,589	16,391
85+	2,364	2,834	3,332	3,829	4,379	5,071	5,785	6,705	7,539	8,246
Missing/unknown	13	14	10	20	22	22	11	15	20	27
0-19	763	790	820	867	1,000	1,064	1,094	1,159	1,215	1,258
20-44	22,877	24,509	25,383	27,451	30,105	32,719	34,658	36,617	38,129	39,010
45-64	43,018	46,669	50,038	55,359	61,889	68,940	75,339	82,215	88,234	93,809
65-74	33,011	36,823	39,659	43,527	46,441	49,776	53,399	56,133	58,325	60,334
75+	19,330	22,374	24,973	28,408	31,808	35,767	39,987	44,072	47,769	51,683
Unknown	13	14	10	20	22	22	11	15	20	27
Male	61,716	67,993	73,191	80,761	88,767	98,282	107,278	116,108	123,816	130,818
Female	57,295	63,185	67,691	74,869	82,493	89,998	97,202	104,090	109,858	115,276
Unknown	*	*	*	*	*	*	*	13	18	27
White	65,738	71,842	76,063	83,650	90,716	98,561	107,707	116,215	124,412	132,301
Black	47,700	53,087	57,709	63,426	69,760	76,611	82,265	87,896	92,229	95,460
Native American	1,697	1,995	2,203	2,518	2,939	3,435	3,516	4,013	4,118	4,187
Asian	3,553	3,917	4,480	5,238	6,063	6,862	7,477	8,207	8,823	9,337
Other/unknown	324	338	428	800	1,787	2,819	3,523	3,880	4,110	4,836
†Hispanic					12,363	16,828	20,572	23,813	26,567	29,434
†Non-Hispanic					158,902	171,460	183,916	196,398	207,125	216,687
Diabetes	34,590	40,199	44,045	50,967	59,126	68,223	77,684	86,361	93,446	100,739
Hypertension	35,413	39,974	42,566	47,293	50,072	53,881	57,597	61,584	64,940	67,821
Glomerulonephritis	16,997	18,064	18,836	20,242	21,926	23,789	25,322	26,737	27,761	28,133
Cystic kidney	4,621	4,759	4,856	5,189	5,441	5,716	5,922	6,297	6,543	6,731
Other urologic	2,407	2,654	2,731	2,937	3,273	3,671	3,988	4,243	4,517	4,665
Other cause	12,783	13,286	13,440	14,579	16,614	18,419	20,123	21,709	23,115	24,374
Unknown cause	5,661	6,056	6,101	6,518	6,837	7,415	7,798	8,335	8,787	9,238
Missing cause	6,540	6,187	8,308	7,907	7,976	7,174	6,054	4,945	4,583	4,420
All	119,012	131,179	140,883	155,632	171,265	188,288	204,488	220,211	233,692	246,121

The point prevalent cohorts and associated modalities are determined on December 31 of each calendar year without applying the 60-day stable modality rule. The age of the point prevalent patient is determined as of December 31 of each calendar year.

†CMS has only started collecting Hispanic ethnicity data as of April, 1995, so Hispanic data in years prior to 1995 has been left blank.

* Values for cells with fewer than 10 patients are suppressed.

Table D.3 (continued)
Prevalent ESRD Patients: by Dialysis/Functioning Graft Status
point prevalent patients on December 31, by demographic characteristics & primary diagnosis

PERITONEAL DIALYSIS	1991	1992	1993	1994	1995	1996	1997	1998	1999	2000
0-4	90	90	85	109	131	161	156	174	176	202
5-9	72	68	77	84	103	114	124	106	106	122
10-14	157	155	163	162	171	195	217	215	182	201
15-19	283	286	303	292	309	352	346	326	298	298
20-24	449	516	563	576	584	571	544	500	487	437
25-29	803	853	900	942	991	992	957	898	787	760
30-34	1,273	1,379	1,522	1,545	1,570	1,521	1,373	1,305	1,206	1,141
35-39	1,532	1,714	1,846	1,951	2,025	1,949	1,835	1,668	1,641	1,516
40-44	1,813	1,964	2,101	2,222	2,470	2,450	2,368	2,100	2,029	1,912
45-49	1,815	2,055	2,283	2,512	2,782	2,826	2,674	2,565	2,372	2,324
50-54	1,775	2,085	2,290	2,625	2,841	2,946	2,780	2,651	2,661	2,659
55-59	1,899	2,105	2,327	2,519	2,827	2,874	2,869	2,702	2,651	2,576
60-64	2,203	2,518	2,560	2,745	3,061	3,016	2,823	2,545	2,463	2,392
65-69	2,476	2,752	3,099	3,202	3,276	3,184	2,902	2,632	2,546	2,358
70-74	1,868	2,206	2,553	2,770	2,769	2,707	2,533	2,418	2,241	2,078
75-79	1,107	1,295	1,410	1,548	1,652	1,654	1,669	1,550	1,473	1,423
80-84	452	523	602	725	765	739	728	668	639	651
85+	139	176	200	209	234	254	269	253	260	236
Missing/unknown	*	*	*	*	*	*	*	.	.	*
0-19	602	599	628	647	714	822	843	821	762	823
20-44	5,870	6,426	6,932	7,236	7,640	7,483	7,077	6,471	6,150	5,766
45-64	7,692	8,763	9,460	10,401	11,511	11,662	11,146	10,463	10,147	9,951
65-74	4,344	4,958	5,652	5,972	6,045	5,891	5,435	5,050	4,787	4,436
75+	1,698	1,994	2,212	2,482	2,651	2,647	2,666	2,471	2,372	2,310
Unknown	*	*	*	*	*	*	*	.	.	*
Male	10,706	12,001	13,114	13,943	14,838	14,686	13,890	12,845	12,288	11,926
Female	9,501	10,739	11,770	12,796	13,725	13,821	13,277	12,431	11,930	11,361
Unknown	*	*	*	*	.	.	*	.	.	.
White	14,229	15,950	17,107	18,131	19,084	18,715	17,855	16,559	15,893	15,266
Black	5,216	5,846	6,652	7,237	7,643	7,631	7,159	6,650	6,345	6,020
Native American	279	324	338	376	423	438	412	403	372	349
Asian	454	582	739	919	1,151	1,266	1,210	1,162	1,126	1,154
Other/unknown	30	40	50	78	262	457	532	502	482	498
†Hispanic					1,887	2,450	2,599	2,619	2,673	2,670
†Non-Hispanic					26,676	26,057	24,569	22,657	21,545	20,617
Diabetes	5,900	6,963	7,401	8,194	9,191	9,601	9,330	8,730	8,432	8,136
Hypertension	4,807	5,474	6,001	6,516	6,585	6,244	5,840	5,426	5,286	5,115
Glomerulonephritis	3,941	4,341	4,698	5,032	5,382	5,492	5,309	4,922	4,608	4,400
Cystic kidney	1,021	1,126	1,203	1,245	1,310	1,331	1,237	1,158	1,122	1,056
Other urologic	299	376	429	461	481	503	524	495	478	468
Other cause	2,576	2,764	3,004	3,085	3,360	3,352	3,236	3,093	2,992	2,836
Unknown cause	885	946	1,037	1,069	1,076	1,085	1,062	993	890	853
Missing cause	779	752	1,113	1,139	1,178	899	630	459	410	423
All	20,208	22,742	24,886	26,741	28,563	28,507	27,168	25,276	24,218	23,287

The point prevalent cohorts and associated modalities are determined on December 31 of each calendar year without applying the 60-day stable modality rule.
 †CMS has only started collecting Hispanic ethnicity data as of April, 1995, so Hispanic data in years prior to 1995 has been left blank.
 * Values for cells with fewer than 10 patients are suppressed.
 "." Zero values in this cell.

Table D.3 (continued)

Prevalent ESRD Patients: by Dialysis/Functioning Graft Status

point prevalent patients on December 31, by demographic characteristics & primary diagnosis

TRANSPLANT	1991	1992	1993	1994	1995	1996	1997	1998	1999	2000
0-4	195	213	225	221	196	222	214	216	230	239
5-9	597	599	629	623	643	652	708	695	709	669
10-14	839	904	957	1,013	1,092	1,163	1,166	1,233	1,279	1,312
15-19	1,402	1,414	1,506	1,691	1,759	1,765	1,859	1,900	1,985	2,090
20-24	2,606	2,698	2,736	2,722	2,609	2,636	2,653	2,691	2,780	2,849
25-29	4,592	4,606	4,696	4,668	4,702	4,791	4,869	4,865	4,735	4,629
30-34	6,701	7,031	7,419	7,622	7,713	7,717	7,726	7,841	7,710	7,845
35-39	7,862	8,380	8,872	9,358	9,683	10,169	10,532	10,873	10,990	11,165
40-44	7,966	8,707	9,518	10,126	10,692	11,201	11,790	12,431	12,964	13,359
45-49	6,610	7,549	8,541	9,555	10,428	11,351	12,033	12,851	13,318	14,294
50-54	5,464	6,238	7,048	7,796	8,430	9,451	10,676	11,848	12,991	14,294
55-59	4,543	5,110	5,747	6,353	6,870	7,524	8,582	9,627	10,446	11,218
60-64	3,501	3,952	4,402	4,891	5,399	6,085	6,706	7,482	8,123	8,890
65-69	1,744	2,191	2,692	3,149	3,587	4,101	4,632	5,159	5,703	6,305
70-74	568	771	978	1,219	1,467	1,731	2,104	2,491	2,814	3,338
75-79	115	136	179	241	341	437	566	716	868	1,034
80-84	19	20	26	37	42	68	80	105	155	195
85+	12	*	*	*	12	16	20	31	37	42
Missing/unknown	57	56	53	51	46	45	45	43	42	42
0-19	3,033	3,130	3,317	3,548	3,690	3,802	3,947	4,044	4,203	4,310
20-44	29,727	31,422	33,241	34,496	35,399	36,514	37,570	38,701	39,179	39,847
45-64	20,118	22,849	25,738	28,595	31,127	34,411	37,997	41,808	44,878	48,696
65-74	2,312	2,962	3,670	4,368	5,054	5,832	6,736	7,650	8,517	9,643
75+	146	165	213	286	395	521	666	852	1,060	1,271
Unknown	57	56	53	51	46	45	45	43	42	42
Male	33,319	36,326	39,609	42,639	45,254	48,366	51,900	55,387	58,119	61,637
Female	22,069	24,250	26,611	28,695	30,447	32,749	35,051	37,701	39,750	42,162
Unknown	*	*	12	10	10	10	10	10	10	10
White	42,236	46,152	50,448	53,951	57,084	60,821	64,822	69,165	72,530	76,714
Black	8,934	9,804	10,769	11,989	12,913	14,168	15,490	16,703	17,618	18,724
Native American	632	706	778	865	928	986	1,050	1,119	1,189	1,256
Asian	1,864	2,134	2,381	2,667	2,886	3,168	3,480	3,816	4,081	4,422
Other/unknown	1,727	1,788	1,856	1,872	1,900	1,982	2,119	2,295	2,461	2,693
†Hispanic					4,138	4,678	5,390	6,285	7,044	7,947
†Non-Hispanic					71,573	76,447	81,571	86,813	90,835	95,862
Diabetes	9,474	10,512	11,946	13,179	14,185	15,608	17,039	18,584	19,754	21,132
Hypertension	6,413	7,151	7,903	8,767	9,430	10,246	11,039	11,859	12,423	13,108
Glomerulonephritis	14,833	16,108	17,281	18,417	19,370	20,541	21,953	23,417	24,551	25,874
Cystic kidney	3,810	4,315	4,753	5,162	5,612	6,095	6,689	7,298	7,787	8,370
Other urologic	886	982	1,145	1,256	1,392	1,550	1,696	1,885	2,035	2,185
Other cause	8,580	9,321	10,095	10,716	11,284	11,996	12,757	13,556	14,101	14,890
Unknown cause	2,577	2,720	2,886	3,056	3,179	3,375	3,631	3,911	4,086	4,383
Missing cause	8,820	9,475	10,223	10,791	11,259	11,714	12,157	12,588	13,142	13,867
All	55,393	60,584	66,232	71,344	75,711	81,125	86,961	93,098	97,879	103,809

The point prevalent cohorts and associated modalities are determined on December 31 of each calendar year without applying the 60-day stable modality rule. The age of the point prevalent patient is determined as of December 31 of each calendar year.

†CMS has only started collecting Hispanic ethnicity data as of April, 1995, so Hispanic data in years prior to 1995 has been left blank.

* Values for cells with fewer than 10 patients are suppressed.

Table D.4
Incident ESRD Patients for Selected Years: by Age & Modality

1992	Center hemo	Center self hemo	Home hemo	CAPD	CCPD	Other PD	Uncertain dialysis	Transplant	All
0-4	17	.	.	33	*	*	43	46	148
5-9	21	.	.	13	*	*	26	59	122
10-14	85	.	.	34	*	*	45	76	245
15-19	292	*	.	73	*	*	49	75	503
20-24	698	*	.	121	*	15	67	109	1,019
25-29	1,104	*	*	187	12	14	105	173	1,598
30-34	1,513	*	.	362	*	25	121	212	2,246
35-39	2,036	*	*	416	10	28	181	217	2,891
40-44	2,424	*	*	490	15	19	231	237	3,425
45-49	2,730	*	*	496	27	37	226	184	3,706
50-54	3,213	*	*	536	24	43	279	132	4,233
55-59	4,037	*	*	613	21	53	323	107	5,163
60-64	5,345	*	*	740	31	71	409	71	6,674
65-69	6,919	*	10	582	74	127	827	30	8,576
70-74	6,541	10	*	509	57	140	822	*	8,089
75-79	4,901	*	*	326	57	87	601	*	5,983
80-84	2,689	*	.	136	23	41	324	*	3,219
85+	1,051	.	.	53	*	15	146	*	1,273
Unknown	*	.	.	*	.	.	*	.	14
0-19	415	*	.	153	20	10	163	256	1,018
20-44	7,775	16	*	1,576	53	101	705	948	11,179
45-64	15,325	17	11	2,385	103	204	1,237	494	19,776
65-74	13,460	17	12	1,091	131	267	1,649	38	16,665
75+	8,641	12	*	515	86	143	1,071	*	10,475
Unknown	*	.	.	*	.	.	*	.	14
All	45,623	63	31	5,722	393	725	4,830	1,740	59,127
1996									
0-4	29	.	*	12	96	10	*	49	204
5-9	40	.	.	13	52	*	*	43	154
10-14	97	.	*	19	90	*	18	59	289
15-19	353	.	*	52	71	*	24	93	604
20-24	778	.	18	102	35	*	43	78	1,059
25-29	1,256	*	26	193	75	*	58	99	1,713
30-34	1,834	*	34	306	89	14	78	146	2,504
35-39	2,519	*	67	371	137	12	114	189	3,413
40-44	3,255	*	77	485	172	24	160	211	4,386
45-49	4,123	*	91	575	198	17	218	196	5,419
50-54	4,516	*	111	589	194	38	244	156	5,853
55-59	5,501	*	122	662	170	37	311	105	6,915
60-64	6,976	*	134	700	207	38	395	87	8,543
65-69	8,374	12	88	608	198	96	864	36	10,276
70-74	8,759	*	89	625	157	84	902	*	10,629
75-79	7,063	14	62	377	91	58	736	*	8,403
80-84	4,177	*	53	174	48	46	432	.	4,931
85+	1,878	*	28	69	23	19	214	.	2,232
Unknown	*	*	*	*
0-19	519	.	10	96	309	19	54	244	1,251
20-44	9,642	10	222	1,457	508	60	453	723	13,075
45-64	21,116	19	458	2,526	769	130	1,168	544	26,730
65-74	17,133	19	177	1,233	355	180	1,766	42	20,905
75+	13,118	16	143	620	162	123	1,382	*	15,566
Unknown	*	*	*	*
All	61,530	64	1,010	5,932	2,103	512	4,824	1,556	77,531

The incident cohorts and associated modalities are determined at the time of ESRD initiation without applying the 60-day stable modality rule.
 The age of the incident patient is determined as of the date of ESRD initiation.
 * Values for cells with fewer than 10 patients are suppressed.
 "." Zero values in this cell.

Table D.4 (continued)

Incident ESRD Patients for Selected Years: by Age & Modality

2000	Center hemo	Center self hemo	Home hemo	CAPD	CCPD	Other PD	Uncertain dialysis	Transplant	All
0-4	78	.	*	16	119	*	*	34	254
5-9	55	.	.	*	57	.	.	45	163
10-14	134	.	.	16	82	*	*	82	321
15-19	388	.	*	24	78	*	20	102	620
20-24	826	.	*	54	46	*	70	75	1,075
25-29	1,410	.	*	128	70	*	111	121	1,844
30-34	2,025	.	*	206	79	*	137	188	2,647
35-39	2,763	.	*	247	129	*	174	210	3,534
40-44	3,948	*	13	340	167	*	216	273	4,967
45-49	5,302	*	19	445	179	14	306	279	6,546
50-54	6,777	*	22	526	215	19	350	263	8,176
55-59	7,876	.	25	532	219	14	342	216	9,224
60-64	9,000	*	34	547	256	22	275	142	10,277
65-69	10,287	*	35	527	229	41	236	93	11,452
70-74	11,220	*	32	478	194	47	270	42	12,292
75-79	10,604	*	54	385	149	41	258	*	11,497
80-84	6,888	*	43	171	77	27	172	*	7,384
85+	3,683	.	14	75	32	14	69	.	3,887
Unknown	30	.	.	*	.	.	.	*	32
0-19	655	.	*	62	336	*	29	263	1,358
20-44	10,972	*	30	975	491	23	708	867	14,067
45-64	28,955	*	100	2,050	869	69	1,273	900	34,223
65-74	21,507	13	67	1,005	423	88	506	135	23,744
75+	21,175	*	111	631	258	82	499	*	22,768
Unknown	30	.	.	*	.	.	.	*	32
All	83,294	29	312	4,724	2,377	271	3,015	2,170	96,192

The incident cohorts and associated modalities are determined at the time of ESRD initiation without applying the 60-day stable modality rule.

The age of the incident patient is determined as of the date of ESRD initiation.

* Values for cells with fewer than 10 patients are suppressed.

." Zero values in this cell.

Table D.5
Prevalent ESRD Patients for Selected Years: by Age & Modality
point prevalent patients on December 31

	Center hemo	Center self hemo	Home hemo	CAPD	CCPD	Other PD	Uncertain dialysis	Transplant	All
1992									
0-4	24	.	.	48	42	12	44	213	383
5-9	46	.	.	33	35	10	38	599	761
10-14	138	.	*	74	81	18	60	904	1,279
15-19	566	*	*	178	108	21	116	1,414	2,415
20-24	1,754	28	24	412	104	*	165	2,698	5,191
25-29	3,096	33	48	742	111	15	249	4,606	8,900
30-34	4,713	66	83	1,192	187	34	285	7,031	13,591
35-39	6,356	75	129	1,515	199	43	381	8,380	17,078
40-44	7,841	88	175	1,729	235	42	420	8,707	19,237
45-49	8,706	86	178	1,818	237	37	440	7,549	19,051
50-54	9,933	76	165	1,823	262	57	470	6,238	19,024
55-59	12,089	72	163	1,875	230	60	506	5,110	20,105
60-64	14,955	75	171	2,235	283	66	608	3,952	22,345
65-69	18,758	91	191	2,430	322	84	515	2,191	24,582
70-74	17,574	64	145	1,952	254	78	442	771	21,280
75-79	12,452	32	89	1,145	150	58	281	136	14,343
80-84	6,895	19	53	481	42	21	169	20	7,700
85+	2,810	*	19	150	26	16	55	*	3,090
Unknown	14	.	.	*	*	.	*	56	79
0-19	774	*	12	333	266	61	258	3,130	4,838
20-44	23,760	290	459	5,590	836	140	1,500	31,422	63,997
45-64	45,683	309	677	7,751	1,012	220	2,024	22,849	80,525
65-74	36,332	155	336	4,382	576	162	957	2,962	45,862
75+	22,157	56	161	1,776	218	95	505	165	25,133
Unknown	14	.	.	*	*	.	*	56	79
All	128,720	814	1,645	19,833	2,909	678	5,251	60,584	220,434
1996									
0-4	69	*	*	24	137	11	33	222	499
5-9	80	.	*	21	93	*	25	652	877
10-14	176	*	11	50	145	11	66	1,163	1,624
15-19	702	*	16	135	217	15	100	1,765	2,952
20-24	2,171	*	16	353	218	11	190	2,636	5,604
25-29	3,979	19	34	648	344	18	341	4,791	10,174
30-34	6,098	30	70	1,029	492	31	475	7,717	15,942
35-39	8,650	49	134	1,353	596	48	681	10,169	21,680
40-44	11,227	51	182	1,680	770	65	782	11,201	25,958
45-49	14,219	71	210	1,988	838	70	807	11,351	29,554
50-54	15,343	63	223	2,053	893	59	782	9,451	28,867
55-59	17,539	53	196	2,011	863	54	864	7,524	29,104
60-64	20,757	48	218	2,113	903	76	1,025	6,085	31,225
65-69	24,482	51	235	2,167	1,017	98	1,036	4,101	33,187
70-74	24,714	59	235	1,905	802	67	910	1,731	30,423
75-79	19,449	20	155	1,142	512	53	690	437	22,458
80-84	10,944	20	108	544	195	22	411	68	12,312
85+	4,991	*	76	196	58	12	205	16	5,558
Unknown	22	.	.	*	.	.	*	45	76
0-19	1,027	*	32	230	592	40	224	3,802	5,952
20-44	32,125	158	436	5,063	2,420	173	2,469	36,514	79,358
45-64	67,858	235	847	8,165	3,497	259	3,478	34,411	118,750
65-74	49,196	110	470	4,072	1,819	165	1,946	5,832	63,610
75+	35,384	44	339	1,882	765	87	1,306	521	40,328
Unknown	22	.	.	*	.	.	*	45	76
All	185,612	552	2,124	19,414	9,093	724	9,430	81,125	308,074

The point prevalent cohorts and associated modalities are determined on December 31 of each calendar year without applying the 60-day stable modality rule. The age of the point prevalent patient is determined as of December 31 of each calendar year.
 *Values for cells with fewer than 10 patients are suppressed.
 "." Zero values in this cell.

Table D.5 (continued)

Prevalent ESRD Patients for Selected Years: by Age & Modality

point prevalent patients on December 31

2000	Center hemo	Center self hemo	Home hemo	CAPD	CCPD	Other PD	Uncertain dialysis	Transplant	All
0-4	140	.	*	25	177	*	11	239	603
5-9	97	.	*	17	105	*	11	669	909
10-14	234	.	*	29	172	10	31	1,312	1,793
15-19	765	*	*	57	241	16	76	2,090	3,255
20-24	2,500	*	22	190	247	10	137	2,849	5,958
25-29	4,576	*	42	344	416	13	216	4,629	10,242
30-34	7,011	10	52	607	534	17	285	7,845	16,361
35-39	10,313	23	103	820	696	17	395	11,165	23,532
40-44	14,182	31	136	1,061	851	40	484	13,359	30,144
45-49	18,638	33	159	1,268	1,056	38	542	14,294	36,028
50-54	22,931	36	180	1,514	1,145	37	637	14,294	40,774
55-59	24,371	34	156	1,464	1,112	46	640	11,218	39,041
60-64	27,140	20	111	1,356	1,036	46	519	8,890	39,118
65-69	29,784	28	114	1,246	1,112	45	365	6,305	38,999
70-74	30,257	29	122	1,088	990	53	308	3,338	36,185
75-79	26,919	12	115	753	670	33	280	1,034	29,816
80-84	16,315	14	62	326	325	16	163	195	17,416
85+	8,210	*	35	141	95	*	86	42	8,617
Unknown	27	.	.	*	.	.	*	42	71
0-19	1,236	*	20	128	695	40	129	4,310	6,560
20-44	38,582	73	355	3,022	2,744	97	1,517	39,847	86,237
45-64	93,080	123	606	5,602	4,349	167	2,338	48,696	154,961
65-74	60,041	57	236	2,334	2,102	98	673	9,643	75,184
75+	51,444	27	212	1,220	1,090	56	529	1,271	55,849
Unknown	27	.	.	*	.	.	*	42	71
All	244,410	282	1,429	12,307	10,980	458	5,187	103,809	378,862

The point prevalent cohorts and associated modalities are determined on December 31 of each calendar year without applying the 60-day stable modality rule. The age of the point prevalent patient is determined as of December 31 of each calendar year.

* Values for cells with fewer than 10 patients are suppressed.

." Zero values in this cell.

Table D.6

Incident ESRD Patients for Selected Years: by Gender, Race, Ethnicity, & Primary Diagnosis; Grouped by Modality

incident patients

	Center hemo	Center self hemo	Home hemo	CAPD	CCPD	Other PD	Uncertain dialysis	Transplant	All
1992									
Male	24,229	43	21	3,050	226	385	2,446	961	31,361
Female	21,394	20	10	2,671	167	340	2,383	775	27,760
Unknown	.	.	.	*	.	.	*	*	*
White	28,685	42	11	4,143	297	536	3,086	1,315	38,115
Black	14,892	17	16	1,365	83	168	1,047	252	17,840
Native American	714	*	*	67	*	18	52	14	877
Asian	1,235	*	.	140	*	*	126	81	1,592
Other/unknown	97	.	.	*	.	*	519	78	703
Diabetes	16,745	19	10	2,301	148	288	1,625	254	21,390
Hypertension	14,625	15	*	1,294	107	218	1,361	94	17,722
Glomerulonephritis	4,917	*	*	892	58	70	432	215	6,592
Cystic kidney	1,150	*	*	260	11	13	88	103	1,630
Other urologic	919	*	*	78	*	15	124	30	1,176
Other cause	4,154	*	*	547	36	62	399	175	5,382
Unknown cause	2,061	*	.	208	12	40	185	54	2,567
Missing cause	1,052	*	*	142	15	19	616	815	2,668
All	45,623	63	31	5,722	393	725	4,830	1,740	59,127
1996									
Male	32,882	38	567	3,080	1,182	268	2,466	932	41,415
Female	28,645	26	443	2,852	921	244	2,339	624	36,094
Unknown	*	19	.	22
White	36,949	44	174	4,103	1,453	333	3,101	1,124	47,281
Black	19,960	11	584	1,316	470	145	1,334	337	24,157
Native American	981	*	155	94	41	10	54	11	1,348
Asian	2,132	*	87	273	71	15	125	50	2,760
Other/unknown	1,508	.	10	146	68	*	210	34	1,985
†Hispanic	7,139	*	319	604	244	62	507	86	8,963
†Non-Hispanic	54,391	62	691	5,328	1,859	450	4,317	1,470	68,568
Diabetes	26,250	27	398	2,684	900	214	1,872	316	32,661
Hypertension	16,586	10	291	1,113	303	141	1,219	68	19,731
Glomerulonephritis	6,123	15	116	978	336	49	365	253	8,235
Cystic kidney	1,266	*	30	249	109	12	63	146	1,877
Other urologic	1,256	*	15	102	33	*	92	51	1,560
Other cause	6,885	*	122	561	324	59	389	228	8,575
Unknown cause	2,468	*	33	196	85	16	140	57	2,996
Missing cause	696	.	*	49	13	12	684	437	1,896
All	61,530	64	1,010	5,932	2,103	512	4,824	1,556	77,531

The incident cohorts and associated modalities are determined at the time of ESRD initiation without applying the 60-day stable modality rule.
 †CMS has only started collecting Hispanic ethnicity data as of April, 1995, so Hispanic data in years prior to 1995 has been left blank.
 * Values for cells with fewer than 10 patients are suppressed.
 "." Zero values in this cell.

Table D.6 (continued)

Incident ESRD Patients for Selected Years: by Gender, Race, Ethnicity, & Primary Diagnosis; Grouped by Modality
incident patients

2000	Center hemo	Center self hemo	Home hemo	CAPD	CCPD	Other PD	Uncertain dialysis	Transplant	All
Male	44,487	19	174	2,499	1,304	134	1,724	1,232	51,573
Female	38,799	10	138	2,225	1,073	137	1,291	938	44,611
Unknown	*	*
White	52,439	19	212	3,410	1,734	180	1,766	1,788	61,548
Black	24,819	*	88	956	462	75	925	208	27,540
Native American	1,063	.	*	66	28	*	45	12	1,222
Asian	2,825	*	*	191	71	*	105	78	3,284
Other/unknown	2,148	.	*	101	82	*	174	84	2,598
†Hispanic	10,252	*	25	455	318	22	81	112	11,266
†Non-Hispanic	73,042	28	287	4,269	2,059	249	2,934	2,058	84,926
Diabetes	37,574	*	131	2,159	956	127	538	278	41,772
Hypertension	22,561	12	85	1,008	394	59	338	109	24,566
Glomerulonephritis	6,636	*	14	659	377	18	88	309	8,102
Cystic kidney	1,579	*	*	237	93	*	24	205	2,144
Other urologic	1,409	.	*	75	52	*	20	63	1,630
Other cause	9,243	*	54	398	370	40	84	252	10,444
Unknown cause	3,443	*	18	146	90	11	47	85	3,841
Missing cause	849	*	*	42	45	*	1,876	869	3,693
All	83,294	29	312	4,724	2,377	271	3,015	2,170	96,192

The incident cohorts and associated modalities are determined at the time of ESRD initiation without applying the 60-day stable modality rule.

†CMS has only started collecting Hispanic ethnicity data as of April, 1995, so Hispanic data in years prior to 1995 has been left blank.

* Values for cells with fewer than 10 patients are suppressed.

." Zero values in this cell.

Table D.7
Prevalent ESRD Patients for Selected Years: by Gender, Race, Ethnicity, & Primary Diagnosis; Grouped by Modality
point prevalent patients on December 31

	Center hemo	Center self hemo	Home hemo	CAPD	CCPD	Other PD	Uncertain dialysis	Transplant	All
1992									
Male	66,592	448	953	10,468	1,533	349	2,607	36,326	119,276
Female	62,127	366	692	9,363	1,376	329	2,641	24,250	101,144
Unknown	*	.	.	*	.	.	*	*	14
White	70,336	374	1,132	13,902	2,048	449	2,789	46,152	137,182
Black	52,305	389	393	5,088	758	214	1,155	9,804	70,106
Native American	1,897	12	86	297	27	*	58	706	3,089
Asian	3,849	36	32	508	74	*	212	2,134	6,850
Other/unknown	333	*	*	38	*	*	1,037	1,788	3,207
Diabetes	39,719	160	320	6,162	801	221	1,344	10,512	59,239
Hypertension	39,452	215	307	4,802	672	146	937	7,151	53,682
Glomerulonephritis	17,511	171	382	3,777	564	91	662	16,108	39,266
Cystic kidney	4,593	36	130	946	180	18	176	4,315	10,394
Other urologic	2,600	17	37	329	47	17	116	982	4,145
Other cause	12,948	111	227	2,347	417	112	494	9,321	25,977
Unknown cause	5,936	41	79	823	123	42	206	2,720	9,970
Missing cause	5,961	63	163	647	105	31	1,316	9,475	17,761
All	128,720	814	1,645	19,833	2,909	678	5,251	60,584	220,434
1996									
Male	96,771	333	1,178	9,805	4,881	359	4,554	48,366	166,247
Female	88,833	219	946	9,609	4,212	365	4,791	32,749	141,724
Unknown	*	85	10	103
White	97,076	277	1,208	12,628	6,087	412	4,458	60,821	182,967
Black	75,634	242	735	5,267	2,364	249	1,984	14,168	100,643
Native American	3,316	17	102	323	115	*	68	986	4,936
Asian	6,782	15	65	880	386	28	240	3,168	11,564
Other/unknown	2,804	*	14	316	141	26	2,680	1,982	7,964
†Hispanic	16,621	15	192	1,621	829	70	433	4,678	24,459
†Non-Hispanic	168,991	537	1,932	17,793	8,264	654	8,997	76,447	283,615
Diabetes	67,533	121	569	6,679	2,922	270	1,821	15,608	95,523
Hypertension	53,235	142	504	4,312	1,932	157	1,309	10,246	71,837
Glomerulonephritis	23,295	120	374	3,744	1,748	107	874	20,541	50,803
Cystic kidney	5,557	33	126	887	444	26	192	6,095	13,360
Other urologic	3,607	15	49	333	170	17	132	1,550	5,873
Other cause	18,089	71	259	2,096	1,256	95	672	11,996	34,534
Unknown cause	7,294	20	101	749	336	25	240	3,375	12,140
Missing cause	7,002	30	142	614	285	27	4,190	11,714	24,004
All	185,612	552	2,124	19,414	9,093	724	9,430	81,125	308,074

The point prevalent cohorts and associated modalities are determined on December 31 of each calendar year without applying the 60-day stable modality rule.
 †CMS has only started collecting Hispanic ethnicity data as of April, 1995, so Hispanic data in years prior to 1995 has been left blank.
 * Values for cells with fewer than 10 patients are suppressed.
 "." Zero values in this cell.

Table D.7 (continued)

Prevalent ESRD Patients for Selected Years: by Gender, Race, Ethnicity, & Primary Diagnosis; Grouped by Modality
point prevalent patients on December 31

2000	Center hemo	Center self hemo	Home hemo	CAPD	CCPD	Other PD	Uncertain dialysis	Transplant	All
Male	129,842	159	817	6,072	5,854	217	2,918	61,637	207,516
Female	114,541	123	612	6,235	5,126	241	2,267	42,162	171,307
Unknown	27	*	10	39
White	131,291	138	872	8,007	7,259	260	3,086	76,714	227,627
Black	94,862	124	474	3,179	2,841	167	1,574	18,724	121,945
Native American	4,163	*	21	203	146	*	60	1,256	5,861
Asian	9,274	15	48	662	492	16	192	4,422	15,121
Other/unknown	4,820	*	14	256	242	*	275	2,693	8,308
†Hispanic	29,333	*	93	1,321	1,349	33	299	7,947	40,383
†Non-Hispanic	215,077	274	1,336	10,986	9,631	425	4,888	95,862	338,479
Diabetes	100,302	67	370	4,497	3,639	185	981	21,132	131,173
Hypertension	67,417	79	325	2,764	2,351	83	612	13,108	86,739
Glomerulonephritis	27,809	57	267	2,252	2,148	70	579	25,874	59,056
Cystic kidney	6,641	14	76	603	453	*	132	8,370	16,298
Other urologic	4,624	*	35	229	239	*	62	2,185	7,387
Other cause	24,126	26	222	1,317	1,519	74	404	14,890	42,578
Unknown cause	9,136	21	81	447	406	20	137	4,383	14,631
Missing cause	4,355	12	53	198	225	10	2,280	13,867	21,000
All	244,410	282	1,429	12,307	10,980	458	5,187	103,809	378,862

The point prevalent cohorts and associated modalities are determined on December 31 of each calendar year without applying the 60-day stable modality rule.

†CMS has only started collecting Hispanic ethnicity data as of April, 1995, so Hispanic data in years prior to 1995 has been left blank.

* Values for cells with fewer than 10 patients are suppressed.

." Zero values in this cell.

Table D.8

Treatment Status Two Years after First ESRD Service

incident patients, 1996-1998 combined, by gender, age at first service, & modality at 90 days

ALL	Treatment status two years after first ESRD service (percent of patients)						Patient count
	Center hemo	Home hemo	CAPD & CCPD	Oth/unk dialysis	Transplant	Death	
Age at first service							
0-19	23.8	0.3	14.3	4.0	52.6	5.0	2,526
20-44	50.8	0.3	10.3	2.2	23.7	12.7	31,193
45-64	56.4	0.3	7.9	1.7	9.2	24.4	70,285
65-74	51.4	0.2	5.0	1.0	1.5	40.8	56,475
75+	42.3	0.2	2.4	0.8	0.1	54.3	42,263
Modality on day 90							
Center hemodialysis	58.0	0.1	1.7	0.8	5.1	34.4	166,938
Home hemodialysis	22.0	22.9	4.1	1.6	6.0	43.4	1,126
CAPD/CCPD	16.0	0.3	40.0	1.7	13.3	28.7	24,729
Other/unknown dialysis	34.7	0.2	4.9	17.5	5.1	37.6	5,714
Transplant	1.6	.	0.5	1.6	93.5	2.8	4,235
Interaction of age at first service & modality on day 90							
0-19	Center hemodialysis 48.6 . 4.2 2.8 38.0 6.4 1,025						
	Home hemodialysis 11.1 22.2 16.7 5.6 33.3 11.1 18						
	CAPD/CCPD 8.4 0.3 32.5 4.8 48.2 5.7 912						
	Other/unknown dialysis 15.7 . 15.7 24.1 42.2 2.4 83						
	Transplant 2.5 . 1.2 1.4 94.1 0.8 488						
20-44	Center hemodialysis 63.9 0.1 3.5 1.4 17.1 14.1 22,888						
	Home hemodialysis 35.1 23.0 10.1 3.4 12.8 15.5 148						
	CAPD/CCPD 16.4 0.3 42.6 2.6 27.6 10.5 5,503						
	Other/unknown dialysis 34.4 . 6.2 25.6 16.7 17.0 706						
	Transplant 1.7 . 0.6 1.8 93.9 2.0 1,948						
45-64	Center hemodialysis 65.4 0.1 2.0 1.2 6.4 24.8 56,905						
	Home hemodialysis 25.7 29.2 5.0 2.5 10.2 27.4 401						
	CAPD/CCPD 17.2 0.3 43.9 1.8 12.4 24.5 9,773						
	Other/unknown dialysis 39.3 0.4 6.0 20.0 7.6 26.7 1,618						
	Transplant 1.2 . 0.3 1.4 92.8 4.3 1,588						
65-74	Center hemodialysis 56.6 0.1 1.2 0.5 1.0 40.5 48,405						
	Home hemodialysis 20.4 18.4 2.4 0.7 0.3 57.8 294						
	CAPD/CCPD 16.3 0.3 38.1 0.8 2.2 42.4 5,706						
	Other/unknown dialysis 35.2 0.1 4.8 15.3 0.9 43.7 1,871						
	Transplant 2.5 . . 0.5 93.0 4.0 199						
75+	Center hemodialysis 45.1 0.1 0.5 0.3 0.1 53.9 37,715						
	Home hemodialysis 11.7 18.5 0.4 . . 69.4 265						
	CAPD/CCPD 13.3 0.1 28.0 0.3 0.3 58.0 2,835						
	Other/unknown dialysis 30.1 0.2 2.5 13.1 0.1 54.0 1,436						
	Transplant 100.0 . 12						
All	50.8	0.3	6.4	1.4	7.9	33.2	202,742

The incident cohorts and associated modalities are determined at the time of ESRD initiation without applying the 60-day stable modality rule.

The age of the incident patient is determined as of the date of ESRD initiation.

Includes only incident patients with unknown age & patients who have Medicare as their primary payor, are receiving their first ESRD service, & survive the first 90 days.

." Zero values in this cell.

Table D.8 (continued)

Treatment Status Two Years after First ESRD Service

incident patients, 1996-1998 combined, by gender, age at first service, & modality at 90 days

MALE	Treatment status two years after first ESRD service (percent of patients)						Patient count
	Center hemo	Home hemo	CAPD & CCPD	Oth/unk dialysis	Transplant	Death	
Age at first service							
0-19	23.0	0.1	10.7	4.2	57.7	4.3	1,394
20-44	52.1	0.2	8.8	1.9	23.6	13.3	18,630
45-64	55.6	0.4	7.7	1.8	10.4	24.2	38,146
65-74	49.7	0.3	5.6	0.9	1.9	41.6	28,450
75+	41.7	0.2	2.7	0.7	0.1	54.6	21,798
Modality on day 90							
Center hemodialysis	57.5	0.1	1.6	0.8	6.0	33.9	89,083
Home hemodialysis	22.4	23.4	3.7	1.6	7.2	41.8	625
CAPD/CCPD	15.7	0.3	39.6	1.7	13.9	28.9	13,282
Other/unknown dialysis	34.3	0.2	5.1	17.1	6.0	37.3	2,946
Transplant	1.5	.	0.2	1.5	94.0	2.7	2,482
Interaction of age at first service & modality on day 90							
0-19	Center hemodialysis 47.1 . 2.3 2.3 42.2 6.2 569						
	Home hemodialysis 18.2 9.1 9.1 9.1 45.5 9.1 11						
	CAPD/CCPD 7.6 0.2 27.8 5.8 53.7 4.9 449						
	Other/unknown dialysis 17.6 . 15.7 27.5 37.3 2.0 51						
	Transplant 2.2 . 0.6 1.6 95.2 0.3 314						
20-44	Center hemodialysis 64.0 0.1 2.8 1.3 17.4 14.4 14,109						
	Home hemodialysis 37.9 21.8 10.3 4.6 11.5 13.8 87						
	CAPD/CCPD 16.5 0.2 40.9 2.3 28.3 11.8 2,947						
	Other/unknown dialysis 36.4 . 5.8 24.2 16.4 17.2 396						
	Transplant 1.3 . 0.3 1.5 94.9 2.1 1,091						
45-64	Center hemodialysis 64.6 0.2 2.0 1.2 7.4 24.7 30,948						
	Home hemodialysis 24.8 30.1 4.1 2.0 11.8 27.2 246						
	CAPD/CCPD 16.3 0.3 43.9 1.8 13.1 24.6 5,140						
	Other/unknown dialysis 36.6 0.5 5.8 21.3 9.5 26.3 863						
	Transplant 1.4 . . 1.7 92.7 4.2 949						
65-74	Center hemodialysis 55.1 0.1 1.4 0.5 1.3 41.6 24,126						
	Home hemodialysis 19.4 22.2 2.1 . 0.7 55.6 144						
	CAPD/CCPD 16.0 0.3 38.6 1.0 2.7 41.4 3,153						
	Other/unknown dialysis 35.0 0.2 5.2 13.6 1.1 45.0 907						
	Transplant 2.5 . . 0.8 93.3 3.3 120						
75+	Center hemodialysis 44.6 0.1 0.5 0.3 0.1 54.4 19,331						
	Home hemodialysis 11.7 14.6 . . 73.7 137						
	CAPD/CCPD 14.1 0.2 28.9 0.2 0.3 56.4 1,593						
	Other/unknown dialysis 30.7 0.1 2.9 11.8 0.1 54.3 729						
	Transplant 100.0 . *						
All	50.2	0.3	6.4	1.4	9.0	32.7	108,418

Includes only incident patients with unknown age & patients who have Medicare as their primary payor, are receiving their first ESRD service, & survive the first 90 days.

* Values for cells with fewer than 10 patients are suppressed.

." Zero values in this cell.

Table D.8 (continued)

Treatment Status Two Years after First ESRD Service

incident patients, 1996-1998 combined, by gender, age at first service, & modality at 90 days

FEMALE	Treatment status two years after first ESRD service (percent of patients)						Patient count	
	Center hemo	Home hemo	CAPD & CCPD	Oth/unk dialysis	Transplant	Death		
Age at first service								
0-19	24.9	0.4	18.7	3.7	46.4	5.8	1,132	
20-44	48.9	0.3	12.6	2.5	23.8	11.9	12,563	
45-64	57.4	0.3	8.2	1.6	7.9	24.6	32,139	
65-74	53.2	0.2	4.4	1.1	1.1	40.0	28,025	
75+	42.9	0.2	2.1	0.8	0.0	53.9	20,465	
Modality on day 90								
Center hemodialysis	58.5	0.1	1.7	0.8	4.0	34.8	77,855	
Home hemodialysis	21.6	22.4	4.6	1.6	4.4	45.5	501	
CAPD/CCPD	16.4	0.2	40.5	1.7	12.7	28.4	11,447	
Other/unknown dialysis	35.2	0.1	4.7	17.9	4.2	37.8	2,768	
Transplant	1.8	.	1.0	1.7	92.7	2.9	1,753	
Interaction of age at first service & modality on day 90								
0-19	Center hemodialysis	50.4	.	6.6	3.5	32.7	6.8	456
	Home hemodialysis	.	42.9	28.6	.	14.3	14.3	*
	CAPD/CCPD	9.3	0.4	36.9	3.9	43.0	6.5	463
	Other/unknown dialysis	12.5	.	15.6	18.8	50.0	3.1	32
	Transplant	2.9	.	2.3	1.1	92.0	1.7	174
20-44	Center hemodialysis	63.6	0.1	4.7	1.5	16.5	13.6	8,779
	Home hemodialysis	31.1	24.6	9.8	1.6	14.8	18.0	61
	CAPD/CCPD	16.4	0.3	44.6	2.9	26.8	9.0	2,556
	Other/unknown dialysis	31.9	.	6.8	27.4	17.1	16.8	310
	Transplant	2.2	.	1.1	2.3	92.6	1.8	857
45-64	Center hemodialysis	66.4	0.1	2.1	1.1	5.2	25.0	25,957
	Home hemodialysis	27.1	27.7	6.5	3.2	7.7	27.7	155
	CAPD/CCPD	18.2	0.2	43.9	1.8	11.5	24.4	4,633
	Other/unknown dialysis	42.4	0.3	6.2	18.5	5.4	27.2	755
	Transplant	0.9	.	0.6	1.1	93.0	4.4	639
65-74	Center hemodialysis	58.1	0.1	1.0	0.5	0.8	39.5	24,279
	Home hemodialysis	21.3	14.7	2.7	1.3	.	60.0	150
	CAPD/CCPD	16.7	0.2	37.5	0.6	1.5	43.6	2,553
	Other/unknown dialysis	35.5	.	4.5	16.9	0.7	42.4	964
	Transplant	2.5	.	.	.	92.4	5.1	79
75+	Center hemodialysis	45.7	0.1	0.4	0.3	0.0	53.4	18,384
	Home hemodialysis	11.7	22.7	0.8	.	.	64.8	128
	CAPD/CCPD	12.2	.	27.0	0.4	0.2	60.1	1,242
	Other/unknown dialysis	29.4	0.3	2.1	14.4	.	53.7	707
	Transplant	100.0	.	*
All		51.5	0.2	6.5	1.4	6.7	33.6	94,324

Includes only incident patients with unknown age & patients who have Medicare as their primary payor, are receiving their first ESRD service, & survive the first 90 days.

* Values for cells with fewer than 10 patients are suppressed.

." Zero values in this cell.

Table D.9

Prevalent ESRD Patients for Selected Years: by Patient Vintage

point prevalent patients on December 31, by year-end treatment modality & number of years patient has had ESRD (vintage)

1992	Functioning Dialysis graft	Center hemo	Center self hemo	Home hemo	CAPD	CCPD	Other PD	Uncertain dialysis	All	
0	49,458	2,785	38,992	141	221	6,856	830	282	2,136	52,243
1	32,500	5,048	25,516	98	208	4,632	696	144	1,206	37,548
2	21,768	6,008	17,533	90	169	2,723	496	65	692	27,776
3	14,229	6,434	11,907	72	112	1,620	242	52	224	20,663
4	9,821	5,914	8,211	63	105	1,026	156	34	226	15,735
5	6,786	5,171	5,673	52	94	693	111	20	143	11,957
6	5,084	4,656	4,245	40	90	508	82	20	99	9,740
7	3,868	4,009	3,240	44	62	395	55	12	60	7,877
8	3,098	3,574	2,570	32	59	304	53	11	69	6,672
9	2,519	3,047	2,109	23	54	253	38	*	37	5,566
10	2,051	2,880	1,710	25	61	168	38	10	39	4,931
11	1,685	2,353	1,393	22	56	136	29	*	46	4,038
12	1,361	1,851	1,137	21	47	106	18	*	29	3,212
13	1,206	1,566	1,009	15	51	81	20	*	24	2,772
14	1,013	1,400	833	25	55	65	11	*	23	2,413
15	1,386	1,423	1,159	16	83	90	15	*	21	2,809
16	654	726	520	*	48	47	*	*	26	1,380
17	467	493	338	*	24	39	*	*	49	960
18	377	432	262	12	24	27	*	*	48	809
19	292	408	188	*	13	32	*	.	49	700
20	41	123	29	.	*	*	*	.	*	164
21	31	93	23	.	*	*	*	.	*	124
22	19	61	15	.	.	*	*	.	*	80
23	14	44	11	.	*	.	.	.	*	58
24	89	55	73	.	*	12	*	*	*	144
25	*	*	*	.	.	*	.	.	*	14
26	*	*	.	.	.	*	.	.	.	*
27	*	*	*	.	.	*	.	.	.	10
28	21	10	17	.	.	*	.	*	.	31
All	159,850	60,584	128,720	814	1,645	19,833	2,909	678	5,251	220,434
1996										
0	65,142	2,706	53,395	87	654	6,038	2,812	204	1,952	67,848
1	45,887	5,280	37,311	86	281	4,429	2,213	192	1,375	51,167
2	36,444	6,938	27,351	71	210	2,999	1,450	155	4,208	43,382
3	20,263	7,374	17,100	45	160	1,705	753	42	458	27,637
4	14,464	7,421	12,339	41	126	1,163	461	33	301	21,885
5	10,273	6,774	8,810	23	81	760	364	23	212	17,047
6	7,283	6,356	6,310	22	64	505	205	12	165	13,639
7	5,466	5,767	4,759	16	49	372	151	11	108	11,233
8	4,132	5,016	3,537	18	45	271	136	10	115	9,148
9	3,111	4,217	2,622	15	49	215	112	*	92	7,328
10	2,492	3,735	2,119	17	45	172	69	13	57	6,227
11	2,123	3,143	1,798	13	35	141	78	*	54	5,266
12	1,745	2,762	1,485	13	31	119	50	*	41	4,507
13	1,483	2,417	1,243	*	30	110	49	*	42	3,900
14	1,282	2,248	1,051	15	45	91	44	*	34	3,530
15	1,007	1,876	869	11	24	50	21	*	29	2,883
16	879	1,499	724	10	20	65	29	*	27	2,378
17	791	1,282	642	*	39	58	27	*	18	2,073
18	627	1,154	506	12	38	36	14	*	20	1,781
19	788	1,178	665	*	45	35	15	*	20	1,966
20	396	585	330	*	16	22	*	.	18	981
21	296	403	223	*	13	20	*	.	28	699
22	233	354	167	*	14	11	11	.	22	587
23	201	328	150	*	*	11	*	.	25	529
24	38	101	27	.	*	*	*	.	*	139
25	25	72	16	.	*	*	*	.	.	97
26	19	41	15	.	.	*	.	.	*	60
27	*	33	*	.	.	*	*	.	.	42
28	36	39	31	.	.	*	.	.	*	75
29	*	*	*	.	*	10
30	*	*	*	.	.	.	*	.	.	10
31	*	*	*	.	.	*	.	.	*	*
32	*	*	*	12
All	226,949	81,125	185,612	552	2,124	19,414	9,093	724	9,430	308,074

The point prevalent cohorts & associated modalities are determined on December 31 of each calendar year without applying the 60-day stable modality rule.

* Values for cells with fewer than 10 patients are suppressed.

." Zero values in this cell.

Table D.9 (continued)

Prevalent ESRD Patients for Selected Years: by Patient Vintage

point prevalent patients on December 31, by year-end treatment modality & number of years patient has had ESRD (vintage)

2000	Dialysis	Functioning graft	Center hemo	Center self hemo	Home hemo	CAPD	CCPD	Other PD	Uncertain dialysis	All
0	80,084	3,287	70,246	39	265	4,063	3,074	150	2,247	83,371
1	56,548	5,506	50,072	35	205	2,817	2,478	120	821	62,054
2	40,705	7,094	36,053	37	146	1,892	1,669	58	850	47,799
3	23,892	8,124	21,565	31	120	866	1,071	37	202	32,016
4	17,597	8,439	15,941	19	118	652	665	15	187	26,036
5	12,935	8,502	11,728	17	92	417	509	17	155	21,437
6	9,517	8,273	8,618	*	46	340	325	10	172	17,790
7	6,819	7,243	6,161	12	50	252	231	11	102	14,062
8	5,302	6,642	4,812	14	37	183	175	*	75	11,944
9	3,961	5,683	3,567	*	39	154	136	*	53	9,644
10	3,125	5,193	2,824	*	31	101	115	*	39	8,318
11	2,466	4,551	2,230	*	22	88	74	*	41	7,017
12	2,143	3,880	1,900	*	27	83	79	*	45	6,023
13	1,622	3,204	1,424	*	18	73	74	*	24	4,826
14	1,364	2,875	1,211	*	16	59	46	*	25	4,239
15	1,160	2,383	1,035	*	15	42	38	.	27	3,543
16	1,023	2,171	891	*	20	38	52	*	16	3,194
17	881	1,914	783	*	14	34	36	*	10	2,795
18	772	1,747	672	*	20	34	27	.	16	2,519
19	633	1,476	548	*	15	22	26	.	20	2,109
20	516	1,219	449	*	17	16	17	*	10	1,735
21	458	1,011	388	.	24	18	19	*	*	1,469
22	382	943	327	*	18	16	10	.	*	1,325
23	441	941	390	*	18	15	10	.	*	1,382
24	219	469	191	*	10	*	*	*	*	688
25	167	295	136	*	*	10	*	.	*	462
26	133	269	104	*	11	*	*	.	*	402
27	117	234	89	.	*	*	*	.	12	351
28	26	80	21	.	*	*	*	.	*	106
29	15	52	11	.	*	*	*	.	.	67
30	*	32	*	*	37
31	*	23	*	25
32	17	33	12	.	*	*	*	.	.	50
33	*	*	*	.	.	*	.	.	.	*
34	*	*	*	*
35	.	*	*
36	*	*	*	*
All	275,053	103,809	244,410	282	1,429	12,307	10,980	458	5,187	378,862

The point prevalent cohorts & associated modalities are determined on December 31 of each calendar year without applying the 60-day stable modality rule.

* Values for cells with fewer than 10 patients are suppressed.

." Zero values in this cell.

Table D.10

Prevalent ESRD Patients for Selected Years: by Number of Years on Modality

point prevalent patients on December 31; by year-end treatment modality

	Functioning graft	Center hemo	Center self hemo	Home hemo	CAPD	CCPD	Other PD	Uncertain dialysis	All	
1992	Dialysis									
0	120,895	9,079	97,180	677	1,371	14,902	2,528	608	3,629	129,974
1	23,580	8,432	18,951	73	173	3,212	299	39	833	32,012
2	6,687	7,406	5,306	14	22	883	52	*	401	14,093
3	3,116	6,234	2,622	13	17	368	21	*	70	9,350
4	1,985	5,455	1,695	13	17	193	*	*	57	7,440
5	1,127	4,760	974	*	*	102	*	*	28	5,887
6	774	4,291	679	*	13	53	.	*	21	5,065
7	462	3,219	375	*	*	61	.	.	13	3,681
8	299	2,627	256	*	*	29	.	*	*	2,926
9	197	2,101	166	*	*	21	*	*	*	2,298
10	136	1,663	123	*	.	*	.	*	*	1,799
11	86	1,276	73	*	.	*	.	*	*	1,362
12	90	1,009	85	.	.	*	.	.	*	1,099
13	88	862	77	.	*	.	.	*	*	950
14	81	723	72	*	*	804
15	69	526	63	.	*	.	.	.	*	595
16	31	166	13	.	*	.	.	.	17	197
17	50	187	*	46	237
18	47	142	*	46	189
19	44	135	44	179
20	*	95	*	*	97
21	.	70	70
22	*	38	*	42
23	.	28	28
24	.	36	36
25	.	*	*
26	.	*	*
27	.	*	*
28	.	*	*
All	159,850	60,584	128,720	814	1,645	19,833	2,909	678	5,251	220,434
1996										
0	151,008	10,996	124,070	432	1,762	12,888	7,179	529	4,148	162,004
1	58,032	9,632	49,484	103	307	5,344	1,760	94	940	67,664
2	13,864	8,506	8,995	12	34	853	132	84	3,754	22,370
3	1,582	7,753	1,199	.	*	145	13	*	216	9,335
4	719	6,714	548	*	*	53	*	*	105	7,433
5	514	6,259	405	*	*	33	*	*	71	6,773
6	303	5,418	230	.	*	26	*	*	42	5,721
7	194	4,597	150	.	*	23	.	*	19	4,791
8	131	3,923	99	.	.	*	*	.	22	4,054
9	111	3,367	91	.	*	*	.	*	11	3,478
10	91	3,029	73	.	*	*	.	*	*	3,120
11	77	2,214	61	*	*	11	.	.	*	2,291
12	55	1,819	46	.	*	*	.	*	*	1,874
13	38	1,562	28	*	.	*	*	*	*	1,600
14	31	1,274	26	.	.	*	.	.	*	1,305
15	18	983	13	*	.	*	.	.	*	1,001
16	18	801	16	.	.	*	.	.	*	819
17	36	664	29	.	*	.	.	.	*	700
18	24	576	21	*	600
19	25	393	23	.	*	.	.	.	*	418
20	14	117	*	10	131
21	21	143	21	164
22	20	100	20	120
23	21	103	21	124
24	*	71	*	72
25	.	49	49
26	*	18	*	19
27	.	20	20
28	.	10	10
29	.	*	*
30	.	11	11
All	226,949	81,125	185,612	552	2,124	19,414	9,093	724	9,430	308,074

The point prevalent cohorts & associated modalities are determined on December 31 of each calendar year without applying the 60-day stable modality rule.

* Values for cells with fewer than 10 patients are suppressed.

." Zero values in this cell.

Table D.10 (continued)

Prevalent ESRD Patients for Selected Years: by Number of Years on Modality

point prevalent patients on December 31; by year-end treatment modality

2000	Dialysis	Functioning graft	Center hemo	Center self hemo	Home hemo	CAPD	CCPD	Other PD	Uncertain dialysis	All
0	139,829	12,958	121,022	128	985	6,822	6,639	381	3,852	152,787
1	82,615	10,756	75,119	90	321	3,314	2,905	52	814	93,371
2	32,101	11,152	29,264	27	69	1,411	997	19	314	43,253
3	10,245	9,647	9,502	17	22	400	280	*	23	19,892
4	4,939	8,486	4,584	16	22	179	112	*	24	13,425
5	4,175	7,354	3,970	*	*	132	40	*	19	11,529
6	735	6,286	632	.	.	25	*	*	71	7,021
7	144	5,680	117	.	.	*	.	.	20	5,824
8	67	4,696	52	.	.	*	.	.	13	4,763
9	58	4,438	50	.	.	*	.	*	*	4,496
10	31	3,856	22	.	.	*	.	.	*	3,887
11	21	3,237	16	.	.	*	.	.	*	3,258
12	15	2,708	10	.	.	*	*	.	*	2,723
13	12	2,321	10	.	.	*	.	.	*	2,333
14	*	2,092	*	.	*	*	.	.	.	2,099
15	11	1,559	10	.	.	*	.	.	.	1,570
16	*	1,327	*	*	1,335
17	*	1,179	*	.	.	*	.	.	.	1,183
18	*	989	*	.	.	*	.	.	.	993
19	*	766	*	*	769
20	.	636	636
21	*	531	*	536
22	*	453	*	*	457
23	*	300	*	303
24	*	71	*	*	74
25	*	84	*	88
26	*	67	*	69
27	*	61	*	69
28	.	51	51
29	.	31	31
30	.	13	13
31	.	11	11
32	.	*	*
33	.	*	*
34	.	*	*
All	275,053	103,809	244,410	282	1,429	12,307	10,980	458	5,187	378,862

The point prevalent cohorts and associated modalities are determined on December 31 of each calendar year without applying the 60-day stable modality rule.

* Values for cells with fewer than 10 patients are suppressed.

." Zero values in this cell.

Table D.11

Incident ESRD Patients for Selected Years: by Diabetic Status & State/Territory

1992 ALL	Center hemo	Center self hemo	Home hemo	CAPD	CCPD	Other PD	Uncertain dialysis	Transplant	All
Alabama	951	*	.	109	*	20	16	37	1,141
Alaska	44	.	.	*	*	*	.	*	55
Arizona	843	.	.	106	*	*	33	16	1,003
Arkansas	394	.	.	85	*	*	29	*	526
California	4,325	*	*	428	14	*	1,147	251	6,178
Colorado	418	*	*	55	*	*	20	15	518
Connecticut	455	.	*	145	21	20	22	20	684
Delaware	162	.	.	31	.	.	*	*	202
District of Columbia	296	.	.	10	.	*	11	*	327
Florida	3,150	*	.	248	45	30	128	65	3,667
Georgia	1,593	.	.	183	12	25	48	40	1,901
Hawaii	262	.	.	*	.	.	10	*	287
Idaho	83	.	.	22	*	.	10	*	126
Illinois	2,282	*	*	238	14	28	100	81	2,748
Indiana	863	.	.	198	*	*	71	34	1,174
Iowa	367	*	.	70	*	*	48	28	526
Kansas	332	.	.	71	*	*	16	13	437
Kentucky	616	*	.	89	*	16	26	26	779
Louisiana	1,171	.	*	97	*	*	34	20	1,327
Maine	129	.	.	14	*	.	*	*	154
Maryland	1,083	.	*	129	*	*	17	39	1,282
Massachusetts	998	.	.	103	*	15	36	32	1,190
Michigan	1,641	*	*	331	13	17	33	67	2,106
Minnesota	610	*	.	34	*	14	54	44	760
Mississippi	644	.	*	50	*	*	30	13	745
Missouri	873	*	.	129	*	*	107	28	1,158
Montana	98	.	.	11	*	*	*	*	125
Nebraska	166	.	*	76	*	*	19	24	295
Nevada	190	*	.	26	*	*	12	10	243
New Hampshire	122	.	.	16	.	*	*	*	152
New Jersey	1,421	*	*	266	22	34	277	23	2,048
New Mexico	257	.	.	38	*	*	26	*	334
New York	3,291	*	.	516	18	75	77	80	4,061
North Carolina	1,286	*	.	229	71	56	139	14	1,798
North Dakota	104	.	.	10	.	.	*	*	124
Ohio	1,873	*	.	254	13	87	105	96	2,429
Oklahoma	478	.	*	74	.	56	26	18	654
Oregon	282	.	.	66	*	17	95	16	479
Pennsylvania	2,419	*	.	223	11	37	148	94	2,936
Rhode Island	186	.	.	10	*	.	*	*	205
South Carolina	776	*	*	99	*	12	124	20	1,046
South Dakota	84	.	*	*	.	.	18	*	113
Tennessee	838	.	*	108	*	*	224	33	1,217
Texas	3,081	*	.	221	26	24	752	76	4,185
Utah	162	.	.	13	.	*	*	20	199
Vermont	51	.	.	*	.	.	*	*	64
Virginia	1,338	.	.	157	*	16	16	42	1,575
Washington	644	18	*	50	*	11	21	23	774
West Virginia	302	.	.	65	*	*	*	*	385
Wisconsin	681	*	.	106	*	11	102	51	959
Wyoming	26	*	.	10	.	*	*	*	42
American Samoa	12	*	.	13
Guam	37	*	.	40
Puerto Rico	641	.	.	68	.	*	*	*	725
Saipan	*	*
Virgin Islands	21	.	.	*	.	.	.	*	24
Foreign	*	.	.	*	.	.	*	40	53
Unknown	153	.	.	*	.	*	564	91	820
All	45,623	63	31	5,722	393	725	4,830	1,740	59,127

The incident cohorts and associated modalities are determined at the time of ESRD initiation without applying the 60-day stable modality rule.

* Values for cells with fewer than 10 patients are suppressed.

." Zero values in this cell.

Table D.11 (continued)

Incident ESRD Patients for Selected Years: by Diabetic Status & State/Territory

1992 DIABETIC	Center hemo	Center self hemo	Home hemo	CAPD	CCPD	Other PD	Uncertain		All
							dialysis	Transplant	
Alabama	317	.	.	46	*	10	*	*	382
Alaska	15	.	.	*	.	.	.	*	17
Arizona	417	.	.	44	*	*	14	*	479
Arkansas	126	.	.	38	.	*	*	*	174
California	1,708	*	*	162	*	*	469	17	2,366
Colorado	171	*	*	25	*	.	*	*	212
Connecticut	154	.	*	55	*	*	*	*	237
Delaware	48	.	.	*	55
District of Columbia	103	.	.	*	.	*	*	.	110
Florida	958	*	.	96	13	*	43	12	1,129
Georgia	536	.	.	74	*	10	17	*	644
Hawaii	137	.	.	*	.	.	*	.	146
Idaho	38	.	.	12	*	.	*	*	60
Illinois	691	.	.	85	*	11	24	15	831
Indiana	315	.	.	69	*	*	16	*	411
Iowa	128	.	.	32	.	*	13	13	187
Kansas	112	.	.	30	*	.	*	.	148
Kentucky	198	.	.	26	*	*	*	*	243
Louisiana	440	.	*	37	*	.	16	*	499
Maine	46	.	.	*	*	.	*	.	52
Maryland	405	.	.	48	*	*	*	*	470
Massachusetts	335	.	.	55	*	*	10	*	407
Michigan	635	.	.	151	*	*	*	20	825
Minnesota	221	*	.	16	.	*	21	*	272
Mississippi	228	.	.	23	*	*	*	*	261
Missouri	318	*	.	48	*	*	35	*	419
Montana	52	.	.	*	*	*	.	*	66
Nebraska	59	.	.	31	*	*	*	13	114
Nevada	59	.	.	12	*	.	*	*	79
New Hampshire	35	.	.	*	.	*	.	*	43
New Jersey	464	.	.	92	*	12	76	*	657
New Mexico	148	.	.	26	*	*	13	.	191
New York	1,089	*	.	201	*	26	21	*	1,349
North Carolina	451	*	.	92	27	18	40	.	629
North Dakota	52	.	.	*	.	.	*	.	60
Ohio	704	.	.	109	*	42	31	12	901
Oklahoma	229	.	*	32	.	31	15	*	310
Oregon	108	.	.	30	*	*	33	.	175
Pennsylvania	879	*	.	98	*	17	37	11	1,048
Rhode Island	62	.	.	*	.	.	.	*	66
South Carolina	269	*	*	36	*	*	50	*	370
South Dakota	46	.	*	*	.	.	*	*	57
Tennessee	314	.	.	34	*	*	62	*	423
Texas	1,289	*	.	92	*	13	319	*	1,732
Utah	70	.	.	*	.	.	*	*	84
Vermont	12	.	.	*	.	.	*	.	17
Virginia	437	.	.	60	*	*	*	*	516
Washington	265	*	.	18	*	*	10	12	318
West Virginia	123	.	.	32	*	*	.	*	158
Wisconsin	270	.	.	47	*	*	38	16	378
Wyoming	*	.	.	*	.	*	.	.	*
American Samoa	*	*	.	*
Guam	24	24
Puerto Rico	357	.	.	33	.	*	*	*	398
Saipan	*	*
Virgin Islands	*	.	.	*	*
Foreign	*	.	.	*	.	.	*	.	*
Unknown	56	.	.	*	.	*	101	*	162
All	16,745	19	10	2,301	148	288	1,625	254	21,390

The incident cohorts and associated modalities are determined at the time of ESRD initiation without applying the 60-day stable modality rule.

* Values for cells with fewer than 10 patients are suppressed.

." Zero values in this cell.

Table D.11 (continued)

Incident ESRD Patients for Selected Years: by Diabetic Status & State/Territory

1992 NON-DIABETIC	Center hemo	Center self hemo	Home hemo	CAPD	CCPD	Other PD	Uncertain dialysis	Transplant	All
Alabama	634	*	.	63	*	10	13	33	759
Alaska	29	.	.	*	*	*	.	*	38
Arizona	426	.	.	62	.	*	19	14	524
Arkansas	268	.	.	47	*	*	24	*	352
California	2,617	*	*	266	11	*	678	234	3,812
Colorado	247	.	.	30	*	*	11	12	306
Connecticut	301	.	.	90	15	11	15	15	447
Delaware	114	.	.	24	.	.	*	*	147
District of Columbia	193	.	.	*	.	.	*	*	217
Florida	2,192	.	.	152	32	24	85	53	2,538
Georgia	1,057	.	.	109	*	15	31	37	1,257
Hawaii	125	.	.	*	.	.	*	*	141
Idaho	45	.	.	10	*	.	*	*	66
Illinois	1,591	*	*	153	*	17	76	66	1,917
Indiana	548	.	.	129	*	*	55	28	763
Iowa	239	*	.	38	*	*	35	15	339
Kansas	220	.	.	41	.	*	12	13	289
Kentucky	418	*	.	63	*	10	21	22	536
Louisiana	731	.	.	60	*	*	18	16	828
Maine	83	.	.	11	.	.	*	*	102
Maryland	678	.	*	81	*	*	11	32	812
Massachusetts	663	.	.	48	*	12	26	29	783
Michigan	1,006	*	*	180	*	13	26	47	1,281
Minnesota	389	*	.	18	*	*	33	37	488
Mississippi	416	.	*	27	*	*	23	12	484
Missouri	555	*	.	81	*	*	72	21	739
Montana	46	.	.	*	.	*	*	*	59
Nebraska	107	.	*	45	*	*	12	11	181
Nevada	131	*	.	14	*	*	*	*	164
New Hampshire	87	.	.	12	.	*	*	*	109
New Jersey	957	*	*	174	13	22	201	19	1,391
New Mexico	109	.	.	12	.	.	13	*	143
New York	2,202	*	.	315	12	49	56	75	2,712
North Carolina	835	*	.	137	44	38	99	14	1,169
North Dakota	52	.	.	*	.	.	*	*	64
Ohio	1,169	*	.	145	10	45	74	84	1,528
Oklahoma	249	.	*	42	.	25	11	16	344
Oregon	174	.	.	36	*	15	62	16	304
Pennsylvania	1,540	*	.	125	*	20	111	83	1,888
Rhode Island	124	.	.	*	*	.	*	*	139
South Carolina	507	.	*	63	*	*	74	16	676
South Dakota	38	.	*	*	.	.	12	*	56
Tennessee	524	.	*	74	*	*	162	28	794
Texas	1,792	*	.	129	17	11	433	67	2,453
Utah	92	.	.	*	.	*	*	13	115
Vermont	39	.	.	*	.	.	*	*	47
Virginia	901	.	.	97	*	*	12	37	1,059
Washington	379	12	*	32	*	*	11	11	456
West Virginia	179	.	.	33	*	*	*	*	227
Wisconsin	411	*	.	59	*	*	64	35	581
Wyoming	20	*	.	*	.	*	*	*	33
American Samoa	*	*
Guam	13	*	.	16
Puerto Rico	284	.	.	35	.	*	*	*	327
Saipan	*	*
Virgin Islands	16	.	.	*	.	.	.	*	18
Foreign	*	.	.	*	.	.	*	40	48
Unknown	97	.	.	*	.	*	463	90	658
All	28,878	44	21	3,421	245	437	3,205	1,486	37,737

The incident cohorts and associated modalities are determined at the time of ESRD initiation without applying the 60-day stable modality rule.

* Values for cells with fewer than 10 patients are suppressed.

." Zero values in this cell.

Table D.11 (continued)

Incident ESRD Patients for Selected Years: by Diabetic Status & State/Territory

1996	Center	Center	Home				Uncertain		
ALL	hemo	self hemo	hemo	CAPD	CCPD	Other PD	dialysis	Transplant	All
Alabama	1,258	.	*	133	35	*	19	27	1,485
Alaska	56	.	.	*	.	.	.	*	65
Arizona	1,072	.	11	111	18	.	64	18	1,294
Arkansas	586	*	*	80	20	*	24	10	723
California	6,589	*	*	680	136	23	991	152	8,579
Colorado	554	*	*	65	16	*	14	18	673
Connecticut	651	.	*	153	24	*	52	12	896
Delaware	191	.	*	17	*	.	*	*	224
District of Columbia	373	.	*	*	*	.	*	*	398
Florida	3,535	.	590	254	137	36	280	70	4,902
Georgia	1,753	.	115	216	70	13	284	26	2,477
Hawaii	329	.	*	33	*	.	10	*	382
Idaho	151	.	.	30	*	.	19	10	212
Illinois	3,055	.	21	264	81	17	132	86	3,656
Indiana	1,157	*	*	207	11	*	98	32	1,512
Iowa	443	*	*	69	18	.	48	26	606
Kansas	435	*	.	91	18	.	*	14	568
Kentucky	754	.	*	73	39	13	67	25	972
Louisiana	1,490	.	*	66	24	.	21	17	1,622
Maine	191	.	*	14	*	.	*	*	216
Maryland	1,579	.	16	118	48	*	80	38	1,882
Massachusetts	1,283	.	*	90	40	*	36	34	1,491
Michigan	2,387	.	*	354	24	*	68	64	2,914
Minnesota	756	*	*	36	28	*	34	64	927
Mississippi	878	.	*	87	18	*	19	*	1,021
Missouri	1,256	.	*	156	64	*	*	19	1,513
Montana	120	*	.	16	*	*	*	*	148
Nebraska	231	.	.	79	11	*	17	14	354
Nevada	297	.	*	34	10	.	24	*	376
New Hampshire	129	.	.	20	*	.	*	*	162
New Jersey	1,967	*	*	186	175	14	250	42	2,638
New Mexico	356	.	.	64	22	*	25	13	483
New York	4,760	15	20	394	126	60	96	75	5,546
North Carolina	1,595	.	68	171	147	52	354	20	2,407
North Dakota	109	.	.	*	*	.	*	12	138
Ohio	2,572	.	*	330	129	73	160	87	3,360
Oklahoma	668	*	*	57	22	28	27	14	820
Oregon	430	.	*	80	*	*	71	13	603
Pennsylvania	3,342	.	*	181	93	28	187	82	3,918
Rhode Island	245	.	.	10	*	.	12	*	275
South Carolina	944	.	59	73	46	*	223	27	1,374
South Dakota	133	.	.	12	16	.	31	*	194
Tennessee	1,326	*	*	89	58	11	39	29	1,554
Texas	4,458	*	*	246	134	14	575	73	5,516
Utah	218	.	.	21	*	*	*	16	268
Vermont	80	.	*	14	*	.	10	*	109
Virginia	1,765	.	12	120	52	30	74	51	2,104
Washington	811	24	*	79	18	*	20	43	1,004
West Virginia	444	.	*	32	28	*	20	11	539
Wisconsin	839	.	*	94	50	*	116	52	1,164
Wyoming	52	.	.	*	*	.	*	*	68
American Samoa	11	11
Guam	48	.	.	*	.	*	*	*	63
Puerto Rico	762	.	*	96	54	16	20	*	954
Saipan	13	13
Virgin Islands	31	.	.	*	.	.	*	.	34
Foreign	*	*	*	*	.	.	*	30	44
Unknown	*	44	32	80
All	61,530	64	1,010	5,932	2,103	512	4,824	1,556	77,531

The incident cohorts and associated modalities are determined at the time of ESRD initiation without applying the 60-day stable modality rule.

* Values for cells with fewer than 10 patients are suppressed.

." Zero values in this cell.

Table D.11 (continued)

Incident ESRD Patients for Selected Years: by Diabetic Status & State/Territory

1996 DIABETIC	Center hemo	Center self hemo	Home hemo	CAPD	CCPD	Other PD	Uncertain dialysis	Transplant	All
Alabama	497	.	*	55	11	*	*	*	583
Alaska	27	.	.	*	.	.	.	*	31
Arizona	538	.	*	51	*	.	27	*	633
Arkansas	238	.	*	44	*	.	11	*	305
California	2,912	*	.	307	52	*	418	27	3,726
Colorado	277	*	*	31	*	*	*	*	328
Connecticut	242	.	.	60	10	*	23	*	340
Delaware	68	.	*	*	*	.	*	*	81
District of Columbia	157	.	.	*	*	.	.	.	161
Florida	1,304	.	218	110	59	*	74	*	1,783
Georgia	740	.	49	99	31	*	134	*	1,064
Hawaii	207	.	.	18	*	.	*	.	232
Idaho	63	.	.	13	.	.	*	*	87
Illinois	1,187	.	13	122	32	*	44	19	1,425
Indiana	495	*	*	89	*	*	40	*	636
Iowa	189	*	*	31	*	.	23	*	256
Kansas	183	*	.	34	*	.	*	*	231
Kentucky	320	.	*	32	18	*	32	*	411
Louisiana	656	.	*	39	*	.	*	*	711
Maine	70	.	.	*	*	.	*	.	76
Maryland	613	.	*	57	18	*	10	11	719
Massachusetts	501	.	*	36	22	*	16	*	585
Michigan	1,059	.	*	165	14	*	29	18	1,292
Minnesota	300	*	*	12	*	*	*	17	351
Mississippi	395	.	*	36	*	*	.	.	443
Missouri	539	.	*	72	33	*	*	*	649
Montana	55	.	.	*	*	*	*	*	65
Nebraska	90	.	.	45	*	*	*	*	155
Nevada	98	.	*	10	*	.	12	*	131
New Hampshire	48	.	.	*	*	.	*	.	57
New Jersey	786	*	*	89	81	*	112	*	1,079
New Mexico	217	.	.	36	13	*	16	*	286
New York	1,944	*	*	147	58	18	24	13	2,213
North Carolina	674	.	34	70	54	22	162	*	1,022
North Dakota	44	*	*	50
Ohio	1,113	.	*	181	59	27	53	16	1,451
Oklahoma	321	.	*	24	14	11	*	*	382
Oregon	171	.	.	32	*	*	23	*	233
Pennsylvania	1,339	.	*	70	40	13	61	*	1,534
Rhode Island	94	.	.	*	*	.	*	*	107
South Carolina	393	.	24	33	21	*	105	*	587
South Dakota	70	.	.	*	*	.	*	.	91
Tennessee	523	.	.	36	28	*	16	*	615
Texas	2,251	*	*	124	63	11	257	*	2,721
Utah	110	.	.	12	*	.	*	*	129
Vermont	32	.	*	*	*	.	*	*	42
Virginia	694	.	*	55	16	16	13	*	805
Washington	327	11	.	43	*	*	*	15	408
West Virginia	195	.	.	19	14	*	*	*	236
Wisconsin	347	.	*	41	24	*	41	15	471
Wyoming	25	.	.	*	.	.	.	*	30
American Samoa	*	*
Guam	34	.	.	*	.	.	*	.	45
Puerto Rico	443	.	*	50	26	11	*	*	536
Saipan	*	*
Virgin Islands	18	.	.	*	19
Foreign	*	*	.	*	.	.	.	*	*
Unknown	*	*	*
All	26,250	27	398	2,684	900	214	1,872	316	32,661

The incident cohorts and associated modalities are determined at the time of ESRD initiation without applying the 60-day stable modality rule.

* Values for cells with fewer than 10 patients are suppressed.

." Zero values in this cell.

Table D.11 (continued)

Incident ESRD Patients for Selected Years: by Diabetic Status & State/Territory

1996 NON-DIABETIC	Center hemo	Center self hemo	Home hemo	CAPD	CCPD	Other PD	Uncertain dialysis	Transplant	All
Alabama	761	.	*	78	24	*	15	21	902
Alaska	29	.	.	*	.	.	.	*	34
Arizona	534	.	*	60	15	.	37	10	661
Arkansas	348	*	*	36	12	*	13	*	418
California	3,677	*	*	373	84	14	573	125	4,853
Colorado	277	*	.	34	10	*	*	12	345
Connecticut	409	.	*	93	14	*	29	*	556
Delaware	123	.	*	*	.	.	*	*	143
District of Columbia	216	.	*	*	*	.	*	*	237
Florida	2,231	.	372	144	78	27	206	61	3,119
Georgia	1,013	.	66	117	39	*	150	21	1,413
Hawaii	122	.	*	15	*	.	*	*	150
Idaho	88	.	.	17	*	.	10	*	125
Illinois	1,868	.	*	142	49	*	88	67	2,231
Indiana	662	.	*	118	*	*	58	28	876
Iowa	254	.	.	38	12	.	25	21	350
Kansas	252	.	.	57	10	.	*	12	337
Kentucky	434	.	.	41	21	*	35	21	561
Louisiana	834	.	*	27	18	.	19	12	911
Maine	121	.	*	11	*	.	*	*	140
Maryland	966	.	*	61	30	*	70	27	1,163
Massachusetts	782	.	*	54	18	*	20	27	906
Michigan	1,328	.	*	189	10	*	39	46	1,622
Minnesota	456	.	*	24	20	*	26	47	576
Mississippi	483	.	*	51	15	*	17	*	578
Missouri	717	.	*	84	31	*	84	16	864
Montana	65	*	.	12	*	.	*	*	83
Nebraska	141	.	.	34	*	.	*	*	199
Nevada	199	.	.	24	*	.	12	*	245
New Hampshire	81	.	.	14	*	.	*	*	105
New Jersey	1,181	.	*	97	94	12	138	35	1,559
New Mexico	139	.	.	28	*	*	*	11	197
New York	2,816	11	15	247	68	42	72	62	3,333
North Carolina	921	.	34	101	93	30	192	14	1,385
North Dakota	65	.	.	*	*	.	*	*	88
Ohio	1,459	.	*	149	70	46	107	71	1,909
Oklahoma	347	*	*	33	*	17	20	11	438
Oregon	259	.	*	48	*	.	48	*	370
Pennsylvania	2,003	.	*	111	53	15	126	74	2,384
Rhode Island	151	.	.	*	*	.	*	*	168
South Carolina	551	.	35	40	25	.	118	18	787
South Dakota	63	.	.	*	*	.	23	*	103
Tennessee	803	*	*	53	30	*	23	21	939
Texas	2,207	*	*	122	71	*	318	64	2,795
Utah	108	.	.	*	.	*	*	12	139
Vermont	48	.	.	10	.	.	*	*	67
Virginia	1,071	.	*	65	36	14	61	44	1,299
Washington	484	13	*	36	14	*	16	28	596
West Virginia	249	.	*	13	14	*	15	*	303
Wisconsin	492	.	*	53	26	*	75	37	693
Wyoming	27	.	.	*	*	.	*	*	38
American Samoa	*	*
Guam	14	.	.	*	.	*	*	*	18
Puerto Rico	319	.	*	46	28	*	16	*	418
Saipan	*	*
Virgin Islands	13	*	.	15
Foreign	*	.	*	*	.	.	*	29	39
Unknown	*	43	31	78
All	35,280	37	612	3,248	1,203	298	2,952	1,240	44,870

The incident cohorts and associated modalities are determined at the time of ESRD initiation without applying the 60-day stable modality rule.

* Values for cells with fewer than 10 patients are suppressed.

." Zero values in this cell.

Table D.11 (continued)

Incident ESRD Patients for Selected Years: by Diabetic Status & State/Territory

2000 ALL	Center hemo	Center self hemo	Home hemo	CAPD	CCPD	Other PD	Uncertain dialysis	Transplant	All
Alabama	1,555	.	.	102	49	*	68	34	1,809
Alaska	66	.	.	19	.	.	*	*	92
Arizona	1,513	.	19	93	15	*	35	37	1,714
Arkansas	740	.	.	75	26	*	34	14	890
California	10,005	*	*	571	156	13	297	236	11,288
Colorado	711	.	*	38	*	.	28	29	815
Connecticut	753	.	*	138	48	*	62	43	1,048
Delaware	262	.	.	*	*	.	*	10	284
District of Columbia	440	.	*	15	.	*	*	*	469
Florida	5,093	.	64	202	133	14	204	85	5,795
Georgia	2,864	.	13	154	70	*	88	47	3,242
Hawaii	454	.	.	32	*	.	29	*	524
Idaho	199	.	.	35	*	.	*	*	255
Illinois	3,731	.	108	212	130	*	109	106	4,401
Indiana	1,640	.	*	148	34	10	93	44	1,971
Iowa	588	.	*	54	19	.	27	33	725
Kansas	572	*	*	81	19	.	23	12	712
Kentucky	1,039	.	.	70	20	*	65	22	1,218
Louisiana	1,911	.	*	86	29	*	40	26	2,099
Maine	272	.	.	12	13	.	*	*	310
Maryland	1,933	.	*	91	49	*	60	92	2,238
Massachusetts	1,477	.	*	65	59	*	72	39	1,723
Michigan	2,914	.	*	254	43	10	80	88	3,392
Minnesota	988	.	.	31	16	*	25	65	1,126
Mississippi	983	.	*	82	51	*	26	*	1,156
Missouri	1,694	.	11	95	54	.	35	38	1,927
Montana	132	*	*	13	*	.	10	*	167
Nebraska	389	.	*	87	*	.	12	15	508
Nevada	483	.	*	29	14	*	21	16	566
New Hampshire	219	.	.	31	11	*	14	11	287
New Jersey	2,701	*	*	96	95	12	101	69	3,079
New Mexico	503	.	.	28	29	.	27	10	597
New York	6,009	*	15	198	102	58	133	114	6,634
North Carolina	2,740	.	*	129	110	44	98	39	3,164
North Dakota	136	.	.	12	*	.	*	*	168
Ohio	3,643	*	*	299	115	10	135	134	4,344
Oklahoma	939	.	*	67	48	11	40	14	1,120
Oregon	647	.	.	66	18	.	18	16	765
Pennsylvania	4,004	*	*	141	103	*	143	102	4,502
Rhode Island	283	.	.	15	*	.	23	*	330
South Carolina	1,402	.	*	57	80	*	57	38	1,639
South Dakota	208	.	.	12	*	.	12	*	242
Tennessee	1,690	.	.	60	80	*	70	34	1,941
Texas	6,263	.	12	242	227	10	222	126	7,102
Utah	271	.	.	28	*	.	17	17	342
Vermont	99	.	.	*	*	.	18	*	126
Virginia	2,058	*	*	120	33	*	82	59	2,363
Washington	1,170	12	*	63	26	*	30	32	1,338
West Virginia	590	.	.	40	60	*	24	16	733
Wisconsin	1,174	.	*	86	35	*	49	66	1,415
Wyoming	73	.	.	*	*	.	*	*	83
American Samoa	21	*	.	22
Guam	67	.	.	*	*	.	.	*	73
Puerto Rico	917	.	*	34	101	*	38	12	1,106
Saipan	19	*	.	20
Virgin Islands	26	.	.	*	*	.	*	.	30
Foreign	13	.	.	*	.	.	*	19	37
Unknown	*	77	41	126
All	83,294	29	312	4,724	2,377	271	3,015	2,170	96,192

The incident cohorts and associated modalities are determined at the time of ESRD initiation without applying the 60-day stable modality rule.

* Values for cells with fewer than 10 patients are suppressed.

." Zero values in this cell.

Table D.11 (continued)

Incident ESRD Patients for Selected Years: by Diabetic Status & State/Territory

2000 DIABETIC	Center hemo	Center self hemo	Home hemo	CAPD	CCPD	Other PD	Uncertain dialysis	Transplant	All
Alabama	698	.	.	44	22	.	11	*	779
Alaska	31	.	.	*	.	.	*	.	39
Arizona	830	.	*	44	*	*	*	*	902
Arkansas	306	.	.	40	*	.	*	.	359
California	4,857	*	*	256	43	*	25	18	5,211
Colorado	364	.	.	22	*	.	.	*	391
Connecticut	318	.	.	64	22	*	24	*	437
Delaware	123	.	.	*	*	.	.	*	127
District of Columbia	158	.	*	*	.	.	*	*	166
Florida	2,101	.	27	75	44	*	39	14	2,304
Georgia	1,225	.	*	67	29	*	*	*	1,338
Hawaii	278	.	.	18	.	.	15	*	312
Idaho	88	.	.	16	*	.	*	*	117
Illinois	1,507	.	53	87	55	.	18	14	1,734
Indiana	678	.	*	64	18	*	21	*	794
Iowa	246	.	.	33	*	.	*	*	297
Kansas	271	.	*	47	10	.	*	*	335
Kentucky	462	.	.	36	*	.	23	*	529
Louisiana	828	.	.	36	*	*	.	*	877
Maine	118	.	.	*	*	.	*	*	132
Maryland	738	.	*	33	20	*	*	*	812
Massachusetts	555	.	*	28	25	*	25	*	639
Michigan	1,315	.	*	115	17	*	13	18	1,486
Minnesota	374	.	.	13	*	*	*	16	411
Mississippi	425	.	*	35	20	*	*	*	486
Missouri	773	.	*	37	17	.	*	*	838
Montana	64	*	.	*	*	.	*	*	78
Nebraska	158	.	.	48	*	.	*	*	215
Nevada	225	.	*	16	*	.	.	*	252
New Hampshire	81	.	.	15	*	.	*	.	103
New Jersey	1,126	.	*	37	44	*	*	12	1,225
New Mexico	321	.	.	15	17	.	10	.	363
New York	2,475	*	*	87	37	29	18	*	2,659
North Carolina	1,221	.	*	46	38	28	15	*	1,359
North Dakota	54	.	.	*	.	.	*	*	64
Ohio	1,727	*	*	156	55	*	39	17	1,999
Oklahoma	465	.	.	44	22	*	*	*	549
Oregon	262	.	.	29	*	.	*	*	306
Pennsylvania	1,726	.	*	52	42	*	48	10	1,882
Rhode Island	87	.	.	*	.	.	*	*	102
South Carolina	639	.	*	23	35	*	*	*	707
South Dakota	95	.	.	*	*	.	*	*	104
Tennessee	748	.	.	24	35	*	13	*	829
Texas	3,365	.	*	125	99	*	37	14	3,649
Utah	132	.	.	18	*	.	*	*	163
Vermont	34	.	.	.	*	.	10	*	46
Virginia	900	.	.	59	15	*	12	*	997
Washington	515	*	.	33	*	*	*	*	569
West Virginia	320	.	.	18	31	*	*	*	381
Wisconsin	480	.	.	35	10	*	11	14	551
Wyoming	39	.	.	*	.	.	*	.	42
American Samoa	18	18
Guam	39	.	.	*	40
Puerto Rico	559	.	*	23	49	*	.	.	634
Saipan	14	14
Virgin Islands	15	.	.	*	16
Foreign	*	*
Unknown	*	.	*
All	37,574	*	131	2,159	956	127	538	278	41,772

The incident cohorts and associated modalities are determined at the time of ESRD initiation without applying the 60-day stable modality rule.

* Values for cells with fewer than 10 patients are suppressed.

." Zero values in this cell.

Table D.11 (continued)

Incident ESRD Patients for Selected Years: by Diabetic Status & State/Territory

2000 NON-DIABETIC	Center hemo	Center self hemo	Home hemo	CAPD	CCPD	Other PD	Uncertain dialysis	Transplant	All
Alabama	857	.	.	58	27	*	57	30	1,030
Alaska	35	.	.	12	.	.	*	*	53
Arizona	683	.	10	49	*	.	32	30	812
Arkansas	434	.	.	35	19	*	28	14	531
California	5,148	*	*	315	113	*	272	218	6,077
Colorado	347	.	*	16	*	.	28	26	424
Connecticut	435	.	*	74	26	*	38	35	611
Delaware	139	.	.	*	*	*	*	*	157
District of Columbia	282	.	.	11	.	*	*	*	303
Florida	2,992	.	37	127	89	10	165	71	3,491
Georgia	1,639	.	10	87	41	*	81	45	1,904
Hawaii	176	.	.	14	*	.	14	*	212
Idaho	111	.	.	19	*	.	*	*	138
Illinois	2,224	.	55	125	75	*	91	92	2,667
Indiana	962	.	.	84	16	*	72	38	1,177
Iowa	342	.	*	21	13	.	20	28	428
Kansas	301	*	*	34	*	.	18	11	377
Kentucky	577	.	.	34	14	*	42	20	689
Louisiana	1,083	.	*	50	20	*	40	23	1,222
Maine	154	.	.	*	11	.	*	*	178
Maryland	1,195	.	*	58	29	*	52	83	1,426
Massachusetts	922	.	*	37	34	*	47	37	1,084
Michigan	1,599	.	*	139	26	*	67	70	1,906
Minnesota	614	.	.	18	11	.	23	49	715
Mississippi	558	.	*	47	31	*	24	*	670
Missouri	921	.	*	58	37	.	32	32	1,089
Montana	68	*	*	*	.	.	*	*	89
Nebraska	231	.	*	39	*	.	*	11	293
Nevada	258	.	*	13	*	*	21	15	314
New Hampshire	138	.	.	16	*	*	*	11	184
New Jersey	1,575	*	*	59	51	10	98	57	1,854
New Mexico	182	.	.	13	12	.	17	10	234
New York	3,534	*	*	111	65	29	115	108	3,975
North Carolina	1,519	.	*	83	72	16	83	30	1,805
North Dakota	82	.	.	*	*	.	*	*	104
Ohio	1,916	*	*	143	60	*	96	117	2,345
Oklahoma	474	.	*	23	26	*	33	*	571
Oregon	385	.	.	37	*	.	13	15	459
Pennsylvania	2,278	*	*	89	61	*	95	92	2,620
Rhode Island	196	.	.	*	*	.	16	*	228
South Carolina	763	.	.	34	45	*	53	35	932
South Dakota	113	.	.	*	*	.	10	*	138
Tennessee	942	.	.	36	45	*	57	31	1,112
Texas	2,898	.	*	117	128	*	185	112	3,453
Utah	139	.	.	10	*	.	14	*	179
Vermont	65	.	.	*	*	.	*	*	80
Virginia	1,158	*	*	61	18	*	70	50	1,366
Washington	655	*	*	30	20	*	26	29	769
West Virginia	270	.	.	22	29	*	16	14	352
Wisconsin	694	.	*	51	25	*	38	52	864
Wyoming	34	.	.	*	*	.	*	*	41
American Samoa	*	*	.	*
Guam	28	.	.	*	*	.	.	*	33
Puerto Rico	358	.	.	11	52	*	38	12	472
Saipan	*	*	.	*
Virgin Islands	11	.	.	.	*	.	*	.	14
Foreign	10	.	.	*	.	.	*	19	34
Unknown	*	76	41	125
All	45,720	20	181	2,565	1,421	144	2,477	1,892	54,420

The incident cohorts and associated modalities are determined at the time of ESRD initiation without applying the 60-day stable modality rule.

* Values for cells with fewer than 10 patients are suppressed.

." Zero values in this cell.

Table D.12

Prevalent ESRD Patients for Selected Years: by Diabetic Status & State/Territory

point prevalent patients on December 31

1992 ALL	Center hemo	Center self hemo	Home hemo	CAPD	CCPD	Other PD	Uncertain dialysis	Transplant	All
Alabama	2,773	*	*	345	85	*	36	1,123	4,377
Alaska	87	*	*	23	*	*	*	92	210
Arizona	2,101	21	15	272	86	*	42	813	3,352
Arkansas	1,075	*	*	219	32	*	27	504	1,870
California	13,885	74	37	1,659	269	*	1,118	6,470	23,521
Colorado	1,103	*	15	181	46	*	39	707	2,095
Connecticut	1,331	*	33	408	80	14	32	720	2,620
Delaware	459	*	*	86	*	*	*	158	714
District of Columbia	939	*	*	31	*	*	*	179	1,164
Florida	8,245	20	25	851	194	33	147	2,978	12,493
Georgia	4,797	*	15	508	144	27	67	1,465	7,027
Hawaii	794	15	*	49	*	*	13	236	1,110
Idaho	245	.	10	79	*	*	13	233	590
Illinois	6,013	57	40	843	96	27	142	3,264	10,482
Indiana	2,336	*	24	720	33	*	50	1,233	4,407
Iowa	914	*	66	227	35	*	31	779	2,056
Kansas	921	.	25	267	31	*	*	535	1,789
Kentucky	1,460	*	12	340	17	21	27	928	2,807
Louisiana	3,227	*	26	321	74	*	38	808	4,499
Maine	308	.	*	49	16	.	*	304	687
Maryland	3,011	43	23	435	34	12	40	1,130	4,728
Massachusetts	2,573	15	39	322	46	29	65	1,350	4,439
Michigan	4,121	*	55	1,149	66	22	57	2,339	7,812
Minnesota	1,611	*	26	150	13	*	44	1,437	3,288
Mississippi	2,044	*	120	165	*	*	26	439	2,804
Missouri	2,420	*	27	467	91	*	42	1,310	4,368
Montana	244	.	11	56	*	*	*	200	520
Nebraska	418	*	*	246	26	*	*	424	1,136
Nevada	439	*	17	76	21	*	12	329	897
New Hampshire	297	.	*	57	26	*	*	233	630
New Jersey	4,225	35	104	795	142	51	263	1,409	7,024
New Mexico	769	*	*	186	15	*	21	368	1,366
New York	9,339	224	140	1,662	154	43	161	3,206	14,929
North Carolina	4,110	17	44	797	258	65	106	1,297	6,694
North Dakota	205	.	*	28	*	.	*	216	463
Ohio	4,872	*	16	918	82	62	140	2,965	9,058
Oklahoma	1,241	*	16	290	*	19	17	673	2,265
Oregon	860	*	37	290	35	*	80	658	1,964
Pennsylvania	6,465	67	48	818	87	57	192	3,069	10,803
Rhode Island	504	.	.	33	10	.	*	202	756
South Carolina	2,599	61	22	295	59	*	110	799	3,953
South Dakota	206	.	49	19	*	.	*	189	475
Tennessee	2,809	*	57	383	48	*	71	1,232	4,609
Texas	9,871	56	63	946	258	42	449	3,744	15,429
Utah	389	*	*	49	.	*	10	506	963
Vermont	129	.	12	40	*	.	*	118	302
Virginia	3,726	*	66	553	56	17	43	1,374	5,838
Washington	1,606	17	192	211	28	31	34	1,148	3,267
West Virginia	646	*	*	171	25	*	*	467	1,333
Wisconsin	1,619	25	30	409	35	*	82	1,734	3,937
Wyoming	63	*	12	39	*	.	*	102	225
American Samoa	30	*	.	31
Guam	75	*	*	14	103
Puerto Rico	1,786	.	16	265	13	*	18	386	2,486
Saipan	15	*	*	21
Virgin Islands	58	.	.	*	.	.	.	*	74
Foreign	43	.	*	*	*	.	10	150	215
Unknown	269	.	.	17	*	*	1,236	1,829	3,359
All	128,720	814	1,645	19,833	2,909	678	5,251	60,584	220,434

The point prevalent cohorts and associated modalities are determined on December 31 of each calendar year without applying the 60-day stable modality rule.

* Values for cells with fewer than 10 patients are suppressed.

." Zero values in this cell.

Table D.12 (continued)

Prevalent ESRD Patients for Selected Years: by Diabetic Status & State/Territory

point prevalent patients on December 31

1992 DIABETIC	Center hemo	Center self hemo	Home hemo	CAPD	CCPD	Other PD	Uncertain dialysis	Transplant	All
Alabama	701	*	*	92	16	*	*	112	928
Alaska	21	*	.	*	.	*	.	14	45
Arizona	952	*	*	91	23	*	10	165	1,255
Arkansas	288	.	.	77	*	.	*	75	448
California	4,761	20	*	513	62	*	406	987	6,760
Colorado	402	.	*	74	12	.	12	184	689
Connecticut	389	*	*	115	27	*	*	124	669
Delaware	139	*	.	17	.	.	.	23	180
District of Columbia	264	.	.	*	.	*	*	18	293
Florida	2,203	*	*	232	49	13	31	527	3,063
Georgia	1,343	*	*	140	37	*	16	239	1,785
Hawaii	364	*	.	23	.	.	*	19	415
Idaho	75	.	*	29	*	.	*	64	176
Illinois	1,536	*	*	242	24	*	25	526	2,370
Indiana	700	.	*	219	*	*	10	200	1,146
Iowa	249	.	10	71	13	*	*	174	525
Kansas	266	.	*	109	*	*	*	101	488
Kentucky	439	.	*	95	*	*	*	125	682
Louisiana	982	.	*	91	24	*	12	124	1,241
Maine	90	.	.	13	*	.	*	43	151
Maryland	911	*	*	132	10	*	12	173	1,250
Massachusetts	683	*	.	99	11	*	10	234	1,046
Michigan	1,283	.	17	390	14	10	11	473	2,198
Minnesota	517	*	12	48	*	*	14	417	1,018
Mississippi	549	.	22	54	*	.	*	51	684
Missouri	711	.	*	145	21	*	11	231	1,127
Montana	94	.	*	25	*	*	.	58	186
Nebraska	128	*	*	78	*	.	*	137	359
Nevada	124	.	*	24	*	.	*	68	230
New Hampshire	71	.	*	14	*	*	.	54	148
New Jersey	1,168	*	19	224	51	19	76	208	1,767
New Mexico	383	.	.	91	*	*	*	63	552
New York	2,611	32	12	473	48	12	26	409	3,623
North Carolina	1,206	*	*	232	83	13	28	207	1,781
North Dakota	90	.	.	13	.	.	*	63	172
Ohio	1,494	*	*	298	27	25	35	607	2,491
Oklahoma	495	*	*	117	*	11	*	126	762
Oregon	248	.	*	98	*	*	21	140	525
Pennsylvania	1,920	23	*	267	33	18	35	538	2,840
Rhode Island	129	.	.	*	*	.	*	32	169
South Carolina	794	15	*	78	10	*	33	112	1,049
South Dakota	84	.	32	*	.	.	*	55	179
Tennessee	735	*	*	91	16	.	22	196	1,064
Texas	3,709	13	*	324	73	14	154	686	4,979
Utah	148	.	*	15	.	.	*	168	336
Vermont	39	.	*	15	.	.	.	25	80
Virginia	1,043	.	13	154	16	10	*	201	1,446
Washington	517	*	39	66	11	*	*	283	938
West Virginia	216	.	.	55	*	*	*	98	379
Wisconsin	517	*	*	159	*	*	27	480	1,196
Wyoming	18	.	*	10	.	.	*	24	58
American Samoa	12	*	.	13
Guam	35	*	*	*	43
Puerto Rico	755	.	*	92	*	*	*	32	890
Saipan	*	*	.	*
Virgin Islands	*	.	.	*	*
Foreign	*	.	.	*	.	.	*	*	15
Unknown	99	.	.	*	*	*	198	17	321
All	39,719	160	320	6,162	801	221	1,344	10,512	59,239

The point prevalent cohorts and associated modalities are determined on December 31 of each calendar year without applying the 60-day stable modality rule.

* Values for cells with fewer than 10 patients are suppressed.

." Zero values in this cell.

Table D.12 (continued)

Prevalent ESRD Patients for Selected Years: by Diabetic Status & State/Territory

point prevalent patients on December 31

1992 NON-DIABETIC	Center hemo	Center self hemo	Home hemo	CAPD	CCPD	Other PD	Uncertain dialysis	Transplant	All
Alabama	2,072	.	*	253	69	*	33	1,011	3,449
Alaska	66	.	*	15	*	*	*	78	165
Arizona	1,149	14	*	181	63	*	32	648	2,097
Arkansas	787	*	*	142	29	*	22	429	1,422
California	9,124	54	28	1,146	207	*	712	5,483	16,761
Colorado	701	*	10	107	34	*	27	523	1,406
Connecticut	942	*	29	293	53	11	26	596	1,951
Delaware	320	*	*	69	*	*	*	135	534
District of Columbia	675	*	*	24	*	*	*	161	871
Florida	6,042	15	22	619	145	20	116	2,451	9,430
Georgia	3,454	*	14	368	107	20	51	1,226	5,242
Hawaii	430	14	*	26	*	.	*	217	695
Idaho	170	.	*	50	*	*	*	169	414
Illinois	4,477	51	34	601	72	22	117	2,738	8,112
Indiana	1,636	*	20	501	24	*	40	1,033	3,261
Iowa	665	*	56	156	22	*	24	605	1,531
Kansas	655	.	21	158	26	.	*	434	1,301
Kentucky	1,021	*	*	245	10	12	24	803	2,125
Louisiana	2,245	*	20	230	50	*	26	684	3,258
Maine	218	.	*	36	12	.	*	261	536
Maryland	2,100	38	18	303	24	10	28	957	3,478
Massachusetts	1,890	14	39	223	35	21	55	1,116	3,393
Michigan	2,838	*	38	759	52	12	46	1,866	5,614
Minnesota	1,094	*	14	102	*	.	30	1,020	2,270
Mississippi	1,495	*	98	111	*	*	21	388	2,120
Missouri	1,709	*	22	322	70	*	31	1,079	3,241
Montana	150	.	*	31	*	*	*	142	334
Nebraska	290	*	*	168	18	*	*	287	777
Nevada	315	*	*	52	17	*	10	261	667
New Hampshire	226	.	*	43	19	*	*	179	482
New Jersey	3,057	33	85	571	91	32	187	1,201	5,257
New Mexico	386	*	*	95	10	*	13	305	814
New York	6,728	192	128	1,189	106	31	135	2,797	11,306
North Carolina	2,904	13	36	565	175	52	78	1,090	4,913
North Dakota	115	.	*	15	*	.	*	153	291
Ohio	3,378	*	12	620	55	37	105	2,358	6,567
Oklahoma	746	*	10	173	*	*	12	547	1,503
Oregon	612	*	28	192	27	*	59	518	1,439
Pennsylvania	4,545	44	42	551	54	39	157	2,531	7,963
Rhode Island	375	.	.	27	*	.	*	170	587
South Carolina	1,805	46	17	217	49	*	77	687	2,904
South Dakota	122	.	17	13	*	.	*	134	296
Tennessee	2,074	*	54	292	32	*	49	1,036	3,545
Texas	6,162	43	57	622	185	28	295	3,058	10,450
Utah	241	*	*	34	.	*	*	338	627
Vermont	90	.	11	25	*	.	*	93	222
Virginia	2,683	*	53	399	40	*	34	1,173	4,392
Washington	1,089	10	153	145	17	24	26	865	2,329
West Virginia	430	*	*	116	19	*	*	369	954
Wisconsin	1,102	22	26	250	31	*	55	1,254	2,741
Wyoming	45	*	*	29	*	.	*	78	167
American Samoa	18	18
Guam	40	*	.	.	.	*	*	13	60
Puerto Rico	1,031	.	15	173	10	*	12	354	1,596
Saipan	11	*	*	14
Virgin Islands	52	.	.	*	.	.	.	*	65
Foreign	34	.	*	*	*	.	*	149	200
Unknown	170	.	.	13	.	*	1,038	1,812	3,038
All	89,001	654	1,325	13,671	2,108	457	3,907	50,072	161,195

The point prevalent cohorts and associated modalities are determined on December 31 of each calendar year without applying the 60-day stable modality rule.

* Values for cells with fewer than 10 patients are suppressed.

." Zero values in this cell.

Table D.12 (continued)

Prevalent ESRD Patients for Selected Years: by Diabetic Status & State/Territory

point prevalent patients on December 31

1996 ALL	Center hemo	Center self hemo	Home hemo	CAPD	CCPD	Other PD	Uncertain dialysis	Transplant	All
Alabama	3,945	*	15	376	205	17	41	1,538	6,138
Alaska	386	*	*	39	*	*	15	151	607
Arizona	3,069	20	30	229	193	*	86	1,156	4,787
Arkansas	1,724	.	14	210	53	*	33	753	2,791
California	20,005	46	35	2,003	946	54	1,157	8,974	33,220
Colorado	1,572	*	12	172	92	*	35	1,010	2,897
Connecticut	1,901	.	*	323	241	*	88	968	3,534
Delaware	596	*	.	78	25	*	25	240	966
District of Columbia	1,216	*	*	41	11	*	30	197	1,512
Florida	10,935	23	461	808	519	25	361	4,034	17,166
Georgia	7,000	*	115	536	344	16	97	1,967	10,076
Hawaii	1,163	*	*	94	14	.	19	297	1,591
Idaho	359	.	*	92	30	.	18	334	841
Illinois	8,402	*	125	852	340	19	279	4,341	14,360
Indiana	3,354	*	23	694	147	12	99	1,626	5,957
Iowa	1,209	.	32	194	77	*	55	1,058	2,628
Kansas	1,248	*	16	270	78	*	29	668	2,313
Kentucky	2,006	.	*	231	123	17	92	1,224	3,702
Louisiana	4,534	.	20	252	193	*	43	1,182	6,233
Maine	531	.	*	42	32	*	15	371	997
Maryland	4,332	21	50	428	199	13	207	1,320	6,570
Massachusetts	3,472	*	48	253	200	15	71	1,824	5,889
Michigan	6,373	*	61	1,207	214	11	128	3,217	11,212
Minnesota	2,154	.	*	131	74	*	64	2,014	4,447
Mississippi	3,078	*	68	211	62	*	51	641	4,121
Missouri	3,433	*	68	441	230	*	55	1,765	5,998
Montana	348	.	*	40	10	*	*	313	724
Nebraska	630	.	*	274	35	.	20	542	1,505
Nevada	789	.	10	97	39	*	36	430	1,403
New Hampshire	360	.	*	76	39	*	16	307	801
New Jersey	5,923	77	46	626	386	68	317	2,037	9,480
New Mexico	1,112	.	*	173	91	21	22	528	1,950
New York	13,458	211	105	1,444	507	76	321	4,486	20,608
North Carolina	6,430	*	52	720	439	75	116	1,796	9,631
North Dakota	243	.	.	37	16	.	12	252	560
Ohio	6,870	*	76	1,072	396	53	217	3,730	12,415
Oklahoma	1,884	*	13	224	81	12	28	932	3,177
Oregon	1,258	*	47	259	76	*	52	911	2,606
Pennsylvania	9,085	36	66	655	338	34	427	4,020	14,661
Rhode Island	662	.	*	51	28	.	18	287	1,047
South Carolina	3,835	*	35	307	203	*	89	1,154	5,632
South Dakota	411	.	*	40	24	.	13	234	724
Tennessee	4,055	18	42	308	199	*	89	1,635	6,355
Texas	14,613	33	81	984	684	18	505	5,179	22,097
Utah	575	.	*	62	20	*	18	695	1,378
Vermont	194	.	*	47	12	.	*	179	443
Virginia	5,286	*	67	460	249	15	201	1,515	7,796
Washington	2,311	18	144	276	85	11	47	1,553	4,445
West Virginia	940	*	*	117	86	*	62	470	1,693
Wisconsin	2,342	*	40	326	174	*	85	2,302	5,279
Wyoming	131	.	*	23	12	.	*	131	311
American Samoa	39	39
Guam	130	*	.	31	*	.	*	19	185
Puerto Rico	2,204	.	11	398	200	17	46	449	3,325
Saipan	40	*	44
Virgin Islands	90	.	.	*	*	.	*	11	114
Foreign	62	*	*	*	.	.	*	266	343
Unknown	1,305	.	*	69	14	29	3,442	1,888	6,750
All	185,612	552	2,124	19,414	9,093	724	9,430	81,125	308,074

The point prevalent cohorts and associated modalities are determined on December 31 of each calendar year without applying the 60-day stable modality rule.

* Values for cells with fewer than 10 patients are suppressed.

." Zero values in this cell.

Table D.12 (continued)

Prevalent ESRD Patients for Selected Years: by Diabetic Status & State/Territory

point prevalent patients on December 31

1996 DIABETIC	Center hemo	Center self hemo	Home hemo	CAPD	CCPD	Other PD	Uncertain dialysis	Transplant	All
Alabama	1,311	.	*	112	55	*	*	215	1,710
Alaska	136	.	*	19	*	*	.	31	190
Arizona	1,543	*	10	97	69	*	32	285	2,043
Arkansas	557	.	*	95	15	*	10	150	832
California	7,924	11	*	708	269	15	402	1,528	10,861
Colorado	668	.	*	63	21	*	*	281	1,042
Connecticut	648	.	*	110	78	*	30	177	1,047
Delaware	188	.	.	20	*	*	*	51	270
District of Columbia	376	*	*	*	*	*	*	19	415
Florida	3,395	*	165	240	175	*	72	750	4,812
Georgia	2,404	*	34	182	107	*	14	348	3,097
Hawaii	610	*	.	45	*	.	*	31	701
Idaho	135	.	*	42	*	.	*	86	281
Illinois	2,717	.	51	281	103	*	49	777	3,985
Indiana	1,125	.	*	223	47	*	26	310	1,740
Iowa	432	.	*	78	28	*	20	250	816
Kansas	447	*	*	98	28	*	*	139	724
Kentucky	702	.	*	69	41	*	25	195	1,038
Louisiana	1,616	.	*	68	49	*	*	233	1,985
Maine	163	.	.	12	*	*	*	75	262
Maryland	1,434	.	19	110	56	*	41	229	1,890
Massachusetts	1,127	*	11	68	67	*	16	328	1,623
Michigan	2,405	*	16	435	73	*	29	724	3,687
Minnesota	725	.	.	40	30	*	*	590	1,396
Mississippi	1,052	*	*	70	18	*	*	85	1,238
Missouri	1,205	.	25	162	81	*	*	359	1,841
Montana	155	.	*	18	*	.	*	100	281
Nebraska	239	.	*	113	13	.	*	180	552
Nevada	266	.	*	40	18	*	*	85	423
New Hampshire	108	.	.	25	10	.	*	65	212
New Jersey	2,039	22	*	206	129	24	107	364	2,899
New Mexico	648	.	.	81	43	12	10	128	922
New York	4,484	34	18	441	156	29	52	687	5,901
North Carolina	2,307	*	11	221	139	32	23	341	3,075
North Dakota	99	.	.	19	.	.	*	72	195
Ohio	2,449	.	19	419	151	19	53	796	3,906
Oklahoma	836	.	*	94	31	*	*	199	1,177
Oregon	440	.	*	88	24	*	15	223	800
Pennsylvania	3,139	*	13	211	107	15	87	741	4,319
Rhode Island	207	.	*	10	10	.	*	47	282
South Carolina	1,335	.	*	92	65	*	22	199	1,723
South Dakota	203	.	.	13	10	.	*	67	296
Tennessee	1,392	*	*	91	72	*	18	271	1,857
Texas	6,425	15	14	385	231	*	202	1,016	8,296
Utah	263	.	*	28	*	.	*	234	537
Vermont	69	.	*	19	*	.	*	48	142
Virginia	1,722	*	18	134	60	*	26	247	2,213
Washington	834	*	29	107	19	*	*	401	1,400
West Virginia	362	*	*	37	32	*	14	88	538
Wisconsin	817	.	*	108	54	*	31	651	1,671
Wyoming	62	.	*	12	*	.	.	37	117
American Samoa	19	19
Guam	77	*	.	19	.	*	*	*	98
Puerto Rico	957	.	*	161	90	*	*	34	1,260
Saipan	22	22
Virgin Islands	37	37
Foreign	13	*	.	*	.	.	*	*	21
Unknown	463	.	.	31	*	*	266	36	806
All	67,533	121	569	6,679	2,922	270	1,821	15,608	95,523

The point prevalent cohorts and associated modalities are determined on December 31 of each calendar year without applying the 60-day stable modality rule.

* Values for cells with fewer than 10 patients are suppressed.

." Zero values in this cell.

Table D.12 (continued)

Prevalent ESRD Patients for Selected Years: by Diabetic Status & State/Territory

point prevalent patients on December 31

1996 NON-DIABETIC	Center hemo	Center self hemo	Home hemo	CAPD	CCPD	Other PD	Uncertain dialysis	Transplant	All
Alabama	2,634	*	11	264	150	*	36	1,323	4,428
Alaska	250	*	*	20	*	*	15	120	417
Arizona	1,526	16	20	132	124	*	54	871	2,744
Arkansas	1,167	.	10	115	38	*	23	603	1,959
California	12,081	35	31	1,295	677	39	755	7,446	22,359
Colorado	904	*	*	109	71	*	30	729	1,855
Connecticut	1,253	.	*	213	163	*	58	791	2,487
Delaware	408	*	.	58	19	.	21	189	696
District of Columbia	840	*	*	33	10	*	26	178	1,097
Florida	7,540	17	296	568	344	16	289	3,284	12,354
Georgia	4,596	.	81	354	237	*	83	1,619	6,979
Hawaii	553	.	*	49	*	.	10	266	890
Idaho	224	.	*	50	23	.	10	248	560
Illinois	5,685	*	74	571	237	12	230	3,564	10,375
Indiana	2,229	*	16	471	100	10	73	1,316	4,217
Iowa	777	.	26	116	49	*	35	808	1,812
Kansas	801	.	13	172	50	*	23	529	1,589
Kentucky	1,304	.	*	162	82	13	67	1,029	2,664
Louisiana	2,918	.	14	184	144	*	35	949	4,248
Maine	368	.	*	30	26	.	10	296	735
Maryland	2,898	21	31	318	143	12	166	1,091	4,680
Massachusetts	2,345	*	37	185	133	10	55	1,496	4,266
Michigan	3,968	.	45	772	141	*	99	2,493	7,525
Minnesota	1,429	.	*	91	44	*	55	1,424	3,051
Mississippi	2,026	*	62	141	44	*	48	556	2,883
Missouri	2,228	*	43	279	149	*	49	1,406	4,157
Montana	193	.	*	22	*	*	*	213	443
Nebraska	391	.	*	161	22	.	14	362	953
Nevada	523	.	*	57	21	*	28	345	980
New Hampshire	252	.	*	51	29	*	12	242	589
New Jersey	3,884	55	38	420	257	44	210	1,673	6,581
New Mexico	464	.	*	92	48	*	12	400	1,028
New York	8,974	177	87	1,003	351	47	269	3,799	14,707
North Carolina	4,123	*	41	499	300	43	93	1,455	6,556
North Dakota	144	.	.	18	16	.	*	180	365
Ohio	4,421	*	57	653	245	34	164	2,934	8,509
Oklahoma	1,048	*	*	130	50	*	22	733	2,000
Oregon	818	*	38	171	52	*	37	688	1,806
Pennsylvania	5,946	30	53	444	231	19	340	3,279	10,342
Rhode Island	455	.	.	41	18	.	11	240	765
South Carolina	2,500	*	28	215	138	*	67	955	3,909
South Dakota	208	.	*	27	14	.	10	167	428
Tennessee	2,663	14	38	217	127	*	71	1,364	4,498
Texas	8,188	18	67	599	453	10	303	4,163	13,801
Utah	312	.	*	34	12	*	16	461	841
Vermont	125	.	*	28	*	.	*	131	301
Virginia	3,564	*	49	326	189	10	175	1,268	5,583
Washington	1,477	12	115	169	66	10	44	1,152	3,045
West Virginia	578	*	*	80	54	*	48	382	1,155
Wisconsin	1,525	*	32	218	120	*	54	1,651	3,608
Wyoming	69	.	*	11	*	.	*	94	194
American Samoa	20	20
Guam	53	*	.	12	*	.	*	18	87
Puerto Rico	1,247	.	*	237	110	10	38	415	2,065
Saipan	18	*	22
Virgin Islands	53	.	.	*	*	.	*	11	77
Foreign	49	.	*	*	.	.	*	262	322
Unknown	842	.	*	38	12	21	3,176	1,852	5,944
All	118,079	431	1,555	12,735	6,171	454	7,609	65,517	212,551

The point prevalent cohorts and associated modalities are determined on December 31 of each calendar year without applying the 60-day stable modality rule.

* Values for cells with fewer than 10 patients are suppressed.

." Zero values in this cell.

Table D.12 (continued)

Prevalent ESRD Patients for Selected Years: by Diabetic Status & State/Territory

point prevalent patients on December 31

2000	Center	Center	Home				Uncertain		
All	hemo	self hemo	hemo	CAPD	CCPD	Other PD	dialysis	Transplant	All
Alabama	5,131	.	22	241	255	*	84	1,887	7,629
Alaska	169	.	*	46	*	.	*	193	421
Arizona	4,220	10	27	197	156	*	75	1,525	6,213
Arkansas	2,138	.	*	196	83	*	30	925	3,379
California	26,937	20	51	1,471	1,060	24	632	11,546	41,741
Colorado	2,112	.	18	92	91	.	57	1,315	3,685
Connecticut	2,472	*	11	282	244	*	55	1,157	4,228
Delaware	756	.	.	15	40	.	30	335	1,176
District of Columbia	1,499	.	*	28	20	*	21	241	1,812
Florida	13,836	*	179	426	670	17	255	5,204	20,594
Georgia	9,225	*	135	399	344	15	204	2,472	12,795
Hawaii	1,470	*	*	82	31	.	27	396	2,012
Idaho	498	.	*	91	46	.	10	444	1,096
Illinois	10,956	*	103	569	478	12	249	5,339	17,709
Indiana	4,778	*	26	410	235	*	99	2,081	7,637
Iowa	1,661	*	23	128	96	.	43	1,274	3,226
Kansas	1,737	.	17	193	80	*	41	815	2,885
Kentucky	2,837	.	*	131	122	13	80	1,426	4,615
Louisiana	6,183	*	29	167	193	*	61	1,496	8,138
Maine	705	.	*	26	37	*	13	454	1,240
Maryland	5,488	*	21	293	219	*	116	2,274	8,421
Massachusetts	4,020	.	19	173	184	14	104	2,316	6,830
Michigan	8,472	.	28	694	293	*	178	4,064	13,731
Minnesota	2,606	.	*	74	84	*	51	2,607	5,426
Mississippi	3,836	*	53	163	176	*	45	847	5,127
Missouri	4,645	*	45	264	252	*	82	2,122	7,413
Montana	456	.	*	35	20	*	*	388	912
Nebraska	984	*	*	196	52	.	24	658	1,919
Nevada	1,138	.	*	85	53	*	41	580	1,906
New Hampshire	536	.	*	55	29	.	26	394	1,044
New Jersey	7,994	17	33	262	372	15	192	2,875	11,760
New Mexico	1,688	.	*	104	118	*	34	631	2,582
New York	17,973	199	77	732	607	54	333	6,028	26,003
North Carolina	8,769	.	40	396	461	127	138	2,377	12,308
North Dakota	301	.	.	36	10	.	13	327	687
Ohio	9,693	.	24	707	494	*	186	4,488	15,600
Oklahoma	2,627	.	11	184	167	24	49	1,094	4,156
Oregon	1,637	.	19	150	111	*	30	1,196	3,144
Pennsylvania	11,237	*	35	366	382	12	287	5,038	17,358
Rhode Island	763	.	.	31	15	.	25	391	1,225
South Carolina	4,932	*	15	170	319	*	88	1,499	7,028
South Dakota	551	.	*	30	34	.	16	300	932
Tennessee	5,294	.	51	197	263	10	117	1,997	7,929
Texas	20,307	.	81	639	871	10	328	6,691	28,927
Utah	768	*	*	67	18	.	30	901	1,789
Vermont	322	.	*	*	17	.	12	217	580
Virginia	6,930	*	47	345	300	12	143	2,272	10,051
Washington	3,035	*	86	178	127	*	57	2,092	5,591
West Virginia	1,361	*	*	93	129	*	29	699	2,317
Wisconsin	3,168	*	15	215	157	*	80	2,832	6,472
Wyoming	213	.	*	14	17	.	*	163	414
American Samoa	57	*	*	59
Guam	224	.	.	10	*	.	*	17	255
Puerto Rico	2,730	.	16	142	333	*	47	659	3,932
Saipan	77	*	86
Virgin Islands	119	.	.	*	*	.	*	15	141
Foreign	49	.	.	*	.	.	*	341	401
Unknown	90	.	.	*	*	.	195	1,884	2,175
All	244,410	282	1,429	12,307	10,980	458	5,187	103,809	378,862

The point prevalent cohorts and associated modalities are determined on December 31 of each calendar year without applying the 60-day stable modality rule.

* Values for cells with fewer than 10 patients are suppressed.

." Zero values in this cell.

Table D.12 (continued)

Prevalent ESRD Patients for Selected Years: by Diabetic Status & State/Territory

point prevalent patients on December 31

2000 DIABETIC	Center hemo	Center self hemo	Home hemo	CAPD	CCPD	Other PD	Uncertain dialysis	Transplant	All
Alabama	2,064	.	*	64	80	.	17	302	2,537
Alaska	76	.	*	20	.	.	.	42	139
Arizona	2,275	.	13	72	58	*	17	396	2,833
Arkansas	778	.	*	77	21	.	*	201	1,084
California	11,874	*	12	554	306	*	133	2,094	14,987
Colorado	986	.	*	38	23	.	*	360	1,419
Connecticut	947	.	*	117	88	*	11	198	1,368
Delaware	309	.	.	*	11	.	11	79	412
District of Columbia	463	.	.	*	*	.	*	42	517
Florida	5,015	*	53	134	199	10	39	1,001	6,452
Georgia	3,566	.	43	115	96	*	42	474	4,344
Hawaii	814	.	*	41	16	.	10	55	937
Idaho	219	.	*	45	23	.	*	119	410
Illinois	4,067	*	44	202	160	*	51	1,053	5,584
Indiana	1,859	.	12	152	85	*	11	396	2,520
Iowa	655	.	*	58	30	.	*	318	1,073
Kansas	748	.	*	75	28	*	*	170	1,033
Kentucky	1,158	.	*	52	35	*	13	248	1,514
Louisiana	2,426	.	*	56	56	*	*	321	2,873
Maine	266	.	.	*	15	*	*	91	381
Maryland	1,942	.	*	76	68	*	19	478	2,590
Massachusetts	1,423	.	*	57	60	*	28	421	1,997
Michigan	3,533	.	*	257	94	*	33	966	4,892
Minnesota	966	.	.	20	20	.	11	742	1,759
Mississippi	1,447	*	*	58	63	*	*	123	1,708
Missouri	1,902	.	10	90	84	.	21	443	2,550
Montana	208	.	.	18	13	.	*	134	375
Nebraska	414	*	*	71	20	.	*	215	730
Nevada	465	.	*	39	23	*	*	128	665
New Hampshire	180	.	*	19	*	.	*	86	299
New Jersey	3,100	*	*	76	130	*	33	560	3,910
New Mexico	1,048	.	*	55	61	*	11	183	1,360
New York	6,689	49	12	258	178	24	49	1,013	8,272
North Carolina	3,561	.	*	109	117	47	26	508	4,373
North Dakota	131	.	.	13	*	.	*	90	242
Ohio	4,092	.	*	312	196	*	41	1,008	5,655
Oklahoma	1,243	.	*	99	74	12	*	259	1,698
Oregon	609	.	*	57	42	*	*	286	1,008
Pennsylvania	4,314	.	*	122	121	*	72	975	5,616
Rhode Island	277	.	.	11	*	.	*	68	367
South Carolina	2,006	.	*	46	104	*	15	266	2,442
South Dakota	272	.	.	16	*	.	*	88	390
Tennessee	2,007	.	*	57	92	*	28	359	2,554
Texas	9,920	.	26	259	333	*	58	1,416	12,014
Utah	351	*	*	34	*	.	*	291	691
Vermont	132	.	*	*	*	.	*	48	195
Virginia	2,674	.	10	131	85	*	20	406	3,330
Washington	1,235	*	23	64	42	*	13	525	1,906
West Virginia	659	*	.	37	57	*	*	147	908
Wisconsin	1,177	.	*	72	44	*	19	792	2,107
Wyoming	102	.	*	*	*	.	*	47	163
American Samoa	34	34
Guam	130	.	.	*	*	.	.	.	135
Puerto Rico	1,363	.	*	63	138	*	*	66	1,641
Saipan	53	53
Virgin Islands	49	.	.	*	*	.	.	*	53
Foreign	14	.	.	*	.	.	.	*	19
Unknown	15	10	30	55
All	100,302	67	370	4,497	3,639	185	981	21,132	131,173

The point prevalent cohorts and associated modalities are determined on December 31 of each calendar year without applying the 60-day stable modality rule.

* Values for cells with fewer than 10 patients are suppressed.

." Zero values in this cell.

Table D.12 (continued)

Prevalent ESRD Patients for Selected Years: by Diabetic Status & State/Territory

point prevalent patients on December 31

2000 NON-DIABETIC	Center hemo	Center self hemo	Home hemo	CAPD	CCPD	Other PD	Uncertain dialysis	Transplant	All
Alabama	3,067	.	16	177	175	*	67	1,585	5,092
Alaska	93	.	*	26	*	.	*	151	282
Arizona	1,945	10	14	125	98	*	58	1,129	3,380
Arkansas	1,360	.	*	119	62	*	24	724	2,295
California	15,063	12	39	917	754	18	499	9,452	26,754
Colorado	1,126	.	11	54	68	.	52	955	2,266
Connecticut	1,525	*	*	165	156	*	44	959	2,860
Delaware	447	.	.	13	29	.	19	256	764
District of Columbia	1,036	.	*	20	18	*	19	199	1,295
Florida	8,821	*	126	292	471	*	216	4,203	14,142
Georgia	5,659	*	92	284	248	*	162	1,998	8,451
Hawaii	656	*	*	41	15	.	17	341	1,075
Idaho	279	.	*	46	23	.	*	325	686
Illinois	6,889	*	59	367	318	*	198	4,286	12,125
Indiana	2,919	*	14	258	150	*	88	1,685	5,117
Iowa	1,006	*	18	70	66	.	36	956	2,153
Kansas	989	.	15	118	52	*	32	645	1,852
Kentucky	1,679	.	*	79	87	*	67	1,178	3,101
Louisiana	3,757	*	21	111	137	*	57	1,175	5,265
Maine	439	.	*	19	22	.	12	363	859
Maryland	3,546	*	17	217	151	*	97	1,796	5,831
Massachusetts	2,597	.	17	116	124	*	76	1,895	4,833
Michigan	4,939	.	20	437	199	*	145	3,098	8,839
Minnesota	1,640	.	*	54	64	*	40	1,865	3,667
Mississippi	2,389	*	45	105	113	*	38	724	3,419
Missouri	2,743	*	35	174	168	*	61	1,679	4,863
Montana	248	.	*	17	*	*	*	254	537
Nebraska	570	.	*	125	32	.	17	443	1,189
Nevada	673	.	*	46	30	*	35	452	1,241
New Hampshire	356	.	*	36	20	.	22	308	745
New Jersey	4,894	15	27	186	242	12	159	2,315	7,850
New Mexico	640	.	*	49	57	*	23	448	1,222
New York	11,284	150	65	474	429	30	284	5,015	17,731
North Carolina	5,208	.	35	287	344	80	112	1,869	7,935
North Dakota	170	.	.	23	*	.	*	237	445
Ohio	5,601	.	20	395	298	*	145	3,480	9,945
Oklahoma	1,384	.	*	85	93	12	41	835	2,458
Oregon	1,028	.	15	93	69	.	21	910	2,136
Pennsylvania	6,923	*	28	244	261	*	215	4,063	11,742
Rhode Island	486	.	.	20	11	.	18	323	858
South Carolina	2,926	*	13	124	215	*	73	1,233	4,586
South Dakota	279	.	*	14	25	.	11	212	542
Tennessee	3,287	.	44	140	171	*	89	1,638	5,375
Texas	10,387	.	55	380	538	*	270	5,275	16,913
Utah	417	.	*	33	11	.	24	610	1,098
Vermont	190	.	*	*	*	.	*	169	385
Virginia	4,256	*	37	214	215	*	123	1,866	6,721
Washington	1,800	*	63	114	85	*	44	1,567	3,685
West Virginia	702	.	*	56	72	*	24	552	1,409
Wisconsin	1,991	*	13	143	113	*	61	2,040	4,365
Wyoming	111	.	*	10	*	.	*	116	251
American Samoa	23	*	*	25
Guam	94	.	.	*	*	.	*	17	120
Puerto Rico	1,367	.	11	79	195	*	44	593	2,291
Saipan	24	*	33
Virgin Islands	70	.	.	*	*	.	*	14	88
Foreign	35	.	.	*	.	.	*	337	382
Unknown	75	.	.	*	*	.	185	1,854	2,120
All	144,108	215	1,059	7,810	7,341	273	4,206	82,677	247,689

The point prevalent cohorts and associated modalities are determined on December 31 of each calendar year without applying the 60-day stable modality rule.

* Values for cells with fewer than 10 patients are suppressed.

." Zero values in this cell.

Table D.13

Prevalent Dialysis Patients: by Age & State/Territory*point prevalent dialysis patients on December 31, 1998-2000 combined, by state & age; grouped by modality*

ALL	Center hemo	Center self hemo	Home hemo	CAPD	CCPD	Other PD	Uncertain dialysis	All
Alabama	14,618	*	53	815	695	25	169	16,377
Alaska	517	*	*	108	19	.	10	659
Arizona	11,930	36	94	604	465	17	221	13,367
Arkansas	6,188	*	19	589	200	*	87	7,093
California	75,772	80	147	4,800	3,076	69	2,629	86,573
Colorado	6,018	.	45	337	302	*	135	6,841
Connecticut	6,987	*	25	894	745	14	177	8,843
Delaware	2,154	.	*	71	119	.	62	2,408
District of Columbia	4,364	.	*	61	67	*	63	4,568
Florida	40,037	27	536	1,481	1,899	63	879	44,922
Georgia	26,082	*	363	1,326	972	40	389	29,175
Hawaii	4,143	*	14	237	81	.	112	4,590
Idaho	1,451	*	25	255	124	.	35	1,891
Illinois	31,369	*	249	1,796	1,404	39	696	35,560
Indiana	13,350	*	51	1,387	693	20	257	15,759
Iowa	4,706	*	63	428	260	.	172	5,631
Kansas	4,927	.	51	614	235	*	164	5,994
Kentucky	7,914	*	17	437	385	54	245	9,053
Louisiana	17,691	*	62	506	583	23	135	19,001
Maine	2,065	.	*	77	97	*	43	2,294
Maryland	16,001	*	53	965	645	29	341	18,042
Massachusetts	11,861	.	67	577	556	44	311	13,416
Michigan	24,043	*	74	2,341	867	11	405	27,742
Minnesota	7,505	.	10	261	278	*	162	8,220
Mississippi	11,177	*	140	518	417	21	112	12,391
Missouri	13,167	*	136	988	714	16	263	15,285
Montana	1,321	*	11	111	45	*	13	1,504
Nebraska	2,613	*	*	737	163	*	86	3,610
Nevada	3,185	*	14	236	145	12	126	3,720
New Hampshire	1,466	*	*	148	84	.	61	1,768
New Jersey	22,891	76	65	1,100	1,207	23	732	26,094
New Mexico	4,632	*	11	317	343	*	78	5,385
New York	51,242	589	240	2,729	1,751	185	819	57,555
North Carolina	24,588	*	88	1,485	1,355	338	328	28,183
North Dakota	892	.	.	105	41	.	40	1,078
Ohio	27,207	*	55	2,271	1,355	93	507	31,491
Oklahoma	7,451	*	21	571	434	51	112	8,647
Oregon	4,632	.	70	560	293	*	151	5,709
Pennsylvania	32,460	29	105	1,256	1,159	29	798	35,836
Rhode Island	2,231	.	.	96	40	.	72	2,439
South Carolina	14,168	*	55	600	905	12	271	16,014
South Dakota	1,543	.	*	113	99	*	37	1,795
Tennessee	15,139	*	114	621	758	21	242	16,896
Texas	56,444	43	240	1,975	2,415	43	1,540	62,700
Utah	2,144	*	10	185	69	*	66	2,476
Vermont	882	.	16	42	58	*	25	1,024
Virginia	20,024	*	138	1,073	920	29	376	22,563
Washington	8,626	30	275	585	354	26	134	10,030
West Virginia	3,788	*	14	292	356	*	83	4,539
Wisconsin	9,050	*	65	720	452	*	256	10,551
Wyoming	572	.	16	51	43	*	10	693
American Samoa	138	*	140
Guam	629	*	.	34	*	.	*	674
Puerto Rico	7,903	*	35	587	915	12	131	9,585
Saipan	208	*	214
Virgin Islands	369	*	.	*	*	.	*	386
Foreign	158	*	*	*	.	.	22	190
Unknown	417	.	.	14	13	*	969	1,416
All	695,050	983	3,991	41,102	31,679	1,418	16,377	790,600

The point prevalent cohorts and associated modalities are determined on December 31 of each calendar year without applying the 60-day stable modality rule.

The age of the point prevalent patient is determined as of December 31 of each calendar year.

Patients with unknown age are excluded.

* Values for cells with fewer than 10 patients are suppressed.

." Zero values in this cell.

Table D.13 (continued)

Prevalent Dialysis Patients: by Age & State/Territory

point prevalent dialysis patients on December 31, 1998-2000 combined, by state & age; grouped by modality

UNDER 65	Center hemo	Center self hemo	Home hemo	CAPD	CCPD	Other PD	Uncertain dialysis	All
Alabama	8,374	*	39	634	518	22	123	9,712
Alaska	350	.	*	77	15	.	10	456
Arizona	6,517	34	45	465	328	12	150	7,551
Arkansas	3,600	.	10	409	154	*	59	4,241
California	41,018	55	106	3,518	2,314	44	1,545	48,600
Colorado	3,347	.	34	237	217	*	111	3,948
Connecticut	3,225	.	18	596	475	*	106	4,429
Delaware	1,215	.	*	43	91	.	49	1,400
District of Columbia	2,731	.	*	50	59	*	44	2,896
Florida	20,790	24	294	1,066	1,308	36	507	24,025
Georgia	16,287	*	235	1,089	770	30	288	18,701
Hawaii	2,293	*	12	184	54	.	38	2,582
Idaho	746	*	15	160	98	.	25	1,045
Illinois	16,511	*	101	1,231	899	26	503	19,276
Indiana	6,764	.	28	967	444	18	171	8,392
Iowa	2,141	*	44	218	158	.	119	2,682
Kansas	2,473	.	40	412	151	*	117	3,196
Kentucky	4,305	*	16	350	279	42	148	5,141
Louisiana	10,719	.	37	394	443	22	107	11,722
Maine	924	.	*	46	64	*	31	1,073
Maryland	9,027	*	41	672	435	27	266	10,472
Massachusetts	5,028	.	39	382	357	18	209	6,033
Michigan	12,740	*	50	1,609	586	*	305	15,299
Minnesota	3,243	.	*	132	180	*	117	3,683
Mississippi	6,896	*	127	392	322	21	84	7,848
Missouri	6,764	.	81	656	500	13	193	8,207
Montana	696	*	*	63	29	.	*	804
Nebraska	1,375	*	*	377	99	*	61	1,920
Nevada	1,825	*	13	169	97	*	89	2,203
New Hampshire	652	*	*	101	55	.	41	856
New Jersey	11,504	62	47	727	763	*	521	13,632
New Mexico	2,593	*	*	233	257	*	56	3,151
New York	27,536	426	160	1,955	1,211	126	636	32,050
North Carolina	14,524	.	75	1,140	1,019	233	260	17,251
North Dakota	470	.	.	42	29	.	27	568
Ohio	13,831	.	36	1,589	878	66	348	16,748
Oklahoma	4,234	*	16	402	319	33	90	5,098
Oregon	2,428	.	32	375	199	*	100	3,137
Pennsylvania	15,570	26	78	858	797	21	480	17,830
Rhode Island	913	.	.	70	30	.	35	1,048
South Carolina	8,658	*	54	504	669	*	200	10,097
South Dakota	741	.	.	80	72	.	30	923
Tennessee	8,782	.	99	461	556	15	183	10,096
Texas	33,807	31	169	1,485	1,768	34	1,068	38,362
Utah	1,188	*	*	121	49	*	46	1,413
Vermont	376	.	13	33	31	*	11	465
Virginia	11,219	*	91	801	640	19	284	13,055
Washington	4,728	10	154	394	249	21	97	5,653
West Virginia	1,840	*	*	185	233	.	43	2,309
Wisconsin	4,143	.	51	364	229	*	160	4,950
Wyoming	314	.	12	33	34	*	*	399
American Samoa	107	*	109
Guam	406	.	.	30	*	.	*	445
Puerto Rico	5,058	*	29	459	746	10	101	6,405
Saipan	161	*	163
Virgin Islands	253	.	.	*	*	.	*	265
Foreign	74	*	.	*	.	.	15	92
Unknown	290	.	.	12	10	*	779	1,094
All	378,324	712	2,610	29,059	22,295	990	11,211	445,201

The point prevalent cohorts and associated modalities are determined on December 31 of each calendar year without applying the 60-day stable modality rule. The age of the point prevalent patient is determined as of December 31 of each calendar year. Patients with unknown age are excluded. * Values for cells with fewer than 10 patients are suppressed. "." Zero values in this cell.

Table D.13 (continued)

Prevalent Dialysis Patients: by Age & State/Territory*point prevalent dialysis patients on December 31, 1998-2000 combined, by state & age; grouped by modality*

65+	Center hemo	Center self hemo	Home hemo	CAPD	CCPD	Other PD	Uncertain dialysis	All
Alabama	6,244	.	14	181	177	*	46	6,665
Alaska	167	*	.	31	*	.	.	203
Arizona	5,413	*	49	139	137	*	71	5,816
Arkansas	2,588	*	*	180	46	.	28	2,852
California	34,753	25	41	1,282	762	25	1,084	37,972
Colorado	2,671	.	11	100	85	*	24	2,893
Connecticut	3,762	*	*	298	270	*	71	4,414
Delaware	939	.	.	28	28	.	13	1,008
District of Columbia	1,633	.	.	11	*	*	19	1,672
Florida	19,247	*	242	415	591	27	372	20,897
Georgia	9,795	*	128	237	202	10	101	10,474
Hawaii	1,850	*	*	53	27	.	74	2,008
Idaho	705	.	10	95	26	.	10	846
Illinois	14,858	*	148	565	505	13	192	16,283
Indiana	6,586	*	23	420	249	*	86	7,367
Iowa	2,565	.	19	210	102	.	53	2,949
Kansas	2,454	.	11	202	84	.	47	2,798
Kentucky	3,608	.	*	87	106	12	97	3,911
Louisiana	6,972	*	25	112	140	*	28	7,279
Maine	1,141	.	*	31	33	*	12	1,221
Maryland	6,974	*	12	293	210	*	75	7,570
Massachusetts	6,833	.	28	195	199	26	102	7,383
Michigan	11,303	.	24	732	281	*	100	12,443
Minnesota	4,262	.	*	129	98	*	45	4,537
Mississippi	4,281	.	13	126	95	.	28	4,543
Missouri	6,403	*	55	332	214	*	70	7,078
Montana	625	.	*	48	16	*	*	700
Nebraska	1,238	.	*	360	64	*	25	1,690
Nevada	1,360	.	*	67	48	*	37	1,517
New Hampshire	814	.	*	47	29	.	20	912
New Jersey	11,387	14	18	373	444	15	211	12,462
New Mexico	2,039	.	*	84	86	.	22	2,234
New York	23,703	163	80	773	540	59	183	25,501
North Carolina	10,063	*	13	345	336	105	68	10,931
North Dakota	422	.	.	63	12	.	13	510
Ohio	13,376	*	19	682	477	27	159	14,743
Oklahoma	3,217	*	*	169	115	18	22	3,549
Oregon	2,204	.	38	185	94	.	51	2,572
Pennsylvania	16,889	*	27	398	362	*	318	18,005
Rhode Island	1,318	.	.	26	10	.	37	1,391
South Carolina	5,508	.	*	96	236	*	71	5,915
South Dakota	802	.	*	33	27	*	*	872
Tennessee	6,357	*	15	160	202	*	59	6,800
Texas	22,637	12	71	490	647	*	470	24,336
Utah	956	.	*	64	20	.	20	1,063
Vermont	506	.	*	*	27	.	14	559
Virginia	8,805	*	47	272	280	10	92	9,508
Washington	3,898	20	121	191	105	*	37	4,377
West Virginia	1,948	*	*	107	123	*	40	2,230
Wisconsin	4,907	*	14	356	223	*	96	5,601
Wyoming	258	.	*	18	*	.	*	294
American Samoa	31	31
Guam	223	*	.	*	.	.	*	229
Puerto Rico	2,845	.	*	128	169	*	30	3,180
Saipan	47	*	51
Virgin Islands	116	*	.	*	*	.	*	121
Foreign	84	.	*	*	.	.	*	98
Unknown	74	.	.	*	*	.	189	268
All	316,664	271	1,381	12,042	9,384	428	5,162	345,332

The point prevalent cohorts and associated modalities are determined on December 31 of each calendar year without applying the 60-day stable modality rule.

The age of the point prevalent patient is determined as of December 31 of each calendar year.

Patients with unknown age are excluded.

* Values for cells with fewer than 10 patients are suppressed.

." Zero values in this cell.

