

■

# Reference Section D

Treatment modalities

table d.1 · 311  
counts & percentages of reported ESRD patients, by treatment modality  
*incident & point prevalent patients, 1989-2001 combined*

table d.2 · 313  
incident ESRD patients, by dialysis/functioning graft status, by demographic characteristics & primary diagnosis

table d.3 · 317  
prevalent ESRD patients, by dialysis/functioning graft status, by demographic characteristics & primary diagnosis  
*point prevalent patients on December 31*

table d.4 · 321  
incident ESRD patients for selected years, by age & modality

table d.5 · 323  
prevalent ESRD patients for selected years, by age & modality  
*point prevalent patients on December 31*

table d.6 · 325  
incident ESRD patients for selected years  
*by gender, race, ethnicity, & primary diagnosis; grouped by modality*

table d.7 · 327  
prevalent ESRD patients for selected years  
*point prevalent patients on December 31, by gender, race, ethnicity, & primary diagnosis; grouped by modality*

table d.8 · 329  
treatment status two years after first ESRD service  
*incident patients, 1997-1999 combined, by gender, age at first service, & modality at 90 days*

table d.9 · 332  
prevalent ESRD patients for selected years, by patient vintage  
*point prevalent patients on December 31, by year-end treatment modality & number of years patient has had ESRD (vintage)*

table d.10 · 334  
prevalent ESRD patients for selected years, by number of years on modality  
*point prevalent patients on December 31, by year-end treatment modality*

table d.11 · 336  
incident ESRD patients for selected years, by modality, diabetic status, & state/territory

table d.12 · 345  
prevalent ESRD patients for selected years, by modality, diabetic status, & state/territory  
*point prevalent patients on December 31*

table d.13 · 354  
prevalent dialysis patients, by modality, age, & state/territory  
*point prevalent dialysis patients on December 31, 1999-2001 combined, by state, age, & modality*

Table D.1

**Counts & Percentages of Reported ESRD Patients, by Treatment Modality***incident & point prevalent patients, 1989-2001 combined*

<b>INCIDENT COUNTS</b>	<b>1978</b>	<b>1979</b>	<b>1980</b>	<b>1981</b>	<b>1982</b>	<b>1983</b>	<b>1984</b>	<b>1985</b>	<b>1986</b>	<b>1987</b>	<b>1988</b>
<b>Modality</b>											
Dialysis	14,088	15,806	16,952	19,069	21,542	25,107	26,257	29,028	31,497	34,422	38,090
Functioning graft	386	420	452	450	566	572	642	758	962	1,257	1,847
<b>Dialysis type</b>											
Center hemodialysis	12,265	13,803	14,731	16,278	17,851	20,720	21,580	23,288	25,440	28,053	31,341
Center self hemodialysis	309	479	359	201	119	172	157	102	104	94	95
Home hemodialysis	57	56	52	23	14	22	38	204	225	156	116
CAPD	491	603	912	1,840	3,135	3,791	3,929	4,027	4,069	4,383	4,635
CCPD	*	*	.	*	*	*	23	115	193	269	260
Other PD	405	664	596	364	190	210	220	538	690	619	639
Uncertain dialysis	560	200	302	361	231	189	310	754	776	848	1,004
<b>All</b>	<b>14,474</b>	<b>16,226</b>	<b>17,404</b>	<b>19,519</b>	<b>22,108</b>	<b>25,679</b>	<b>26,899</b>	<b>29,786</b>	<b>32,459</b>	<b>35,679</b>	<b>39,937</b>
<b>PERCENTAGES</b>											
<b>Modality</b>											
Dialysis	97.3	97.4	97.4	97.7	97.4	97.8	97.6	97.5	97.0	96.5	95.4
Functioning graft	2.7	2.6	2.6	2.3	2.6	2.2	2.4	2.5	3.0	3.5	4.6
<b>Dialysis type</b>											
Center hemodialysis	87.1	87.3	86.9	85.4	82.9	82.5	82.2	80.2	80.8	81.5	82.3
Center self hemodialysis	2.2	3.0	2.1	1.1	0.6	0.7	0.6	0.4	0.3	0.3	0.2
Home hemo	0.4	0.4	0.3	0.1	0.1	0.1	0.1	0.7	0.7	0.5	0.3
CAPD	3.5	3.8	5.4	9.6	14.6	15.1	15.0	13.9	12.9	12.7	12.2
CCPD	.	.	.	.	.	.	0.1	0.4	0.6	0.8	0.7
Other PD	2.9	4.2	3.5	1.9	0.9	0.8	0.8	1.9	2.2	1.8	1.7
Uncertain dialysis	4.0	1.3	1.8	1.9	1.1	0.8	1.2	2.6	2.5	2.5	2.6
<b>All</b>	<b>100.0</b>	<b>100.0</b>	<b>100.0</b>	<b>100.0</b>	<b>100.0</b>	<b>100.0</b>	<b>100.0</b>	<b>100.0</b>	<b>100.0</b>	<b>100.0</b>	<b>100.0</b>
<b>POINT PREVALENT COUNTS</b>											
<b>Modality</b>											
Dialysis	36,619	42,734	47,743	53,314	60,055	70,464	77,827	83,938	90,172	97,900	106,577
Functioning graft	5,705	7,638	9,702	11,992	14,579	17,571	21,195	25,445	30,542	35,378	40,244
<b>Dialysis type</b>											
Center hemodialysis	31,412	36,948	40,935	46,590	51,800	58,159	61,958	63,479	68,970	76,415	85,646
Center self hemodialysis	855	965	1,143	1,335	1,461	1,984	1,483	1,325	1,265	1,253	1,253
Home hemodialysis	1,886	2,008	1,854	703	473	1,296	2,653	5,825	5,512	4,679	2,986
CAPD	217	332	736	2,355	3,957	6,550	8,726	10,442	11,217	12,122	12,924
CCPD	.	*	*	*	*	22	218	544	780	934	1,015
Other PD	847	1,254	1,486	1,118	1,081	926	752	895	875	820	772
Uncertain dialysis	1,402	1,226	1,588	1,212	1,280	1,527	2,037	1,428	1,553	1,677	1,981
<b>All</b>	<b>42,324</b>	<b>50,372</b>	<b>57,445</b>	<b>65,306</b>	<b>74,634</b>	<b>88,035</b>	<b>99,022</b>	<b>109,383</b>	<b>120,714</b>	<b>133,278</b>	<b>146,821</b>
<b>PERCENTAGES</b>											
<b>Modality</b>											
Dialysis	86.5	84.8	83.1	81.6	80.5	80.0	78.6	76.7	74.7	73.5	72.6
Functioning graft	13.5	15.2	16.9	18.4	19.5	20.0	21.4	23.3	25.3	26.5	27.4
<b>Dialysis type</b>											
Center hemodialysis	85.8	86.5	85.7	87.4	86.3	82.5	79.6	75.6	76.5	78.1	80.4
Center self hemodialysis	2.3	2.3	2.4	2.5	2.4	2.8	1.9	1.6	1.4	1.3	1.2
Home hemodialysis	5.2	4.7	3.9	1.3	0.8	1.8	3.4	6.9	6.1	4.8	2.8
CAPD	0.6	0.8	1.5	4.4	6.6	9.3	11.2	12.4	12.4	12.4	12.1
CCPD	.	.	.	.	.	.	0.3	0.6	0.9	1.0	1.0
Other PD	2.3	2.9	3.1	2.1	1.8	1.3	1.0	1.1	1.0	0.8	0.7
Uncertain dialysis	3.8	2.9	3.3	2.3	2.1	2.2	2.6	1.7	1.7	1.7	1.9
<b>All</b>	<b>100.0</b>	<b>100.0</b>	<b>100.0</b>	<b>100.0</b>	<b>100.0</b>	<b>100.0</b>	<b>100.0</b>	<b>100.0</b>	<b>100.0</b>	<b>100.0</b>	<b>100.0</b>

The incident cohorts and associated modalities are determined at the time of ESRD initiation without applying the 60-day stable modality rule.

The point prevalent cohorts and associated modalities are determined on December 31 of each calendar year without applying the 60-day stable modality rule.

\* Values for cells with fewer than 10 patients are suppressed.

." Zero values in this cell.

Table D.1 (continued)  
**Counts & Percentages of Reported ESRD Patients, by Treatment Modality**  
*incident & point prevalent patients, 1989-2001 combined*

<b>INCIDENT COUNTS</b>	<b>1989</b>	<b>1990</b>	<b>1991</b>	<b>1992</b>	<b>1993</b>	<b>1994</b>	<b>1995</b>	<b>1996</b>	<b>1997</b>	<b>1998</b>	<b>1999</b>	<b>2000</b>	<b>2001</b>
<b>Modality</b>													
Dialysis	43,254	47,050	52,708	57,628	59,828	70,888	71,194	76,237	80,378	85,407	89,729	92,149	93,883
Functioning graft	1,857	1,897	1,833	1,844	2,102	1,773	1,633	1,596	1,650	1,747	2,218	2,453	2,412
<b>Dialysis type</b>													
Center hemodialysis	36,189	39,358	43,815	48,377	50,238	57,527	60,140	63,961	70,828	76,413	80,521	83,513	86,022
Center self hemodialysis	71	91	74	67	52	41	66	62	53	47	45	31	38
Home hemodialysis	78	89	76	59	59	72	364	1,018	191	228	384	258	229
CAPD	5,042	5,752	5,936	6,091	6,274	6,319	6,538	6,035	5,581	5,143	4,943	4,685	4,711
CCPD	397	512	729	949	891	1,145	1,949	2,217	2,123	2,228	2,254	2,202	2,280
Other PD	642	533	713	752	693	891	653	510	486	428	320	322	299
Uncertain dialysis	835	715	1,365	1,333	1,621	4,893	1,484	2,434	1,116	920	1,262	1,138	304
All	45,111	48,947	54,541	59,472	61,930	72,661	72,827	77,833	82,028	87,154	91,947	94,602	96,295
<b>PERCENTAGES</b>													
<b>Modality</b>													
Dialysis	95.9	96.1	96.6	96.9	96.6	97.6	97.8	97.9	98.0	98.0	97.6	97.4	97.5
Functioning graft	4.1	3.9	3.4	3.1	3.4	2.4	2.2	2.1	2.0	2.0	2.4	2.6	2.5
<b>Dialysis type</b>													
Center hemodialysis	83.7	83.7	83.1	83.9	84.0	81.2	84.5	83.9	88.1	89.5	89.7	90.6	91.6
Center self hemodialysis	0.2	0.2	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.1	.	.
Home hemo	0.2	0.2	0.1	0.1	0.1	0.1	0.5	1.3	0.2	0.3	0.4	0.3	0.2
CAPD	11.7	12.2	11.3	10.6	10.5	8.9	9.2	7.9	6.9	6.0	5.5	5.1	5.0
CCPD	0.9	1.1	1.4	1.6	1.5	1.6	2.7	2.9	2.6	2.6	2.5	2.4	2.4
Other PD	1.5	1.1	1.4	1.3	1.2	1.3	0.9	0.7	0.6	0.5	0.4	0.3	0.3
Uncertain dialysis	1.9	1.5	2.6	2.3	2.7	6.9	2.1	3.2	1.4	1.1	1.4	1.2	0.3
All	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0
<b>POINT PREVALENT COUNTS</b>													
<b>Modality</b>													
Dialysis	118,381	130,952	145,832	161,266	174,873	196,009	211,444	227,410	241,956	256,056	269,545	282,504	292,215
Functioning graft	44,914	50,180	55,626	60,901	66,695	72,027	77,501	83,154	88,851	94,941	100,808	107,151	113,866
<b>Dialysis type</b>													
Center hemodialysis	95,961	105,448	117,551	130,099	143,186	157,753	171,077	187,288	204,868	221,972	236,860	251,655	263,255
Center self hemodialysis	1,069	941	736	810	149	226	523	550	485	370	342	283	333
Home hemodialysis	2,710	2,502	2,195	2,254	672	993	1,985	2,298	1,831	1,523	1,444	1,338	1,122
CAPD	14,499	16,512	18,210	20,029	21,512	22,475	21,532	19,483	17,439	15,413	14,030	12,800	11,927
CCPD	1,343	2,206	2,832	3,709	4,355	5,355	7,790	9,687	10,411	10,616	11,253	11,811	12,341
Other PD	714	573	631	658	626	842	770	647	566	479	445	454	445
Uncertain dialysis	2,085	2,770	3,677	3,707	4,373	8,365	7,767	7,457	6,356	5,683	5,171	4,163	2,792
All	163,295	181,132	201,458	222,167	241,568	268,036	288,945	310,564	330,807	350,997	370,353	389,655	406,081
<b>PERCENTAGES</b>													
<b>Modality</b>													
Dialysis	72.5	72.3	72.4	72.6	72.4	73.1	73.2	73.2	73.1	73.0	72.8	72.5	72.0
Functioning graft	27.5	27.7	27.6	27.4	27.6	26.9	26.8	26.8	26.9	27.0	27.2	27.5	28.0
<b>Dialysis type</b>													
Center hemodialysis	81.1	80.5	80.6	80.7	81.9	80.5	80.9	82.4	84.7	86.7	87.9	89.1	90.1
Center self hemodialysis	0.9	0.7	0.5	0.5	0.1	0.1	0.2	0.2	0.2	0.1	0.1	0.1	0.1
Home hemodialysis	2.3	1.9	1.5	1.4	0.4	0.5	0.9	1.0	0.8	0.6	0.5	0.5	0.4
CAPD	12.2	12.6	12.5	12.4	12.3	11.5	10.2	8.6	7.2	6.0	5.2	4.5	4.1
CCPD	1.1	1.7	1.9	2.3	2.5	2.7	3.7	4.3	4.3	4.1	4.2	4.2	4.2
Other PD	0.6	0.4	0.4	0.4	0.4	0.4	0.4	0.3	0.2	0.2	0.2	0.2	0.2
Uncertain dialysis	1.8	2.1	2.5	2.3	2.5	4.3	3.7	3.3	2.6	2.2	1.9	1.5	1.0
All	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0

The incident cohorts are determined at the time of ESRD initiation without applying the 60-day stable modality rule.

The point prevalent cohorts and associated modalities are determined on December 31 of each calendar year without applying the 60-day stable modality rule.

\* Values for cells with fewer than 10 patients are suppressed.

.. Zero values in this cell.

Table D.2

## Incident ESRD Patients, by Dialysis/Functioning Graft Status, by demographic characteristics &amp; primary diagnosis

ALL DIALYSIS	1978	1979	1980	1981	1982	1983	1984	1985	1986	1987	1988	1989	1990	1991	1992	1993	1994	1995	1996	1997	1998	1999	2000	2001
0-4	21	22	34	52	74	83	85	99	101	93	103	105	98	124	103	117	118	141	145	147	154	144	177	203
5-9	63	69	71	73	83	80	86	73	91	76	70	75	96	80	65	85	122	112	111	92	102	120	120	115
10-14	182	155	147	163	187	177	184	162	142	151	152	138	165	168	171	174	209	218	221	228	217	237	239	247
15-19	374	391	381	427	408	394	365	357	364	385	394	434	409	391	425	404	455	486	509	460	486	522	496	519
20-24	582	576	573	652	699	639	706	703	685	734	751	752	786	827	913	871	1,004	1,012	972	899	956	906	919	971
25-29	764	783	786	899	1,056	1,046	1,096	1,123	1,194	1,144	1,208	1,358	1,383	1,429	1,436	1,505	1,755	1,571	1,607	1,610	1,650	1,578	1,635	1,562
30-34	802	921	1,032	1,193	1,307	1,325	1,398	1,394	1,561	1,471	1,628	1,865	1,877	1,960	2,046	2,093	2,538	2,416	2,359	2,310	2,339	2,362	2,329	2,239
35-39	901	885	984	1,135	1,271	1,321	1,415	1,608	1,692	1,795	2,009	2,103	2,236	2,429	2,682	2,636	3,396	3,188	3,220	3,205	3,192	3,311	3,157	3,211
40-44	964	1,022	1,029	1,247	1,277	1,431	1,461	1,596	1,728	1,954	2,090	2,372	2,662	3,016	3,212	3,216	3,866	4,033	4,150	4,101	4,434	4,396	4,499	4,468
45-49	1,111	1,254	1,219	1,377	1,442	1,637	1,685	1,840	1,932	2,107	2,318	2,698	2,795	3,221	3,540	3,808	4,769	4,888	5,197	5,259	5,689	5,903	6,019	6,077
50-54	1,471	1,585	1,645	1,869	2,042	2,068	2,079	2,264	2,363	2,528	2,885	3,138	3,330	3,606	4,122	4,292	5,593	5,508	5,697	6,253	6,820	7,446	7,642	8,196
55-59	1,663	1,830	1,941	2,358	2,498	2,641	2,819	2,993	3,086	3,275	3,654	3,948	4,278	4,673	5,104	5,223	6,548	6,580	6,796	7,266	7,924	8,472	8,754	9,021
60-64	1,605	1,853	2,066	2,366	2,712	3,170	3,452	3,817	3,970	4,399	4,815	5,265	5,703	6,342	6,661	6,831	8,059	8,239	8,464	8,872	9,423	9,653	9,943	10,252
65-69	1,636	1,905	2,143	2,176	2,534	3,347	3,449	3,897	4,410	5,032	5,541	6,442	7,150	7,953	8,559	8,848	9,823	9,606	10,352	10,774	11,017	11,163	11,325	11,436
70-74	1,152	1,400	1,565	1,671	2,103	2,894	2,902	3,445	3,790	4,299	4,705	5,462	6,148	7,255	8,085	8,523	9,661	9,665	10,748	11,238	11,637	11,975	12,231	12,332
75-79	568	791	906	958	1,206	1,824	1,978	2,271	2,735	3,043	3,483	4,219	4,585	5,365	5,996	6,288	7,240	7,401	8,479	9,335	10,129	11,169	11,404	11,400
80-84	173	278	331	349	455	761	810	1,013	1,208	1,436	1,673	2,065	2,386	2,753	3,227	3,427	4,050	4,290	4,959	5,725	6,124	6,784	7,347	7,475
85+	55	86	99	102	180	259	284	366	440	499	607	811	959	1,112	1,276	1,481	1,659	1,833	2,248	2,598	3,108	3,587	3,870	4,146
Unknown	*	.	.	*	*	10	*	*	*	*	*	*	*	*	*	*	23	*	*	*	*	*	43	13
0-19	640	637	633	715	752	734	720	691	698	705	719	752	768	763	764	780	904	957	986	927	959	1,023	1,032	1,084
20-44	4,013	4,187	4,404	5,126	5,610	5,762	6,076	6,424	6,860	7,098	7,686	8,450	8,944	9,661	10,289	10,321	12,559	12,220	12,308	12,125	12,571	12,553	12,539	12,451
45-64	5,850	6,522	6,871	7,970	8,694	9,516	10,035	10,914	11,351	12,309	13,672	15,049	16,106	17,842	19,427	20,154	24,969	25,215	26,154	27,650	29,856	31,474	32,358	33,546
65-74	2,788	3,305	3,708	3,847	4,637	6,241	6,351	7,342	8,200	9,331	10,246	11,904	13,298	15,208	16,644	17,371	19,484	19,271	21,100	22,012	22,654	23,138	23,556	23,768
75+	796	1,155	1,336	1,409	1,841	2,844	3,072	3,650	4,383	4,978	5,763	7,095	7,930	9,230	10,499	11,196	12,949	13,524	15,686	17,658	19,361	21,540	22,621	23,021
Unknown	*	.	.	*	*	10	*	*	*	*	*	*	*	*	*	*	23	*	*	*	*	*	43	13
Male	7,790	8,750	9,432	10,516	11,831	13,754	14,463	15,763	17,207	18,621	20,524	23,376	25,401	28,207	30,523	31,855	37,177	37,487	40,624	42,678	45,388	47,869	49,165	50,419
Female	6,298	7,055	7,520	8,553	9,709	11,349	11,792	13,263	14,290	15,799	17,565	19,875	21,649	24,498	27,103	27,966	33,687	33,700	35,611	37,696	40,015	41,860	42,946	43,450
Unknown	.	*	.	.	*	*	*	*	.	*	*	*	.	*	*	*	24	*	*	*	*	.	38	14
White	8,847	9,749	10,436	12,103	14,545	16,554	17,353	19,177	20,769	22,465	24,648	28,074	30,622	34,117	36,978	37,535	43,495	43,591	46,525	50,673	53,412	57,037	58,800	59,967
Black	4,701	5,448	5,824	6,211	6,306	7,620	7,775	8,578	9,228	10,253	11,470	13,439	14,358	16,147	17,675	18,813	21,739	22,365	23,828	24,276	25,912	26,571	26,730	27,430
Native American	125	131	174	198	221	288	283	298	371	426	517	616	678	743	904	859	1,042	1,218	1,380	1,025	1,469	1,222	1,258	1,206
Asian	96	124	154	219	366	518	756	862	991	1,135	1,146	944	1,184	1,349	1,541	1,789	2,293	2,489	2,728	2,664	3,074	3,311	3,245	3,373
Other/unknown	319	354	364	338	104	127	90	113	138	143	309	181	208	352	530	832	2,319	1,531	1,776	1,740	1,540	1,588	2,116	1,907
†Hispanic																		8,969	9,049	9,995	10,556	11,365	11,664	
†Non-Hispanic																		67,268	71,329	75,412	79,173	80,784	82,219	
Diabetes	1,410	1,630	2,239	3,624	4,991	5,914	7,055	8,189	9,380	10,516	11,846	14,438	16,293	18,980	21,168	21,150	25,772	28,945	32,788	35,262	37,748	39,737	41,625	42,449
Hypertension	1,862	2,073	2,578	4,045	5,507	6,016	6,758	7,769	8,420	9,586	10,832	12,870	13,958	15,779	17,637	16,922	20,283	18,423	19,929	21,420	22,673	24,031	24,544	25,293
Glomerulonephritis	1,890	2,021	2,112	3,054	3,849	4,036	4,280	4,449	4,578	4,836	5,133	5,515	5,815	6,138	6,396	6,305	7,246	7,515	8,038	7,997	8,158	8,109	7,831	7,662
Cystic kidney	537	585	631	845	1,013	1,052	1,083	1,135	1,209	1,207	1,217	1,220	1,367	1,434	1,532	1,480	1,688	1,762	1,739	1,843	1,947	2,003	1,954	1,958
Other urologic	208	238	274	393	405	490	494	641	627	645	706	779	823	1,100	1,148	1,008	1,140	1,303	1,532	1,457	1,512	1,632	1,561	1,530
Other cause	1,211	1,380	1,552	2,241	3,092	3,485	3,699	4,067	4,081	4,245	4,329	4,826	4,687	4,910	5,234	5,166	6,310	7,876	8,419	8,735	9,399	9,875	10,212	10,341
Unknown cause	1,234	1,335	1,354	1,578	1,347	1,350	1,389	1,404	1,640	1,893	1,934	1,753	1,858	2,258	2,514	2,233	2,497	2,570	2,958	3,108	3,418	3,499	3,770	3,846
Missing cause	5,736	6,544	6,212	3,289	1,338	2,764	1,499	1,374	1,562	1,494	2,093	1,853	2,249	2,109	1,999	5,564	5,952	2,800	834	556	552	843	652	804
All	14,088	15,806	16,952	19,069	21,542	25,107	26,257	29,028	31,497	34,422	38,090	43,254	47,050	52,708	57,628	59,828	70,888	71,194	76,237	80,378	85,407	89,729	92,149	93,883

The incident cohorts are determined at the time of ESRD initiation without applying the 60-day stable modality rule. The age of the incident patient is determined as of the date of ESRD initiation. †CMS began collecting †Hispanic ethnicity data in April 1995; cells for years prior to 1996 have therefore been left blank. \* Values for cells with fewer than 10 patients are suppressed. " ." Zero values in this cell.

Table D.2 (continued)

**Incident ESRD Patients, by Dialysis/Functioning Graft Status, by demographic characteristics & primary diagnosis**

HEMODIALYSIS	1978	1979	1980	1981	1982	1983	1984	1985	1986	1987	1988	1989	1990	1991	1992	1993	1994	1995	1996	1997	1998	1999	2000	2001	
0-4	12	14	19	18	21	19	11	20	21	10	16	28	20	24	14	28	18	36	23	40	29	27	51	68	
5-9	48	45	50	46	41	35	33	30	25	24	21	30	38	29	25	34	44	51	40	32	40	47	54	53	
10-14	158	128	123	120	110	98	91	89	77	77	70	74	89	73	90	79	109	108	102	117	111	130	134	144	
15-19	327	368	352	380	334	312	278	269	269	273	305	321	304	297	297	287	316	345	369	339	382	415	387	403	
20-24	537	535	532	574	610	547	569	601	553	592	601	619	627	673	734	706	747	837	810	765	844	805	818	871	
25-29	694	714	700	774	888	871	886	912	949	917	948	1,088	1,081	1,136	1,162	1,173	1,296	1,258	1,301	1,354	1,401	1,365	1,424	1,385	
30-34	703	839	910	1,017	1,065	1,050	1,140	1,101	1,227	1,144	1,283	1,474	1,475	1,510	1,576	1,618	1,821	1,910	1,888	1,954	1,977	2,028	2,034	1,961	
35-39	801	801	863	982	1,038	1,047	1,131	1,252	1,322	1,410	1,552	1,660	1,740	1,870	2,123	2,080	2,455	2,496	2,627	2,728	2,746	2,875	2,780	2,833	
40-44	877	926	927	1,091	1,042	1,159	1,189	1,281	1,363	1,541	1,662	1,909	2,111	2,304	2,535	2,506	2,815	3,251	3,380	3,464	3,890	3,850	3,961	3,964	
45-49	1,009	1,147	1,086	1,222	1,204	1,344	1,384	1,463	1,561	1,720	1,866	2,166	2,256	2,576	2,847	3,044	3,593	3,871	4,278	4,541	4,950	5,193	5,343	5,455	
50-54	1,334	1,450	1,466	1,638	1,727	1,700	1,701	1,846	1,925	2,038	2,416	2,600	2,700	2,932	3,349	3,499	4,265	4,487	4,715	5,365	5,953	6,613	6,836	7,357	
55-59	1,503	1,665	1,724	2,032	2,097	2,202	2,332	2,439	2,566	2,737	3,057	3,268	3,563	3,930	4,241	4,379	5,203	5,506	5,752	6,320	7,030	7,572	7,940	8,144	
60-64	1,447	1,687	1,878	2,041	2,291	2,661	2,915	3,142	3,298	3,680	4,026	4,498	4,858	5,354	5,621	5,817	6,495	6,968	7,290	7,896	8,513	8,722	9,062	9,415	
65-69	1,443	1,730	1,945	1,873	2,164	2,909	2,927	3,202	3,675	4,217	4,700	5,538	6,149	6,800	7,464	7,615	8,450	8,391	8,955	9,648	10,001	10,174	10,354	10,666	
70-74	1,027	1,273	1,387	1,444	1,784	2,498	2,495	2,894	3,217	3,630	4,028	4,778	5,392	6,296	7,067	7,481	8,423	8,673	9,370	10,191	10,681	10,959	11,311	11,604	
75-79	511	695	810	862	1,026	1,573	1,753	1,898	2,326	2,614	3,008	3,708	4,114	4,714	5,313	5,616	6,421	6,713	7,570	8,552	9,404	10,402	10,651	10,830	
80-84	150	245	287	298	396	658	697	840	1,010	1,249	1,466	1,848	2,140	2,449	2,895	3,054	3,651	3,943	4,516	5,328	5,768	6,361	6,935	7,135	
85+	49	76	83	88	141	223	241	308	382	429	524	729	879	996	1,148	1,329	1,498	1,721	2,053	2,432	2,965	3,411	3,687	3,991	
Unknown	*	.	.	*	*	*	*	*	*	*	*	*	*	*	*	*	20	*	*	*	*	*	*	40	10
0-19	545	555	544	564	506	464	413	408	392	384	412	453	451	423	426	428	487	540	534	528	562	619	626	668	
20-44	3,612	3,815	3,932	4,438	4,643	4,674	4,915	5,147	5,414	5,604	6,046	6,750	7,034	7,493	8,130	8,083	9,134	9,752	10,006	10,265	10,858	10,923	11,017	11,014	
45-64	5,293	5,949	6,154	6,933	7,319	7,907	8,332	8,890	9,350	10,175	11,365	12,532	13,377	14,792	16,058	16,739	19,556	20,832	22,035	24,122	26,446	28,100	29,181	30,371	
65-74	2,470	3,003	3,332	3,317	3,948	5,407	5,422	6,096	6,892	7,847	8,728	10,316	11,541	13,096	14,531	15,096	16,873	17,064	18,325	19,839	20,682	21,133	21,665	22,270	
75+	710	1,016	1,180	1,248	1,563	2,454	2,691	3,046	3,718	4,292	4,998	6,285	7,133	8,159	9,356	9,999	11,570	12,377	14,139	16,312	18,137	20,174	21,273	21,956	
Unknown	*	.	.	*	*	*	*	*	*	*	*	*	*	*	*	*	20	*	*	*	*	*	*	40	10
Male	7,026	8,029	8,524	9,247	9,997	11,523	12,008	12,883	14,142	15,339	17,056	19,682	21,394	23,465	25,665	26,931	30,460	31,862	34,698	37,797	40,823	43,249	44,804	46,448	
Female	5,605	6,309	6,618	7,255	7,985	9,388	9,766	10,709	11,627	12,963	14,495	16,653	18,144	20,500	22,838	23,414	27,173	28,705	30,342	33,271	35,864	37,701	38,962	39,831	
Unknown	*	.	.	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	36	10
White	7,889	8,783	9,189	10,310	11,974	13,561	14,183	15,214	16,650	18,083	19,989	23,102	25,152	27,792	30,503	31,030	35,708	36,534	38,950	43,995	47,196	50,655	52,794	54,609	
Black	4,272	5,007	5,326	5,502	5,403	6,537	6,634	7,294	7,867	8,765	9,925	11,752	12,609	14,125	15,576	16,558	18,610	19,669	21,076	22,272	24,054	24,725	25,038	25,789	
Native American	103	116	160	173	191	253	237	243	306	355	445	530	581	633	772	747	872	1,051	1,203	916	1,325	1,135	1,132	1,105	
Asian	80	112	143	200	330	453	654	763	858	999	1,021	833	1,072	1,187	1,337	1,512	1,881	2,089	2,327	2,346	2,744	2,994	2,931	3,065	
Other/unknown	287	320	324	317	86	110	67	80	88	101	172	121	124	228	315	502	569	1,227	1,485	1,543	1,369	1,441	1,907	1,721	
†Hispanic																			7,922	8,187	9,134	9,701	10,505	10,882	
†Non-Hispanic																			57,119	62,885	67,554	71,249	73,297	75,407	
Diabetes	1,195	1,338	1,890	2,987	3,971	4,714	5,701	6,509	7,531	8,533	9,794	12,031	13,595	15,761	17,747	17,875	21,986	24,800	27,883	31,195	33,905	35,900	37,958	39,302	
Hypertension	1,682	1,909	2,294	3,550	4,664	5,089	5,813	6,526	7,150	8,211	9,300	11,207	12,164	13,691	15,480	14,811	17,936	16,391	17,684	19,514	20,926	22,242	22,831	23,771	
Glomerulonephritis	1,706	1,857	1,869	2,670	3,228	3,367	3,531	3,596	3,706	3,891	4,134	4,478	4,719	4,941	5,129	5,101	5,825	6,065	6,462	6,701	6,906	6,979	6,708	6,630	
Cystic kidney	492	532	574	724	871	891	895	943	990	983	976	984	1,068	1,083	1,188	1,117	1,286	1,340	1,325	1,456	1,595	1,616	1,602	1,596	
Other urologic	191	220	252	338	341	418	410	542	538	554	633	708	735	973	990	875	985	1,134	1,323	1,275	1,377	1,482	1,412	1,412	
Other cause	1,106	1,247	1,371	1,953	2,619	2,885	3,045	3,299	3,346	3,434	3,566	4,030	3,908	4,004	4,368	4,301	5,333	6,784	7,268	7,750	8,446	8,909	9,327	9,439	
Unknown cause	1,087	1,202	1,208	1,314	1,123	1,117	1,132	1,138	1,325	1,577	1,601	1,470	1,567	1,921	2,187	1,912	2,146	2,262	2,586	2,768	3,138	3,197	3,494	3,569	
Missing cause	5,172	6,033	5,684	2,966	1,167	2,433	1,248	1,041	1,183	1,120	1,548	1,430	1,782	1,591	1,414	4,357	2,143	1,794	510	413	395	625	470	570	
All	12,631	14,338	15,142	16,502	17,984	20,914	21,775	23,594	25,769	28,303	31,552	36,338	39,538	43,965	48,503	50,349	57,640	60,570	65,041	71,072	76,688	80,950	83,802	86,289	

The incident cohorts are determined at the time of ESRD initiation without applying the 60-day stable modality rule. The age of the incident patient is determined as of the date of ESRD initiation. †CMS began collecting †Hispanic ethnicity data in April 1995; cells for years prior to 1996 have therefore been left blank. \* Values for cells with fewer than 10 patients are suppressed. . Zero values in this cell.

Table D.2 (continued)

## Incident ESRD Patients, by Dialysis/Functioning Graft Status, by demographic characteristics &amp; primary diagnosis

PERITONEAL DIALYSIS	1978	1979	1980	1981	1982	1983	1984	1985	1986	1987	1988	1989	1990	1991	1992	1993	1994	1995	1996	1997	1998	1999	2000	2001
0-4	*	*	10	26	48	61	69	75	75	78	79	67	67	89	78	74	72	96	113	103	121	112	122	127
5-9	*	13	14	22	38	45	51	41	62	49	41	40	53	46	38	43	61	55	69	56	59	69	65	61
10-14	10	12	13	32	66	73	90	69	65	69	78	56	69	82	74	75	78	95	112	105	101	106	104	97
15-19	18	*	10	29	64	75	79	81	78	97	77	105	98	82	111	98	98	129	127	116	103	103	105	105
20-24	13	21	23	51	78	85	125	84	111	124	135	112	143	132	147	141	147	146	141	131	108	94	95	95
25-29	29	34	39	91	142	165	193	186	217	203	229	234	264	252	228	261	258	274	272	239	238	205	199	171
30-34	42	39	68	120	216	258	233	260	281	280	283	343	353	399	411	402	388	420	404	337	344	314	279	264
35-39	38	34	59	112	194	262	255	300	309	336	377	395	447	497	482	471	495	580	517	449	417	408	367	353
40-44	29	41	62	104	208	250	252	261	305	342	369	405	502	624	583	583	542	664	665	595	509	516	505	480
45-49	37	54	77	110	213	273	281	326	323	328	376	459	479	569	598	647	658	861	788	669	684	652	620	590
50-54	52	65	100	182	275	341	349	341	366	423	392	459	551	594	660	656	734	855	802	834	804	742	736	796
55-59	64	75	125	242	350	402	434	451	430	445	484	567	618	607	714	691	747	884	842	859	811	807	740	837
60-64	65	77	103	232	370	464	478	544	508	542	624	614	737	790	856	792	878	1,016	922	864	792	815	784	767
65-69	36	49	76	208	329	375	421	435	453	528	511	614	758	748	735	852	808	915	844	773	784	732	715	691
70-74	32	39	77	162	285	341	325	337	326	411	419	446	565	589	643	683	745	712	822	719	683	676	627	649
75-79	13	30	38	71	169	205	192	225	213	245	261	333	334	367	428	421	446	477	486	514	476	489	490	493
80-84	*	*	13	40	56	88	90	93	106	107	119	138	175	151	185	204	227	230	229	237	235	241	233	294
85+	*	*	*	*	33	29	34	33	33	45	41	50	51	47	67	70	80	76	96	104	99	116	99	119
Unknown	.	.	.	.	*	*	*	.	*	.	.	*	.	.	*	*	*	*	*	.	*	.	*	*
0-19	37	36	47	109	216	254	289	266	280	293	275	268	287	299	301	290	309	375	421	380	384	390	396	390
20-44	151	169	251	478	838	1,020	1,058	1,091	1,223	1,285	1,393	1,489	1,709	1,904	1,851	1,858	1,830	2,084	1,999	1,751	1,616	1,537	1,445	1,363
45-64	218	271	405	766	1,208	1,480	1,542	1,662	1,627	1,738	1,876	2,099	2,385	2,560	2,828	2,786	3,017	3,616	3,354	3,226	3,091	3,016	2,880	2,990
65-74	68	88	153	370	614	716	746	772	779	939	930	1,060	1,323	1,337	1,378	1,535	1,553	1,627	1,666	1,492	1,467	1,408	1,342	1,340
75+	18	40	56	119	258	322	316	351	352	397	421	521	560	565	680	695	753	783	811	855	810	846	822	906
Unknown	.	.	.	.	*	*	*	.	*	.	.	*	.	.	*	*	*	*	*	.	*	.	*	*
Male	232	296	477	901	1,635	2,028	2,163	2,205	2,332	2,489	2,637	2,924	3,383	3,673	3,778	3,777	3,950	4,492	4,382	4,040	3,837	3,770	3,660	3,660
Female	260	308	435	941	1,502	1,765	1,789	1,937	1,930	2,163	2,258	2,515	2,881	2,992	3,261	3,388	3,514	3,993	3,870	3,664	3,531	3,427	3,225	3,328
Unknown	.	.	.	.	.	*	.	.	.	.	.	.	.	.	*	.	.	*	.	.	*	.	*	*
White	332	433	662	1,312	2,276	2,728	2,839	3,090	3,130	3,389	3,584	3,936	4,651	4,828	5,091	5,086	5,243	5,844	5,715	5,554	5,256	5,246	4,975	4,991
Black	129	146	219	482	801	966	970	902	944	1,085	1,131	1,326	1,424	1,603	1,651	1,743	1,786	2,013	1,831	1,611	1,557	1,490	1,395	1,464
Native American	*	*	*	17	23	33	41	48	59	61	64	69	76	87	93	81	107	139	144	91	115	75	102	90
Asian	11	*	*	17	34	59	95	87	114	108	102	97	100	136	176	219	279	344	354	282	290	265	266	295
Other/unknown	13	15	17	14	*	*	*	15	15	*	14	11	13	11	29	36	49	147	208	166	153	121	149	151
†Hispanic																			888	765	778	786	772	738
†Non-Hispanic																			7,364	6,939	6,593	6,411	6,115	6,253
Diabetes	129	182	247	510	943	1,119	1,216	1,332	1,439	1,570	1,621	1,963	2,293	2,543	2,771	2,646	2,898	3,544	3,676	3,386	3,232	3,170	3,024	2,957
Hypertension	81	99	187	389	780	851	814	885	873	953	1,091	1,227	1,483	1,495	1,574	1,575	1,662	1,642	1,480	1,473	1,429	1,374	1,341	1,403
Glomerulonephritis	115	111	177	314	564	624	689	719	733	778	846	901	990	1,048	1,104	1,029	1,114	1,328	1,350	1,210	1,157	1,043	1,032	1,011
Cystic kidney	27	33	39	91	121	147	177	161	192	195	206	216	276	311	314	336	337	385	362	357	334	365	330	357
Other urologic	*	11	13	43	58	69	73	65	60	65	51	55	76	83	106	105	102	135	137	129	118	120	124	112
Other cause	54	79	129	229	424	561	595	617	574	653	592	635	681	715	692	703	716	922	901	836	835	816	758	846
Unknown cause	82	83	97	198	195	207	219	186	218	209	204	201	229	240	246	250	233	254	286	274	234	240	224	255
Missing cause	*	*	23	68	52	216	169	177	173	229	284	241	236	230	233	521	402	277	60	39	32	69	54	50
All	492	604	912	1,842	3,137	3,794	3,952	4,142	4,262	4,652	4,895	5,439	6,264	6,665	7,040	7,165	7,464	8,487	8,252	7,704	7,371	7,197	6,887	6,991

The incident cohorts are determined at the time of ESRD initiation without applying the 60-day stable modality rule. The age of the incident patient is determined as of the date of ESRD initiation. †CMS began collecting †Hispanic ethnicity data in April 1995; cells for years prior to 1996 have therefore been left blank. \*Values for cells with fewer than 10 patients are suppressed. "Zero values in this cell.

Table D.2 (continued)

**Incident ESRD Patients, by Dialysis/Functioning Graft Status, by demographic characteristics & primary diagnosis**

TRANSPLANT	1978	1979	1980	1981	1982	1983	1984	1985	1986	1987	1988	1989	1990	1991	1992	1993	1994	1995	1996	1997	1998	1999	2000	2001	
0-4	*	10	*	12	24	17	20	22	18	34	37	43	41	38	47	53	33	31	45	45	23	45	36	42	
5-9	14	17	18	24	19	13	17	27	27	24	33	63	62	64	61	61	48	46	43	46	39	56	47	39	
10-14	21	27	25	22	24	30	37	40	54	62	79	93	80	79	73	77	64	74	61	58	64	80	92	79	
15-19	31	39	31	40	44	42	57	53	74	93	114	118	119	92	77	124	112	98	94	87	65	108	108	105	
20-24	41	44	42	37	41	47	57	76	67	94	155	122	129	116	117	128	97	73	77	71	80	88	82	93	
25-29	62	64	67	65	84	78	73	99	119	127	210	237	190	188	175	186	150	127	107	109	110	154	138	144	
30-34	42	43	71	98	78	86	88	104	151	162	261	218	245	247	229	271	216	186	152	145	167	167	213	185	
35-39	38	46	52	38	59	74	81	89	117	175	233	226	238	226	231	249	224	220	187	204	215	230	240	206	
40-44	32	40	40	33	54	53	60	79	88	134	203	206	255	256	248	238	227	195	216	193	233	285	307	281	
45-49	38	35	50	33	54	52	53	58	86	104	167	170	166	190	193	217	216	193	203	198	215	288	330	324	
50-54	27	28	31	24	51	40	46	44	69	96	171	123	134	111	151	197	140	152	159	164	209	262	313	321	
55-59	25	19	*	13	23	17	28	29	44	81	100	131	132	117	115	129	130	102	115	158	155	227	250	271	
60-64	*	*	*	*	*	10	*	12	21	41	57	71	61	67	77	107	80	83	92	102	93	134	157	168	
65-69	*	.	*	*	.	*	*	*	11	11	18	21	27	29	38	47	29	37	38	55	61	62	90	109	
70-74	.	.	.	*	.	*	*	*	*	*	10	11	*	*	10	*	13	*	10	11	30	45	35	.	
75-79	*	*	.	.	.	.	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*
80-84	.	.	.	.	.	.	*	.	*	*	.	.	.	*	*	.	*	.	.	.	*	*	*	*	
85+	*	.	*	*	*	*	.	*	*	.	.	*	*	*	*	.	.	.	.	.	*	.	.	.	
Unknown	*	*	*	*	*	*	*	10	*	11	.	*	*	*	*	*	.	.	.	*	.	.	.	.	
0-19	73	93	80	98	111	102	131	142	173	213	263	317	302	273	258	315	257	249	243	236	191	289	283	265	
20-44	215	237	272	271	316	338	359	447	542	692	1,062	1,009	1,057	1,033	1,000	1,072	914	801	739	722	805	924	980	909	
45-64	93	88	93	76	132	119	133	143	220	322	495	495	493	485	536	650	566	530	569	622	672	911	1,050	1,084	
65-74	*	*	*	*	.	*	13	11	16	15	25	31	38	34	46	57	32	50	43	65	72	92	135	144	
75+	*	.	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	10
Unknown	*	*	*	*	*	*	*	10	*	11	.	*	*	*	*	*	.	.	.	*	.	.	.	.	
Male	261	260	301	305	374	363	398	450	570	728	1,076	1,070	1,068	1,056	1,011	1,193	1,001	913	955	975	989	1,266	1,419	1,409	
Female	125	160	151	145	192	207	244	308	392	527	771	784	826	773	832	907	772	720	641	675	758	952	1,034	1,003	
Unknown	.	.	.	.	.	*	.	.	.	*	.	*	*	*	*	.	.	.	.	.	.	.	.	.	
White	270	306	330	324	386	363	407	510	687	862	1,295	1,399	1,415	1,432	1,403	1,620	1,184	1,136	1,160	1,343	1,445	1,805	2,026	2,022	
Black	63	65	61	65	85	64	72	74	91	153	335	301	283	248	272	319	446	379	331	199	187	276	260	235	
Native American	*	*	*	11	*	*	*	10	*	*	23	13	17	27	19	20	16	18	14	11	12	14	19	14	
Asian	*	*	*	*	*	17	14	17	21	51	77	73	87	72	97	89	96	58	56	53	60	72	84	75	
Other/unknown	46	45	57	43	85	121	142	147	160	182	117	71	95	54	53	54	31	42	35	44	43	51	64	66	
†Hispanic																			141	144	168	177	224	186	
†Non-Hispanic																			1,455	1,506	1,579	2,041	2,229	2,226	
Diabetes	18	28	33	46	88	67	111	121	183	185	202	233	225	262	253	262	280	288	322	305	286	323	299	364	
Hypertension	13	16	22	24	38	31	25	36	42	66	79	69	80	77	95	78	101	72	66	88	85	88	116	126	
Glomerulonephritis	46	49	53	64	75	87	96	131	119	163	162	191	218	186	220	215	211	262	260	256	273	288	321	324	
Cystic kidney	*	10	*	14	11	18	22	26	44	68	68	62	90	103	101	96	86	132	149	134	189	171	217	253	
Other urologic	*	*	*	*	*	*	*	*	17	*	12	21	23	42	32	45	38	42	56	52	50	44	62	60	
Other cause	25	29	57	60	66	65	97	100	145	152	153	156	177	172	180	202	188	240	237	267	229	242	260	338	
Unknown cause	10	10	*	13	23	21	20	31	24	32	32	38	50	47	50	62	48	52	57	57	45	28	85	74	
Missing cause	265	271	276	223	256	274	266	306	388	584	1,139	1,087	1,034	944	913	1,142	821	545	449	491	590	1,034	1,093	873	
All	386	420	452	450	566	572	642	758	962	1,257	1,847	1,857	1,897	1,833	1,844	2,102	1,773	1,633	1,596	1,650	1,747	2,218	2,453	2,412	

The incident cohorts are determined at the time of ESRD initiation without applying the 60-day stable modality rule. The age of the incident patient is determined as of the date of ESRD initiation. †CMS began collecting †Hispanic ethnicity data in April 1995; cells for years prior to 1996 have therefore been left blank. \* Values for cells with fewer than 10 patients are suppressed. " ." Zero values in this cell.







Table D.3 (continued)

**Prevalent ESRD Patients, by Dialysis/Functioning Graft Status, by Demographic Characteristics & Primary Diagnosis**

point prevalent patients on December

<b>PERITONEAL DIALYSIS</b>	<b>1978</b>	<b>1979</b>	<b>1980</b>	<b>1981</b>	<b>1982</b>	<b>1983</b>	<b>1984</b>	<b>1985</b>	<b>1986</b>	<b>1987</b>	<b>1988</b>	<b>1989</b>	<b>1990</b>	<b>1991</b>	<b>1992</b>	<b>1993</b>	<b>1994</b>	<b>1995</b>	<b>1996</b>	<b>1997</b>	<b>1998</b>	<b>1999</b>	<b>2000</b>	<b>2001</b>	
0-4	*	*	12	28	48	72	84	100	105	97	91	101	100	115	130	115	133	147	171	153	171	167	188	184	
5-9	.	*	10	24	48	69	85	73	88	83	74	89	98	79	86	91	104	114	126	136	119	117	139	133	
10-14	*	*	11	32	59	123	148	134	156	151	155	153	162	185	185	189	191	191	214	234	237	207	234	227	
15-19	*	*	12	37	84	133	164	229	215	230	233	284	299	309	328	334	324	335	387	363	342	337	322	301	
20-24	*	12	21	57	90	198	270	280	315	333	347	368	434	475	547	602	600	601	587	560	516	520	475	454	
25-29	16	18	26	120	183	324	432	492	530	579	661	718	808	844	902	936	983	1,020	1,012	987	939	842	827	817	
30-34	12	20	59	174	260	472	652	748	774	802	873	1,004	1,196	1,322	1,438	1,595	1,629	1,631	1,573	1,423	1,352	1,285	1,217	1,197	
35-39	19	17	47	164	289	530	706	906	946	1,078	1,147	1,248	1,482	1,618	1,795	1,934	2,024	2,074	1,993	1,892	1,727	1,738	1,611	1,546	
40-44	14	22	63	180	283	524	695	821	956	1,030	1,134	1,298	1,603	1,902	2,052	2,196	2,323	2,527	2,519	2,424	2,185	2,125	2,020	1,925	
45-49	16	34	65	179	315	579	751	921	935	995	1,127	1,371	1,605	1,889	2,158	2,400	2,632	2,877	2,890	2,732	2,626	2,503	2,454	2,362	
50-54	26	48	85	261	392	677	829	990	1,030	1,131	1,175	1,360	1,690	1,867	2,196	2,377	2,726	2,900	2,991	2,843	2,717	2,776	2,775	2,756	
55-59	29	31	104	318	451	766	1,099	1,276	1,374	1,402	1,400	1,585	1,796	1,982	2,204	2,418	2,634	2,899	2,928	2,941	2,775	2,721	2,682	2,746	
60-64	22	32	63	297	488	824	1,091	1,397	1,494	1,649	1,744	1,904	2,192	2,263	2,616	2,677	2,882	3,146	3,080	2,898	2,619	2,548	2,507	2,437	
65-69	16	31	79	221	394	571	888	1,184	1,404	1,539	1,641	1,879	2,225	2,546	2,815	3,161	3,286	3,347	3,263	2,978	2,707	2,617	2,511	2,486	
70-74	12	19	44	152	327	392	595	819	934	1,054	1,170	1,315	1,577	1,917	2,263	2,599	2,826	2,823	2,753	2,576	2,474	2,317	2,196	2,145	
75-79	12	18	27	68	155	211	314	420	501	604	631	749	933	1,130	1,318	1,434	1,579	1,677	1,681	1,698	1,575	1,509	1,515	1,529	
80-84	*	*	*	35	62	77	104	147	194	241	265	329	396	459	528	605	738	775	747	741	690	685	686	760	
85+	*	*	*	*	31	28	35	48	46	58	71	87	122	140	176	203	213	234	253	271	255	268	250	262	
Unknown	.	.	.	.	*	*	*	*	.	.	.	.	.	.	*	*	*	*	*	.	*	*	*	*	*
0-19	12	25	45	121	239	397	481	536	564	561	553	627	659	688	729	729	752	787	898	886	869	828	883	845	
20-44	68	89	216	695	1,105	2,048	2,755	3,247	3,521	3,822	4,162	4,636	5,523	6,161	6,734	7,263	7,559	7,853	7,684	7,286	6,719	6,510	6,150	5,939	
45-64	93	145	317	1,055	1,646	2,846	3,770	4,584	4,833	5,177	5,446	6,220	7,283	8,001	9,174	9,872	10,874	11,822	11,889	11,414	10,737	10,548	10,418	10,301	
65-74	28	50	123	373	721	963	1,483	2,003	2,338	2,593	2,811	3,194	3,802	4,463	5,078	5,760	6,112	6,170	6,016	5,554	5,181	4,934	4,707	4,631	
75+	16	24	36	112	248	316	453	615	741	903	967	1,165	1,451	1,729	2,022	2,242	2,530	2,686	2,681	2,710	2,520	2,462	2,451	2,551	
Unknown	.	.	.	.	*	*	*	*	.	.	.	.	.	.	*	*	*	*	*	.	*	*	*	*	*
Male	105	172	391	1,214	2,064	3,585	4,819	5,877	6,328	6,878	7,309	8,350	9,811	11,177	12,508	13,591	14,501	15,236	15,014	14,223	13,212	12,793	12,579	12,372	
Female	112	161	346	1,142	1,896	2,986	4,124	5,108	5,669	6,178	6,630	7,492	8,907	9,865	11,229	12,275	13,328	14,084	14,156	13,627	12,814	12,489	12,030	11,894	
Unknown	.	.	.	.	.	*	*	*	.	.	.	.	.	.	*	*	*	*	*	.	*	*	*	*	*
White	152	230	547	1,725	2,909	4,894	6,565	8,105	8,738	9,468	10,057	11,276	13,279	14,793	16,563	17,689	18,776	19,513	19,098	18,236	16,920	16,469	16,021	15,830	
Black	55	89	164	573	950	1,478	2,043	2,466	2,771	3,019	3,313	3,917	4,721	5,439	6,130	6,917	7,532	7,860	7,813	7,352	6,894	6,660	6,420	6,266	
Native American	*	*	*	17	37	68	119	134	164	184	184	221	252	293	353	366	412	456	463	442	442	400	388	390	
Asian	*	*	*	20	46	90	177	231	283	341	334	381	416	471	615	781	957	1,190	1,305	1,253	1,215	1,221	1,242	1,244	
Other/unknown	*	*	10	21	18	42	40	50	41	44	51	47	50	46	77	114	153	303	491	567	558	533	540	538	
†Hispanic																			2,697	2,811	2,816	2,906	2,963	2,914	
†Non-Hispanic																			26,473	25,039	23,213	22,377	21,648	21,354	
Diabetes	55	104	167	497	944	1,322	1,940	2,457	2,778	3,146	3,454	4,213	5,153	6,113	7,218	7,670	8,499	9,376	9,739	9,497	8,900	8,685	8,521	8,464	
Hypertension	37	64	141	410	787	1,162	1,564	2,070	2,387	2,659	2,909	3,408	4,256	4,973	5,645	6,161	6,695	6,731	6,377	6,001	5,612	5,538	5,456	5,324	
Glomerulonephritis	48	69	147	479	771	1,414	1,982	2,413	2,614	2,802	3,051	3,331	3,772	4,097	4,539	4,877	5,219	5,520	5,629	5,441	5,079	4,846	4,722	4,516	
Cystic kidney	12	15	39	133	229	400	536	661	736	756	731	807	975	1,068	1,182	1,262	1,301	1,361	1,359	1,266	1,188	1,157	1,099	1,107	
Other urologic	*	*	13	45	81	111	150	189	194	190	203	222	269	313	392	442	485	493	514	534	505	489	492	487	
Other cause	29	44	90	288	537	948	1,320	1,656	1,774	1,918	1,999	2,276	2,542	2,691	2,897	3,144	3,214	3,452	3,467	3,335	3,207	3,172	3,035	3,130	
Unknown cause	31	30	76	225	323	454	571	623	669	716	731	724	834	915	978	1,060	1,109	1,115	1,127	1,092	1,029	962	931	977	
Missing cause	*	*	64	279	288	761	881	917	845	869	861	861	917	872	887	1,251	1,308	1,274	958	684	509	434	355	263	
All	217	333	737	2,356	3,960	6,572	8,944	10,986	11,997	13,056	13,939	15,842	18,718	21,042	23,738	25,867	27,830	29,322	29,170	27,850	26,029	25,283	24,611	24,268	

The point prevalent cohorts are determined on December 31 of each calendar year without applying the 60-day stable modality rule.

The age of the point prevalent patient is determined as of December 31 of each calendar year. †CMS began collecting Hispanic ethnicity data in April 1995; cells for years prior to 1996 have therefore been left blank.

\* Values for cells with fewer than 10 patients are suppressed. †Zero values in this cell.



Table D.4  
**Incident ESRD Patients for Selected Years, by Age & Modality**

<b>1993</b>	<b>Center hemo</b>	<b>Center self hemo</b>	<b>Home hemo</b>	<b>CAPD</b>	<b>CCPD</b>	<b>Other PD</b>	<b>Uncertain dialysis</b>	<b>Transplant</b>	<b>All</b>
0-4	28	.	.	34	40	*	*	53	170
5-9	34	.	.	25	18	*	*	61	146
10-14	79	.	.	45	30	10	10	77	251
15-19	287	.	.	73	25	*	13	124	528
20-24	704	*	*	126	15	*	20	128	999
25-29	1,169	*	*	235	26	14	57	186	1,691
30-34	1,615	*	*	349	53	24	49	271	2,364
35-39	2,073	*	*	412	59	16	69	249	2,885
40-44	2,502	*	*	522	61	34	93	238	3,454
45-49	3,035	*	*	584	63	26	91	217	4,025
50-54	3,493	*	*	589	67	35	102	197	4,489
55-59	4,373	*	*	622	69	39	114	129	5,352
60-64	5,804	*	*	720	72	58	164	107	6,938
65-69	7,600	*	*	763	89	135	246	47	8,895
70-74	7,462	12	*	595	88	131	228	10	8,533
75-79	5,599	*	10	353	68	80	171	*	6,293
80-84	3,051	.	*	169	35	53	116	.	3,427
85+	1,326	.	*	57	13	14	68	.	1,481
Unknown	*	.	.	*	.	.	*	*	*
0-19	428	.	.	177	113	30	32	315	1,095
20-44	8,063	11	*	1,644	214	92	288	1,072	11,393
45-64	16,705	13	21	2,515	271	158	471	650	20,804
65-74	15,062	19	15	1,358	177	266	474	57	17,428
75+	9,976	*	14	579	116	147	355	*	11,201
Unknown	*	.	.	*	.	.	*	*	*
All	50,238	52	59	6,274	891	693	1,621	2,102	61,930
<b>1997</b>									
0-4	40	.	.	*	95	*	*	45	192
5-9	32	.	.	*	48	*	*	46	138
10-14	116	.	*	16	89	*	*	58	286
15-19	339	.	.	55	61	*	*	87	547
20-24	764	.	*	91	40	*	*	71	970
25-29	1,350	*	*	179	60	10	*	109	1,719
30-34	1,950	*	*	234	103	10	*	145	2,455
35-39	2,721	*	*	347	102	12	16	204	3,409
40-44	3,456	.	*	439	156	19	23	193	4,294
45-49	4,527	*	13	516	153	25	24	198	5,457
50-54	5,336	*	23	619	215	15	39	164	6,417
55-59	6,293	*	24	638	221	25	62	158	7,424
60-64	7,873	*	21	655	209	29	83	102	8,974
65-69	9,608	15	25	603	170	93	260	55	10,829
70-74	10,166	*	20	535	184	90	238	10	11,248
75-79	8,524	*	22	392	122	70	199	*	9,339
80-84	5,305	*	16	167	70	51	109	.	5,725
85+	2,422	*	*	79	25	20	42	.	2,598
Unknown	*	.	.	.	.	.	.	*	*
0-19	527	.	*	87	293	15	*	236	1,163
20-44	10,241	*	18	1,290	461	53	56	722	12,847
45-64	24,029	12	81	2,428	798	94	208	622	28,272
65-74	19,774	20	45	1,138	354	183	498	65	22,077
75+	16,251	15	46	638	217	141	350	*	17,662
Unknown	*	.	.	.	.	.	.	*	*
All	70,828	53	191	5,581	2,123	486	1,116	1,650	82,028

The incident cohorts and associated modalities are determined at the time of ESRD initiation without applying the 60-day stable modality rule.

The age of the incident patient is determined as of the date of ESRD initiation.

\* Values for cells with fewer than 10 patients are suppressed.

." Zero values in this cell.

Table D.4 (continued)

## Incident ESRD Patients for Selected Years, by Age &amp; Modality

2001	Center hemo	Center self hemo	Home hemo	CAPD	CCPD	Other PD	Uncertain dialysis	Transplant	All
0-4	67	.	*	16	111	*	*	42	245
5-9	53	.	.	*	54	*	.	39	154
10-14	144	.	.	12	85	*	*	79	326
15-19	403	.	.	33	72	*	*	105	624
20-24	870	.	*	68	27	*	*	93	1,064
25-29	1,385	.	.	117	54	*	*	144	1,706
30-34	1,956	.	*	184	80	*	11	185	2,424
35-39	2,824	.	*	236	117	*	17	206	3,417
40-44	3,949	*	14	315	165	10	14	281	4,749
45-49	5,440	*	13	397	193	11	21	324	6,401
50-54	7,335	*	16	546	250	15	28	321	8,517
55-59	8,113	*	28	583	254	10	30	271	9,292
60-64	9,382	*	31	541	226	21	49	168	10,420
65-69	10,633	*	26	502	189	51	28	109	11,545
70-74	11,572	*	28	467	182	50	29	35	12,367
75-79	10,791	*	30	379	114	57	20	*	11,408
80-84	7,114	*	19	223	71	24	22	*	7,477
85+	3,981	*	*	84	35	21	15	.	4,146
Unknown	10	.	.	*	*	.	*	.	13
0-19	667	.	*	68	322	15	11	265	1,349
20-44	10,984	*	29	920	443	24	50	909	13,360
45-64	30,270	13	88	2,067	923	57	128	1,084	34,630
65-74	22,205	11	54	969	371	101	57	144	23,912
75+	21,886	13	57	686	220	102	57	10	23,031
Unknown	10	.	.	*	*	.	*	.	13
All	86,022	38	229	4,711	2,280	299	304	2,412	96,295

The incident cohorts and associated modalities are determined at the time of ESRD initiation without applying the 60-day stable modality rule.

The age of the incident patient is determined as of the date of ESRD initiation.

\* Values for cells with fewer than 10 patients are suppressed.

." Zero values in this cell.

Table D.5

**Prevalent ESRD Patients for Selected Years, by Age & Modality***point prevalent patients on December 31*

<b>1993</b>	<b>Center hemo</b>	<b>Center self hemo</b>	<b>Home hemo</b>	<b>CAPD</b>	<b>CCPD</b>	<b>Other PD</b>	<b>Uncertain dialysis</b>	<b>Transplant</b>	<b>All</b>
0-4	36	.	.	39	76	15	20	223	409
5-9	52	.	*	32	59	*	20	619	792
10-14	142	.	*	67	122	15	44	951	1,342
15-19	610	*	*	167	167	20	79	1,504	2,551
20-24	1,877	.	13	453	149	10	106	2,735	5,343
25-29	3,325	*	22	759	177	13	183	4,705	9,186
30-34	5,155	11	28	1,306	289	25	268	7,513	14,595
35-39	6,963	21	65	1,606	328	30	313	8,933	18,259
40-44	8,706	11	53	1,874	322	46	382	9,560	20,954
45-49	9,999	16	67	2,032	368	41	360	8,587	21,470
50-54	11,297	13	63	2,009	368	49	362	7,086	21,247
55-59	13,463	*	61	2,041	377	51	376	5,827	22,204
60-64	16,519	10	75	2,274	403	77	420	4,459	24,237
65-69	20,447	13	79	2,713	448	84	472	2,750	27,006
70-74	19,451	23	65	2,200	399	65	453	973	23,629
75-79	13,854	12	36	1,252	182	37	271	183	15,827
80-84	7,927	*	24	519	86	28	174	26	8,789
85+	3,357	*	14	168	35	13	69	*	3,668
Unknown	*	.	.	*	.	.	*	52	60
0-19	840	*	*	305	424	57	163	3,297	5,094
20-44	26,026	45	181	5,998	1,265	124	1,252	33,446	68,337
45-64	51,278	47	266	8,356	1,516	218	1,518	25,959	89,158
65-74	39,898	36	144	4,913	847	149	925	3,723	50,635
75+	25,138	20	74	1,939	303	78	514	218	28,284
Unknown	*	.	.	*	.	.	*	52	60
All	143,186	149	672	21,512	4,355	626	4,373	66,695	241,568
<b>1997</b>									
0-4	47	.	*	15	138	10	13	212	438
5-9	81	.	*	23	113	*	16	699	936
10-14	168	.	*	27	207	*	28	1,168	1,612
15-19	746	*	14	105	258	*	67	1,892	3,091
20-24	2,249	*	17	303	257	*	93	2,686	5,618
25-29	4,332	19	32	608	379	14	179	4,995	10,558
30-34	6,569	18	58	893	530	22	255	7,916	16,261
35-39	9,371	48	111	1,245	647	35	365	10,804	22,626
40-44	12,464	46	146	1,553	871	40	442	12,054	27,616
45-49	15,531	60	210	1,745	987	47	456	12,288	31,324
50-54	17,350	51	205	1,846	997	51	481	10,882	31,863
55-59	19,504	52	177	1,935	1,006	57	552	8,760	32,043
60-64	22,841	47	168	1,901	997	60	583	6,886	33,483
65-69	26,422	41	194	1,852	1,126	66	782	4,742	35,225
70-74	27,109	53	196	1,637	939	66	835	2,161	32,996
75-79	21,870	24	160	1,064	634	44	636	568	25,000
80-84	12,409	11	87	504	237	19	373	80	13,720
85+	5,796	*	46	183	88	*	200	14	6,344
Unknown	*	.	.	.	.	.	.	44	53
0-19	1,042	*	24	170	716	27	124	3,971	6,077
20-44	34,985	135	364	4,602	2,684	120	1,334	38,455	82,679
45-64	75,226	210	760	7,427	3,987	215	2,072	38,816	128,713
65-74	53,531	94	390	3,489	2,065	132	1,617	6,903	68,221
75+	40,075	43	293	1,751	959	72	1,209	662	45,064
Unknown	*	.	.	.	.	.	.	44	53
All	204,868	485	1,831	17,439	10,411	566	6,356	88,851	330,807

The point prevalent cohorts and associated modalities are determined on December 31 of each calendar year without applying the 60-day stable modality rule.

The age of the point prevalent patient is determined as of December 31 of each calendar year.

\*Values for cells with fewer than 10 patients are suppressed.

." Zero values in this cell.

Table D.5 (continued)  
**Prevalent ESRD Patients for Selected Years, by Age & Modality**  
*point prevalent patients on December 31*

2001	Center hemo	Center self hemo	Home hemo	CAPD	CCPD	Other PD	Uncertain dialysis	Transplant	All
0-4	88	.	*	19	165	12	10	262	557
5-9	94	.	*	16	117	*	*	710	949
10-14	272	.	*	31	196	*	22	1,323	1,859
15-19	831	*	*	55	246	20	63	2,208	3,430
20-24	2,674	*	15	170	284	19	59	3,022	6,244
25-29	4,785	*	14	362	455	13	103	4,703	10,440
30-34	7,474	12	41	569	628	14	121	8,234	17,093
35-39	10,851	20	71	777	769	22	195	11,586	24,291
40-44	15,111	29	100	1,021	904	31	237	14,344	31,777
45-49	19,971	45	145	1,244	1,118	53	254	15,611	38,441
50-54	25,084	42	147	1,435	1,321	55	319	15,828	44,231
55-59	27,097	39	133	1,490	1,256	57	307	12,966	43,345
60-64	29,274	23	106	1,298	1,139	39	297	10,207	42,383
65-69	31,739	39	81	1,176	1,310	32	264	7,305	41,946
70-74	32,041	35	86	1,037	1,108	24	195	3,958	38,484
75-79	28,505	19	86	725	804	23	150	1,263	31,575
80-84	18,146	14	48	368	392	12	107	251	19,338
85+	9,206	*	36	133	129	*	83	44	9,644
Unknown	12	.	.	*	.	.	.	41	54
0-19	1,285	*	13	121	724	46	101	4,503	6,795
20-44	40,895	67	241	2,899	3,040	99	715	41,889	89,845
45-64	101,426	149	531	5,467	4,834	204	1,177	54,612	168,400
65-74	63,780	74	167	2,213	2,418	56	459	11,263	80,430
75+	55,857	41	170	1,226	1,325	40	340	1,558	60,557
Unknown	12	.	.	*	.	.	.	41	54
All	263,255	333	1,122	11,927	12,341	445	2,792	113,866	406,081

The point prevalent cohorts and associated modalities are determined on December 31 of each calendar year without applying the 60-day stable modality rule. The age of the point prevalent patient is determined as of December 31 of each calendar year.  
 \*Values for cells with fewer than 10 patients are suppressed.  
 ." Zero values in this cell.



Table D.6  
**Incident ESRD Patients for Selected Years**  
*by gender, race, ethnicity, & primary diagnosis; grouped by modality*

<b>1993†</b>	<b>Center hemo</b>	<b>Center self hemo</b>	<b>Home hemo</b>	<b>CAPD</b>	<b>CCPD</b>	<b>Other PD</b>	<b>Uncertain dialysis</b>	<b>Transplant</b>	<b>All</b>
Male	26,873	31	27	3,291	486	365	782	1,193	33,048
Female	23,361	21	32	2,983	405	328	836	907	28,873
Unknown	*	.	.	.	.	.	*	*	*
White	30,954	42	34	4,436	650	482	937	1,620	39,155
Black	16,532	*	20	1,530	213	185	327	319	19,132
Native American	744	*	*	79	*	16	15	20	879
Asian	1,509	*	.	196	23	*	52	89	1,878
Other/unknown	499	.	*	33	*	*	290	54	886
Diabetes	17,841	17	17	2,362	284	254	375	262	21,412
Hypertension	14,786	13	12	1,393	182	191	345	78	17,000
Glomerulonephritis	5,089	*	*	895	134	64	111	215	6,520
Cystic kidney	1,112	*	*	289	47	12	15	96	1,576
Other urologic	872	*	*	86	19	*	21	45	1,053
Other cause	4,291	*	*	580	123	66	96	202	5,368
Unknown cause	1,906	*	*	217	33	35	36	62	2,295
Missing cause	4,341	*	13	452	69	64	622	1,142	6,706
All	50,238	52	59	6,274	891	693	1,621	2,102	61,930
<b>1997</b>									
Male	37,663	27	107	2,913	1,127	267	574	975	43,653
Female	33,161	26	84	2,668	996	219	542	675	38,371
Unknown	*	.	.	.	.	.	.	.	*
White	43,826	37	132	4,086	1,468	307	817	1,343	52,016
Black	22,210	*	53	1,100	511	147	246	199	24,475
Native American	911	*	*	69	22	12	*	11	1,036
Asian	2,340	*	*	217	65	11	25	53	2,717
Other/unknown	1,541	*	*	109	57	*	22	44	1,784
Hispanic	8,169	*	15	523	242	28	69	144	9,193
Non-Hispanic	62,659	50	176	5,058	1,881	458	1,047	1,506	72,835
Diabetes	31,099	29	67	2,547	839	216	465	305	35,567
Hypertension	19,455	*	56	1,084	389	121	312	88	21,508
Glomerulonephritis	6,675	*	18	864	346	41	45	256	8,253
Cystic kidney	1,445	*	*	255	102	13	17	134	1,977
Other urologic	1,271	.	*	94	35	13	40	52	1,509
Other cause	7,722	*	24	522	314	57	92	267	9,002
Unknown cause	2,751	*	11	193	81	22	44	57	3,165
Missing cause	410	*	*	22	17	*	101	491	1,047
All	70,828	53	191	5,581	2,123	486	1,116	1,650	82,028

The incident cohorts and associated modalities are determined at the time of ESRD initiation without applying the 60-day stable modality rule.

†CMS began collecting Hispanic ethnicity data in April 1995; cells for years prior to 1996 have therefore been left blank.

\*Values for cells with fewer than 10 patients are suppressed.

." Zero values in this cell.

On our website: Incident Patients by Age, Gender, Race, Ethnicity, & Primary Diagnosis, by Age & ESRD Network of Residence, & Point Prevalent Patients on Dec. 31 by Age & ESRD Network (D.a-d).

Table D.6 (continued)

**Incident ESRD Patients for Selected Years***by gender, race, ethnicity, & primary diagnosis; grouped by modality*

2001	Center hemo	Center self hemo	Home hemo	CAPD	CCPD	Other PD	Uncertain dialysis	Transplant	All
Male	46,308	17	123	2,462	1,198	155	156	1,409	51,828
Female	39,704	21	106	2,247	1,081	144	147	1,003	44,453
Unknown	10	.	.	*	*	.	*	.	14
White	54,445	28	136	3,396	1,595	196	171	2,022	61,989
Black	25,697	*	85	949	515	86	91	235	27,665
Native American	1,104	.	*	62	28	10	*	14	1,220
Asian	3,058	*	*	228	67	*	*	75	3,448
Other/unknown	1,718	.	*	76	75	*	33	66	1,973
Hispanic	10,862	.	20	462	276	19	25	186	11,850
Non-Hispanic	75,160	38	209	4,249	2,004	280	279	2,226	84,445
Diabetes	39,201	14	87	2,069	888	131	59	364	42,813
Hypertension	23,681	*	81	1,045	358	92	27	126	25,419
Glomerulonephritis	6,610	*	13	645	366	19	*	324	7,986
Cystic kidney	1,590	*	*	225	132	*	*	253	2,211
Other urologic	1,410	.	*	74	38	*	*	60	1,590
Other cause	9,415	*	21	460	386	38	18	338	10,679
Unknown cause	3,547	*	19	167	88	13	*	74	3,920
Missing cause	568	*	*	26	24	*	183	873	1,677
All	86,022	38	229	4,711	2,280	299	304	2,412	96,295

The incident cohorts and associated modalities are determined at the time of ESRD initiation without applying the 60-day stable modality rule.

\* Values for cells with fewer than 10 patients are suppressed.

." Zero values in this cell.

On our website: Incident Patients by Age, Gender, Race, Ethnicity, & Primary Diagnosis, by Age & ESRD Network of Residence, & Point Prevalent Patients on Dec. 31 by Age & ESRD Network (D.a-d).

Table D.7

**Prevalent ESRD Patients for Selected Years***point prevalent patients on December 31, by gender, race, ethnicity, & primary diagnosis; grouped by modality*

	Center hemo	Center self hemo	Home hemo	CAPD	CCPD	Other PD	Uncertain dialysis	Transplant	All
<b>1993†</b>									
Male	74,209	87	394	11,332	2,259	323	2,168	39,783	130,555
Female	68,971	62	278	10,179	2,096	303	2,198	26,897	110,984
Unknown	*	.	.	*	.	.	*	15	29
White	76,805	95	461	14,671	3,018	406	2,474	51,021	148,951
Black	58,141	47	184	5,745	1,172	190	1,048	10,875	77,402
Native American	2,322	*	11	330	36	14	48	821	3,583
Asian	4,622	*	12	669	112	*	160	2,442	8,031
Other/unknown	1,296	.	*	97	17	*	643	1,536	3,601
Diabetes	44,561	31	127	6,491	1,179	199	1,035	11,980	65,603
Hypertension	42,829	39	143	5,214	947	139	774	7,953	58,038
Glomerulonephritis	18,951	37	155	4,028	849	79	552	17,354	42,005
Cystic kidney	4,827	*	47	1,009	253	23	129	4,781	11,074
Other urologic	2,773	.	13	378	64	16	68	1,150	4,462
Other cause	13,501	22	86	2,499	645	88	381	10,171	27,393
Unknown cause	6,184	*	31	867	193	19	159	2,818	10,275
Missing cause	9,560	11	70	1,026	225	63	1,275	10,488	22,718
All	143,186	149	672	21,512	4,355	626	4,373	66,695	241,568
<b>1997</b>									
Male	107,242	292	1,007	8,671	5,552	280	3,115	52,915	179,074
Female	97,621	193	824	8,768	4,859	286	3,237	35,924	151,712
Unknown	*	.	.	.	.	.	*	12	21
White	107,579	261	1,121	11,368	6,868	318	3,743	66,494	197,752
Black	82,184	195	575	4,617	2,735	194	1,410	15,987	107,897
Native American	3,625	12	69	302	140	10	50	1,109	5,317
Asian	7,535	12	50	788	465	19	193	3,601	12,663
Other/unknown	3,945	*	16	364	203	25	960	1,660	7,178
Hispanic	21,339	26	127	1,634	1,177	54	572	8,517	33,446
Non-Hispanic	183,529	459	1,704	15,805	9,234	512	5,784	80,334	297,361
Diabetes	77,565	123	485	6,096	3,401	205	1,868	17,526	107,269
Hypertension	57,805	106	388	3,815	2,186	126	1,212	11,387	77,025
Glomerulonephritis	25,282	101	346	3,402	2,039	90	605	22,435	54,300
Cystic kidney	5,815	35	116	788	478	11	134	6,851	14,228
Other urologic	3,974	*	51	329	205	*	130	1,731	6,432
Other cause	20,048	64	244	1,905	1,430	70	537	13,134	37,432
Unknown cause	7,857	21	98	681	411	20	210	3,598	12,896
Missing cause	6,522	29	103	423	261	38	1,660	12,189	21,225
All	204,868	485	1,831	17,439	10,411	566	6,356	88,851	330,807

The point prevalent cohorts and associated modalities are determined on December 31 of each calendar year without applying the 60-day stable modality rule.

†CMS began collecting Hispanic ethnicity data in April 1995; cells for years prior to 1996 have therefore been left blank.

\* Values for cells with fewer than 10 patients are suppressed.

." Zero values in this cell.

Table D.7 (continued)

**Prevalent ESRD Patients for Selected Years***point prevalent patients on December 31, by gender, race, ethnicity, & primary diagnosis; grouped by modality*

2001	Center hemo	Center self hemo	Home hemo	CAPD	CCPD	Other PD	Uncertain dialysis	Transplant	All
Male	140,345	191	661	5,887	6,485	219	1,524	67,487	222,799
Female	122,898	142	461	6,038	5,856	226	1,268	46,368	183,257
Unknown	12	.	.	*	.	.	.	11	25
White	141,644	140	716	7,730	8,100	238	1,593	84,261	244,422
Black	101,146	176	313	3,035	3,231	173	858	20,753	129,685
Native American	4,602	*	23	209	181	12	37	1,397	6,466
Asian	10,487	10	55	676	568	14	122	5,040	16,972
Other/unknown	5,376	*	15	277	261	*	182	2,415	8,536
Hispanic	34,075	11	68	1,340	1,574	36	328	12,549	49,981
Non-Hispanic	229,180	322	1,054	10,587	10,767	409	2,464	101,317	356,100
Diabetes	109,495	95	270	4,319	4,145	170	844	23,625	142,963
Hypertension	72,779	102	237	2,675	2,649	68	498	14,600	93,608
Glomerulonephritis	29,541	54	235	2,141	2,375	64	366	28,132	62,908
Cystic kidney	6,919	14	69	579	528	16	83	9,281	17,489
Other urologic	4,852	*	29	238	249	*	48	2,484	7,913
Other cause	25,964	33	176	1,353	1,777	85	274	16,415	46,077
Unknown cause	9,954	14	58	488	489	12	73	4,787	15,875
Missing cause	3,751	12	48	134	129	26	606	14,542	19,248
All	263,255	333	1,122	11,927	12,341	445	2,792	113,866	406,081

The point prevalent cohorts and associated modalities are determined on December 31 of each calendar year without applying the 60-day stable modality rule.

\* Values for cells with fewer than 10 patients are suppressed.

." Zero values in this cell.

Table D.8

**Treatment Status Two Years after First ESRD Service***incident patients, 1997-1999 combined; by gender, age at first service, & modality at 90 days*

ALL	Treatment status two years after first ESRD service (percent of patients)						Patient count
	Center hemo	Home hemo	CAPD & CCPD	Oth/unk dialysis	Transplant	Death	
Age at first service							
0-19	31.0	0.2	15.2	1.6	44.9	7.0	1,315
20-44	58.6	0.2	9.6	0.7	15.8	15.1	19,397
45-64	60.2	0.2	6.4	0.9	5.3	27.2	45,878
65-74	51.8	0.2	4.6	0.9	1.5	41.0	46,746
75+	42.3	0.2	2.1	0.7	0.1	54.6	39,111
Modality on day 90							
Center hemodialysis	57.5	0.1	1.6	0.4	3.1	37.3	131,421
Home hemodialysis	26.6	15.4	4.4	0.8	3.2	49.7	527
CAPD/CCPD	18.9	0.2	38.1	0.6	8.5	33.7	14,953
Other/unknown dialysis	39.9	0.2	4.4	14.6	1.6	39.4	4,082
Transplant	1.8	.	0.3	0.8	94.0	3.0	1,464
Interaction of age at first service & modality on day 90							
0-19	Center hemodialysis						
	51.6	.	5.7	0.7	34.1	7.9	684
	Home hemodialysis						
	.	.	40.0	.	40.0	20.0	*
	CAPD/CCPD						
	10.7	0.7	35.7	1.9	43.2	7.7	428
	Other/unknown dialysis						
	15.2	.	15.2	24.2	36.4	9.1	33
	Transplant						
	2.4	.	0.6	.	96.4	0.6	165
20-44	Center hemodialysis						
	67.8	0.1	3.7	0.5	11.9	16.1	15,739
	Home hemodialysis						
	31.6	26.3	10.5	.	15.8	15.8	38
	CAPD/CCPD						
	21.4	0.3	45.3	0.8	20.7	11.5	2,775
	Other/unknown dialysis						
	37.5	.	9.7	16.7	10.2	25.9	216
	Transplant						
	1.9	.	0.3	1.1	94.6	2.1	629
45-64	Center hemodialysis						
	66.2	0.1	2.0	0.6	3.9	27.2	39,502
	Home hemodialysis						
	35.8	19.0	8.0	1.5	5.1	30.7	137
	CAPD/CCPD						
	20.9	0.2	42.0	0.8	8.0	28.1	4,958
	Other/unknown dialysis						
	46.4	0.3	4.2	14.8	2.4	31.9	788
	Transplant						
	1.6	.	0.4	0.8	92.5	4.7	493
65-74	Center hemodialysis						
	56.2	0.1	1.2	0.3	1.1	41.0	40,388
	Home hemodialysis						
	24.4	14.7	3.2	1.3	1.3	55.1	156
	CAPD/CCPD						
	18.2	0.2	35.6	0.4	2.4	43.1	4,393
	Other/unknown dialysis						
	40.5	0.3	5.1	16.2	0.7	37.2	1,640
	Transplant						
	1.8	.	.	0.6	93.5	4.1	169
75+	Center hemodialysis						
	44.6	0.1	0.4	0.2	0.1	54.5	35,108
	Home hemodialysis						
	21.5	11.5	0.5	.	.	66.5	191
	CAPD/CCPD						
	14.5	0.0	26.5	0.1	0.2	58.6	2,399
	Other/unknown dialysis						
	36.3	0.1	2.5	12.2	.	49.0	1,405
	Transplant						
	.	.	.	.	100.0	.	*
All	52.6	0.2	5.2	0.8	4.5	36.7	152,447

The incident cohorts and associated modalities are determined at the time of ESRD initiation without applying the 60-day stable modality rule.

The age of the incident patient is determined as of the date of ESRD initiation.

Includes only incident patients with known age & patients who have Medicare as their primary payor, are receiving their first ESRD service, & survive the first 90 days.

\* Values for cells with fewer than 10 patients are suppressed.

." Zero values in this cell.

On our website: 1997-1999 Combined, Incident Pediatric Patients by Gender & Age at First Service, & Modality at 90 days (D.e) & by Age, Gender, Race, & Diabetic Status (D.f).

Table D.8 (continued)

**Treatment Status Two Years after First ESRD Service**

incident patients, 1997-1999 combined; by gender, age at first service, &amp; modality at 90 days

MALE	Treatment status two years after first ESRD service (percent of patients)							Patient count
	Center hemo	Home hemo	CAPD & CCPD	Oth/unk dialysis	Transplant	Death		
Age at first service								
0-19	31.1	0.3	11.5	2.1	48.3	6.8	679	
20-44	60.2	0.2	7.8	0.6	15.5	15.7	11,777	
45-64	59.7	0.2	6.1	0.9	6.0	27.1	24,393	
65-74	50.1	0.2	5.3	1.0	2.0	41.4	22,788	
75+	42.2	0.2	2.4	0.7	0.1	54.4	19,848	
Modality on day 90								
Center hemodialysis	57.5	0.1	1.6	0.4	3.7	36.7	68,352	
Home hemodialysis	25.4	14.9	4.1	0.7	4.9	50.0	268	
CAPD/CCPD	18.7	0.2	37.8	0.6	8.9	33.8	7,922	
Other/unknown dialysis	38.8	0.2	4.5	15.0	2.0	39.5	2,130	
Transplant	1.2	.	0.4	0.6	94.8	3.0	813	
Interaction of age at first service & modality on day 90								
0-19	Center hemodialysis	50.1	.	5.0	1.1	36.0	7.8	361
	Home hemodialysis	.	.	50.0	.	50.0	.	*
	CAPD/CCPD	12.2	1.0	25.9	2.4	51.2	7.3	205
	Other/unknown dialysis	13.0	.	21.7	21.7	34.8	8.7	23
	Transplant	2.3	.	1.1	.	95.5	1.1	88
20-44	Center hemodialysis	68.5	0.1	2.8	0.4	11.8	16.4	9,805
	Home hemodialysis	27.3	22.7	13.6	.	18.2	18.2	22
	CAPD/CCPD	22.1	0.2	42.9	0.7	21.1	13.0	1,482
	Other/unknown dialysis	37.0	.	6.7	15.1	10.1	31.1	119
	Transplant	1.4	.	0.6	0.6	94.8	2.6	349
45-64	Center hemodialysis	65.6	0.1	2.0	0.6	4.5	27.1	21,087
	Home hemodialysis	33.3	16.7	6.0	1.2	7.1	35.7	84
	CAPD/CCPD	20.3	0.3	41.5	0.9	8.6	28.4	2,548
	Other/unknown dialysis	44.6	0.5	3.8	15.8	3.5	31.8	399
	Transplant	0.7	.	.	0.7	94.9	3.6	275
65-74	Center hemodialysis	55.0	0.1	1.5	0.3	1.5	41.6	19,397
	Home hemodialysis	20.5	16.4	2.7	1.4	2.7	56.2	73
	CAPD/CCPD	17.1	0.2	36.9	0.5	2.8	42.6	2,390
	Other/unknown dialysis	39.3	0.4	5.6	17.4	1.0	36.3	832
	Transplant	1.0	.	.	1.0	93.8	4.2	96
75+	Center hemodialysis	44.5	0.1	0.5	0.3	0.1	54.5	17,702
	Home hemodialysis	21.8	10.3	.	.	.	67.8	87
	CAPD/CCPD	15.7	.	28.1	.	0.3	56.0	1,297
	Other/unknown dialysis	36.3	.	2.8	11.6	.	49.3	757
	Transplant	.	.	.	.	100.0	.	*
All		52.4	0.2	5.3	0.8	5.1	36.2	79,485

The incident cohorts and associated modalities are determined at the time of ESRD initiation without applying the 60-day stable modality rule.

The age of the incident patient is determined as of the date of ESRD initiation.

Includes only incident patients with known age & patients who have Medicare as their primary payor, are receiving their first ESRD service, & survive the first 90 days.

\* Values for cells with fewer than 10 patients are suppressed.

." Zero values in this cell.

On our website: 1997-1999 Combined, Incident Pediatric Patients by Gender & Age at First Service, & Modality at 90 days (B.e) & by Age, Gender, Race, & Diabetic Status (B.f).

Table D.8 (continued)

**Treatment Status Two Years after First ESRD Service***incident patients, 1997-1999 combined; by gender, age at first service, & modality at 90 days*

FEMALE	Treatment status two years after first ESRD service (percent of patients)						Patient count	
	Center hemo	Home hemo	CAPD & CCPD	Oth/unk dialysis	Transplant	Death		
Age at first service								
0-19	31.0	0.2	19.2	1.1	41.4	7.2	636	
20-44	56.1	0.2	12.3	0.9	16.3	14.1	7,620	
45-64	60.7	0.1	6.6	0.8	4.5	27.2	21,485	
65-74	53.4	0.2	3.9	0.8	1.1	40.6	23,958	
75+	42.5	0.2	1.8	0.6	0.0	54.8	19,263	
Modality on day 90								
Center hemodialysis	57.5	0.1	1.5	0.4	2.5	38.0	63,069	
Home hemodialysis	27.8	15.8	4.6	0.8	1.5	49.4	259	
CAPD/CCPD	19.2	0.2	38.4	0.6	7.9	33.7	7,031	
Other/unknown dialysis	41.0	0.2	4.2	14.2	1.1	39.3	1,952	
Transplant	2.6	.	0.3	1.1	92.9	3.1	651	
Interaction of age at first service & modality on day 90								
0-19	Center hemodialysis	53.3	.	6.5	0.3	31.9	8.0	323
	Home hemodialysis	.	.	33.3	.	33.3	33.3	*
	CAPD/CCPD	9.4	0.4	44.8	1.3	35.9	8.1	223
	Other/unknown dialysis	20.0	.	.	30.0	40.0	10.0	10
	Transplant	2.6	.	.	.	97.4	.	77
20-44	Center hemodialysis	66.8	0.1	5.1	0.5	11.9	15.6	5,934
	Home hemodialysis	37.5	31.3	6.3	.	12.5	12.5	16
	CAPD/CCPD	20.6	0.3	48.0	0.9	20.3	9.7	1,293
	Other/unknown dialysis	38.1	.	13.4	18.6	10.3	19.6	97
	Transplant	2.5	.	.	1.8	94.3	1.4	280
45-64	Center hemodialysis	66.8	0.1	2.0	0.6	3.2	27.3	18,415
	Home hemodialysis	39.6	22.6	11.3	1.9	1.9	22.6	53
	CAPD/CCPD	21.7	0.1	42.4	0.7	7.3	27.8	2,410
	Other/unknown dialysis	48.3	.	4.6	13.9	1.3	31.9	389
	Transplant	2.8	.	0.9	0.9	89.4	6.0	218
65-74	Center hemodialysis	57.4	0.2	1.0	0.3	0.7	40.4	20,991
	Home hemodialysis	27.7	13.3	3.6	1.2	.	54.2	83
	CAPD/CCPD	19.6	0.2	34.2	0.3	1.9	43.7	2,003
	Other/unknown dialysis	41.8	0.2	4.6	14.9	0.4	38.1	808
	Transplant	2.7	.	.	.	93.2	4.1	73
75+	Center hemodialysis	44.7	0.1	0.4	0.2	0.0	54.6	17,406
	Home hemodialysis	21.2	12.5	1.0	.	.	65.4	104
	CAPD/CCPD	13.2	0.1	24.7	0.3	0.1	61.6	1,102
	Other/unknown dialysis	36.3	0.2	2.2	12.8	.	48.6	648
	Transplant	.	.	.	.	100.0	.	*
All		52.8	0.2	5.2	0.8	3.8	37.4	72,962

The incident cohorts and associated modalities are determined at the time of ESRD initiation without applying the 60-day stable modality rule.

The age of the incident patient is determined as of the date of ESRD initiation.

Includes only incident patients with known age & patients who have Medicare as their primary payor, are receiving their first ESRD service, & survive the first 90 days.

\* Values for cells with fewer than 10 patients are suppressed.

." Zero values in this cell.

On our website: 1997-1999 Combined, Incident Pediatric Patients by Gender & Age at First Service, & Modality at 90 days (B.e) & by Age, Gender, Race, & Diabetic Status (B.f).

Table D.9

**Prevalent ESRD Patients for Selected Years, by Patient Vintage***point prevalent patients on December 31, by year-end treatment modality & number of years patient has had ESRD (vintage)*

1993	Dialysis	Functioning graft	Center hemo	Center self hemo	Home hemo	CAPD	CCPD	Other PD	Uncertain dialysis	All
0	51,916	3,132	41,674	32	99	7,108	1,276	217	1,510	55,048
1	36,115	5,255	28,747	24	118	5,019	1,107	167	933	41,370
2	24,319	6,453	19,737	13	73	3,167	635	86	608	30,772
3	15,941	6,743	13,362	14	52	1,781	407	47	278	22,684
4	11,292	6,646	9,677	*	39	1,138	218	19	192	17,938
5	7,771	5,838	6,629	*	22	784	152	26	152	13,609
6	5,556	4,977	4,717	*	38	538	134	17	109	10,533
7	4,195	4,408	3,560	*	30	418	90	14	82	8,603
8	3,352	3,804	2,857	11	23	334	54	*	65	7,156
9	2,654	3,333	2,257	*	19	241	63	*	59	5,987
10	2,166	2,906	1,846	*	14	199	46	*	50	5,072
11	1,849	2,706	1,561	*	13	178	41	*	50	4,555
12	1,504	2,249	1,316	*	21	110	20	*	33	3,753
13	1,197	1,719	1,021	*	17	98	25	*	31	2,916
14	1,090	1,478	939	*	19	85	19	*	23	2,568
15	904	1,318	783	*	11	67	13	*	24	2,222
16	1,405	1,527	1,236	*	35	87	28	*	16	2,932
17	409	537	342	*	*	32	*	.	20	946
18	415	482	310	*	10	38	*	*	49	897
19	332	428	260	*	*	24	*	.	38	760
20	280	384	191	*	*	29	*	*	46	664
21	35	116	29	.	.	*	*	.	*	151
22	26	98	22	.	.	*	.	.	.	124
23	18	60	14	.	.	*	.	.	.	78
24	95	42	76	.	*	16	*	.	*	137
25	*	27	*	.	.	*	.	.	*	34
26	*	*	*	.	*	*	.	.	*	13
27	*	*	.	.	.	*	.	.	.	*
28	21	16	14	.	.	*	*	.	.	37
All	174,873	66,695	143,186	149	672	21,512	4,355	626	4,373	241,568
<b>1997</b>										
0	68,702	2,844	58,508	86	250	5,376	2,849	147	1,486	71,546
1	48,724	5,240	40,555	66	402	3,809	2,344	140	1,408	53,964
2	34,651	7,161	29,108	67	209	2,583	1,827	78	779	41,812
3	24,880	8,055	20,554	47	155	1,675	1,016	66	1,367	32,935
4	15,851	8,066	13,742	28	115	1,043	601	23	299	23,917
5	11,363	7,786	9,923	28	76	725	402	29	180	19,149
6	8,353	6,836	7,249	18	73	507	313	18	175	15,189
7	6,014	6,259	5,268	17	49	333	217	17	113	12,273
8	4,562	5,678	3,987	13	38	267	146	12	99	10,240
9	3,562	4,779	3,082	12	41	210	125	*	84	8,341
10	2,581	4,043	2,210	11	31	154	106	10	59	6,624
11	2,144	3,509	1,835	13	36	143	76	*	35	5,653
12	1,851	2,999	1,600	*	36	107	70	*	27	4,850
13	1,546	2,613	1,320	*	32	93	64	*	25	4,159
14	1,295	2,298	1,100	*	24	93	48	*	23	3,593
15	1,154	2,117	959	*	38	67	45	*	35	3,271
16	899	1,809	762	*	32	44	26	.	27	2,708
17	755	1,416	625	*	22	55	27	*	16	2,171
18	694	1,225	582	*	36	34	30	.	10	1,919
19	553	1,103	449	*	34	28	21	.	14	1,656
20	791	1,240	673	*	51	28	21	*	10	2,031
21	247	437	206	*	*	14	*	*	12	684
22	273	389	200	*	13	20	11	*	24	662
23	209	340	149	*	17	*	10	.	17	549
24	177	311	124	*	*	12	*	.	27	488
25	33	92	24	.	*	*	*	.	*	125
26	25	71	17	.	*	*	*	.	*	96
27	13	42	12	.	.	.	.	.	*	55
28	35	48	31	.	.	.	*	.	.	83
29	*	18	*	.	.	.	.	.	.	26
30	*	*	*	.	.	*	.	.	.	*
31	*	*	*	.	.	*	.	.	.	10
32	*	11	*	.	.	*	.	.	.	18
All	241,956	88,851	204,868	485	1,831	17,439	10,411	566	6,356	330,807

The point prevalent cohorts &amp; associated modalities are determined on December 31 of each calendar year without applying the 60-day stable modality rule.

\* Values for cells with fewer than 10 patients are suppressed.

." Zero values in this cell.



Table D.9 (continued)

**Prevalent ESRD Patients for Selected Years, by Patient Vintage***point prevalent patients on December 31, by year-end treatment modality & number of years patient has had ESRD (vintage)*

2001	Dialysis	Functioning graft	Center hemo	Center self hemo	Home hemo	CAPD	CCPD	Other PD	Uncertain dialysis	All
0	79,555	3,501	71,317	53	179	4,037	3,290	97	582	83,056
1	58,133	6,255	51,973	43	136	2,625	2,680	113	563	64,388
2	43,226	7,808	38,878	36	130	1,733	1,897	90	462	51,034
3	29,526	8,677	26,922	32	85	990	1,307	23	167	38,203
4	19,296	8,967	17,607	32	70	584	837	24	142	28,263
5	14,121	8,938	12,945	30	63	429	528	*	117	23,059
6	10,498	8,695	9,595	16	59	288	396	14	130	19,193
7	8,021	8,218	7,268	*	35	257	297	11	147	16,239
8	5,823	7,261	5,291	*	37	179	198	*	101	13,084
9	4,501	6,527	4,112	11	28	127	155	10	58	11,028
10	3,461	5,589	3,110	*	27	115	120	16	70	9,050
11	2,856	4,999	2,594	*	20	82	112	*	37	7,855
12	2,281	4,377	2,065	*	14	78	82	*	32	6,658
13	1,984	3,690	1,748	*	17	68	95	*	45	5,674
14	1,471	3,048	1,285	*	20	61	63	*	30	4,519
15	1,245	2,679	1,104	*	14	50	51	*	18	3,924
16	1,052	2,282	933	*	18	45	39	*	11	3,334
17	908	2,064	794	*	20	27	44	*	16	2,972
18	782	1,812	698	*	14	27	25	*	12	2,594
19	668	1,679	588	*	20	24	23	.	*	2,347
20	582	1,403	511	*	19	13	27	.	10	1,985
21	465	1,138	408	*	16	21	11	*	*	1,603
22	412	962	358	.	20	*	20	.	*	1,374
23	331	909	280	*	19	17	10	.	*	1,240
24	458	985	410	*	19	16	*	.	*	1,443
25	134	368	123	*	*	*	*	.	.	502
26	152	304	125	*	*	*	*	.	*	456
27	112	274	92	*	*	*	*	.	*	386
28	94	237	71	.	*	*	*	.	*	331
29	23	68	16	.	*	*	*	.	*	91
30	16	51	13	.	*	.	*	.	.	67
31	*	33	*	.	.	.	.	.	.	37
32	12	42	10	.	.	*	*	.	.	54
33	*	12	*	.	.	*	*	.	*	18
34	*	*	*	.	.	*	.	.	.	*
35	*	*	*	.	.	.	.	.	.	*
36	*	*	.	.	.	.	*	.	.	*
All	292,215	113,866	263,255	333	1,122	11,927	12,341	445	2,792	406,081

The point prevalent cohorts &amp; associated modalities are determined on December 31 of each calendar year without applying the 60-day stable modality rule.

\* Values for cells with fewer than 10 patients are suppressed.

"." Zero values in this cell.

Table D.10

## Prevalent ESRD Patients for Selected Years, by Number of Years on Modality

point prevalent patients on December 31; by year-end treatment modality

1993	Dialysis	Functioning graft	Center hemo	Center self hemo	Home hemo	CAPD	CCPD	Other PD	Uncertain dialysis	All
0	157,525	10,049	129,303	113	597	19,761	4,144	534	3,073	167,574
1	9,343	8,541	7,435	20	34	1,092	156	46	560	17,884
2	3,730	7,960	3,000	*	14	319	31	11	351	11,690
3	1,366	6,971	1,140	*	*	107	12	14	84	8,337
4	834	5,919	700	*	*	73	*	*	51	6,753
5	488	5,146	396	*	*	34	*	*	45	5,634
6	361	4,416	293	*	*	34	*	*	20	4,777
7	255	3,917	217	*	*	16	*	*	15	4,172
8	220	2,962	167	*	*	39	.	*	10	3,182
9	148	2,375	123	.	*	16	.	*	*	2,523
10	90	1,955	73	*	.	13	*	*	*	2,045
11	70	1,557	57	*	.	*	.	.	*	1,627
12	48	1,193	42	*	.	*	.	.	*	1,241
13	51	921	46	.	.	*	.	.	*	972
14	68	799	60	.	*	.	.	*	*	867
15	54	670	49	.	.	.	.	.	*	724
16	67	492	60	.	*	.	.	.	*	559
17	17	159	*	.	.	.	.	.	12	176
18	47	187	*	.	.	.	.	.	43	234
19	40	130	*	.	.	.	.	.	32	170
20	47	125	*	.	.	.	.	.	42	172
21	*	87	*	.	.	.	.	.	.	88
22	.	72	.	.	.	.	.	.	.	72
23	*	37	*	.	.	.	.	.	.	39
24	.	22	.	.	.	.	.	.	.	22
25	.	13	.	.	.	.	.	.	.	13
26	.	*	.	.	.	.	.	.	.	*
27	.	*	.	.	.	.	.	.	.	*
28	.	*	.	.	.	.	.	.	.	*
All	174,873	66,695	143,186	149	672	21,512	4,355	626	4,373	241,568
<b>1997</b>										
0	160,056	11,575	133,988	345	1,328	11,977	8,592	420	3,406	171,631
1	44,752	10,447	38,579	94	380	3,253	1,355	46	1,045	55,199
2	27,338	9,548	24,765	38	84	1,615	387	34	415	36,886
3	5,934	8,296	4,403	*	18	331	43	27	1,108	14,230
4	1,303	7,630	1,060	.	*	102	14	*	114	8,933
5	716	6,389	592	*	*	42	*	*	64	7,105
6	538	5,855	446	*	.	28	*	*	54	6,393
7	380	5,075	322	.	*	18	*	*	25	5,455
8	232	4,317	195	.	*	20	*	*	11	4,549
9	136	3,688	101	.	.	10	*	.	21	3,824
10	111	3,095	88	.	*	*	.	*	*	3,206
11	76	2,740	59	.	*	10	*	.	*	2,816
12	80	2,050	65	*	*	11	.	*	*	2,130
13	56	1,701	46	.	*	*	.	.	*	1,757
14	28	1,456	23	.	.	*	*	.	.	1,484
15	22	1,200	19	.	.	*	.	.	*	1,222
16	16	925	12	*	.	*	.	.	*	941
17	22	739	18	.	.	*	.	.	*	761
18	34	633	29	.	*	.	.	.	*	667
19	23	527	21	.	.	.	.	.	*	550
20	33	374	31	.	*	.	.	.	*	407
21	*	105	*	.	.	.	.	.	*	114
22	19	129	.	.	.	.	.	.	19	148
23	16	100	*	.	.	.	.	.	15	116
24	24	88	*	.	.	.	.	.	23	112
25	*	65	*	.	.	.	.	.	.	66
26	.	48	.	.	.	.	.	.	.	48
27	*	20	*	.	.	.	.	.	.	21
28	.	15	.	.	.	.	.	.	.	15
29	.	*	.	.	.	.	.	.	.	*
30	.	14	.	.	.	.	.	.	.	14
All	241,956	88,851	204,868	485	1,831	17,439	10,411	566	6,356	330,807

The point prevalent cohorts and associated modalities are determined on December 31 of each calendar year without applying the 60-day stable modality rule.

\* Values for cells with fewer than 10 patients are suppressed.

." Zero values in this cell.

Table D.10 (continued)

**Prevalent ESRD Patients for Selected Years, by Number of Years on Modality***point prevalent patients on December 31; by year-end treatment modality*

2001	Dialysis	Functioning graft	Center hemo	Center self hemo	Home hemo	CAPD	CCPD	Other PD	Uncertain dialysis	All
0	184,873	14,297	164,125	246	829	7,685	9,572	331	2,085	199,170
1	56,766	12,753	52,107	37	161	2,339	1,731	48	343	69,519
2	27,904	11,336	25,697	23	77	1,224	714	25	144	39,240
3	12,088	10,512	11,435	*	28	370	213	*	31	22,600
4	4,528	9,066	4,300	*	*	137	57	*	16	13,594
5	2,400	8,042	2,267	*	*	71	31	*	12	10,442
6	2,153	7,103	2,065	*	*	45	12	*	18	9,256
7	535	6,025	436	.	*	12	*	*	80	6,560
8	237	5,479	205	.	.	*	*	*	16	5,716
9	189	4,419	163	.	.	10	*	*	*	4,608
10	164	4,086	144	.	.	*	.	*	11	4,250
11	152	3,538	134	.	*	*	*	*	*	3,690
12	80	3,047	72	.	.	*	.	*	.	3,127
13	42	2,554	29	.	.	*	*	.	*	2,596
14	27	2,120	18	.	*	*	*	*	*	2,147
15	11	1,897	*	.	.	*	.	.	*	1,908
16	14	1,445	11	.	.	*	.	.	.	1,459
17	10	1,232	*	.	.	*	.	.	.	1,242
18	*	1,100	*	.	.	*	.	.	.	1,104
19	*	904	*	.	.	*	.	.	.	908
20	*	718	*	.	.	.	.	.	.	720
21	*	579	*	.	.	.	.	.	*	586
22	*	506	*	.	.	.	.	.	.	511
23	*	420	*	.	.	.	.	.	.	424
24	*	287	*	.	.	.	.	.	.	294
25	.	79	.	.	.	.	.	.	.	79
26	*	96	.	.	.	.	.	.	*	99
27	*	70	.	.	.	.	.	.	*	71
28	*	56	.	.	.	.	.	.	*	61
29	.	42	.	.	.	.	.	.	.	42
30	.	28	.	.	.	.	.	.	.	28
31	.	14	.	.	.	.	.	.	.	14
32	.	*	.	.	.	.	.	.	.	*
33	.	*	.	.	.	.	.	.	.	*
34	.	*	.	.	.	.	.	.	.	*
All	292,215	113,866	263,255	333	1,122	11,927	12,341	445	2,792	406,081

The point prevalent cohorts and associated modalities are determined on December 31 of each calendar year without applying the 60-day stable modality rule.

\* Values for cells with fewer than 10 patients are suppressed.

." Zero values in this cell.

Table D.11  
**Incident ESRD Patients for Selected Years, by Modality, Diabetic Status, & State/Territory**

1993	Center hemo	Center self hemo	Home hemo	CAPD	CCPD	Other PD	Uncertain dialysis	Transplant	All
Alabama	949	*	.	96	17	15	12	45	1,135
Alaska	44	.	.	*	.	*	.	*	50
Arizona	885	.	*	90	*	*	15	27	1,032
Arkansas	435	.	.	93	14	*	*	18	570
California	5,236	10	*	580	44	19	498	267	6,657
Colorado	436	.	.	58	13	*	*	23	539
Connecticut	469	.	*	148	19	*	*	33	683
Delaware	153	.	.	22	*	*	12	*	195
District of Columbia	295	.	*	15	*	*	*	*	324
Florida	3,195	*	*	276	75	38	36	67	3,697
Georgia	1,548	.	*	207	36	32	22	42	1,889
Hawaii	295	.	.	17	.	.	*	*	323
Idaho	99	.	.	28	*	*	*	*	137
Illinois	2,507	*	*	280	28	18	148	126	3,114
Indiana	953	.	.	214	*	*	19	55	1,257
Iowa	427	*	.	59	*	*	*	27	527
Kansas	329	.	.	68	*	*	.	12	416
Kentucky	665	.	.	85	13	12	*	36	816
Louisiana	1,150	.	.	112	18	*	10	20	1,313
Maine	138	.	.	16	*	*	.	*	165
Maryland	1,135	.	*	125	12	*	17	38	1,334
Massachusetts	1,055	*	.	101	*	12	14	50	1,238
Michigan	1,798	*	*	323	*	*	*	93	2,241
Minnesota	666	.	.	35	*	*	*	81	795
Mississippi	676	*	.	69	16	*	*	*	775
Missouri	1,030	.	*	137	23	*	*	37	1,243
Montana	104	.	.	14	.	*	.	16	142
Nebraska	173	.	.	82	*	*	*	11	276
Nevada	236	*	.	28	*	*	*	*	281
New Hampshire	116	.	.	19	*	*	*	*	149
New Jersey	1,725	*	*	294	78	37	49	49	2,234
New Mexico	280	.	.	52	*	.	*	13	360
New York	3,474	*	*	550	54	66	28	127	4,304
North Carolina	1,457	*	*	232	126	71	21	33	1,945
North Dakota	115	.	.	*	.	.	.	*	126
Ohio	1,967	.	*	302	27	77	20	105	2,501
Oklahoma	555	.	.	87	*	37	*	25	709
Oregon	391	.	.	81	*	13	41	*	539
Pennsylvania	2,352	*	*	221	28	51	95	115	2,867
Rhode Island	208	.	.	12	*	.	*	*	229
South Carolina	900	*	*	102	18	*	20	28	1,082
South Dakota	111	.	*	*	*	.	*	*	128
Tennessee	1,044	.	.	92	*	*	*	27	1,184
Texas	3,909	*	*	291	77	31	47	95	4,454
Utah	220	.	.	11	*	*	*	24	265
Vermont	57	.	.	*	*	.	.	*	74
Virginia	1,316	.	.	172	*	29	12	40	1,577
Washington	672	17	.	58	19	*	*	35	813
West Virginia	306	.	.	81	*	*	*	*	408
Wisconsin	724	*	.	115	15	*	15	54	930
Wyoming	43	.	.	*	*	*	.	*	53
American Samoa	*	.	.	.	.	.	.	.	*
Guam	22	.	.	11	.	.	.	*	34
Puerto Rico	644	.	.	52	13	*	14	*	740
Saipan	*	.	.	.	.	.	.	.	*
Virgin Islands	16	.	.	*	.	.	.	.	17
Foreign	12	.	.	.	.	.	.	41	53
Unknown	506	.	*	32	*	*	361	59	976
All	50,238	52	59	6,274	891	693	1,621	2,102	61,930

The incident cohorts and associated modalities are determined at the time of ESRD initiation without applying the 60-day stable modality rule.

\* Values for cells with fewer than 10 patients are suppressed.

." Zero values in this cell.

On our website: Incident ESRD Patients for Selected Years, by State (D.g).

Table D.11 (continued)

## Incident ESRD Patients for Selected Years, by Modality, Diabetic Status, &amp; State/Territory

1993 DIABETIC	Center hemo	Center self hemo	Home hemo	CAPD	CCPD	Other PD	Uncertain dialysis	Transplant	All
Alabama	316	.	.	29	*	*	*	*	365
Alaska	21	.	.	*	.	*	.	*	24
Arizona	432	.	*	38	*	*	*	*	483
Arkansas	125	.	.	41	*	*	.	*	178
California	2,014	*	*	212	10	*	157	13	2,417
Colorado	201	.	.	26	*	*	*	10	244
Connecticut	143	.	*	61	*	*	*	*	220
Delaware	51	.	.	*	.	*	*	.	63
District of Columbia	79	.	.	*	.	.	*	.	86
Florida	942	*	*	86	18	16	13	12	1,090
Georgia	514	.	*	69	*	12	*	*	612
Hawaii	170	.	.	10	.	.	*	.	181
Idaho	42	.	.	11	.	.	*	.	55
Illinois	725	.	*	99	*	*	26	18	882
Indiana	291	.	.	69	*	*	*	*	373
Iowa	156	*	.	28	*	*	.	*	196
Kansas	119	.	.	34	*	.	.	.	154
Kentucky	230	.	.	30	*	*	*	*	271
Louisiana	374	.	.	40	*	*	*	.	428
Maine	48	.	.	*	.	.	.	*	59
Maryland	381	.	.	38	*	*	*	*	432
Massachusetts	339	.	.	23	*	*	*	*	378
Michigan	636	*	.	125	*	*	*	16	786
Minnesota	242	.	.	22	*	*	*	12	283
Mississippi	230	.	.	24	*	*	*	*	262
Missouri	372	.	.	63	14	*	*	*	461
Montana	37	.	.	*	.	*	.	*	54
Nebraska	70	.	.	34	*	*	.	*	110
Nevada	80	.	.	*	*	.	.	*	92
New Hampshire	35	.	.	*	*	.	.	.	41
New Jersey	530	.	*	117	28	15	*	*	703
New Mexico	159	.	.	34	*	.	*	*	202
New York	1,086	.	*	194	18	29	*	*	1,339
North Carolina	526	*	.	96	42	21	*	*	693
North Dakota	47	.	.	*	.	.	.	*	50
Ohio	718	.	.	110	*	31	*	19	889
Oklahoma	248	.	.	40	.	14	*	*	307
Oregon	137	.	.	33	*	*	12	*	188
Pennsylvania	798	.	*	86	10	17	21	*	938
Rhode Island	66	.	.	*	.	.	.	.	67
South Carolina	314	.	*	34	*	*	*	*	364
South Dakota	57	.	*	.	*	.	*	*	61
Tennessee	357	.	.	28	*	.	*	*	394
Texas	1,710	.	*	106	25	12	17	12	1,883
Utah	94	.	.	*	*	.	*	*	111
Vermont	23	.	.	*	.	.	.	*	28
Virginia	435	.	.	68	*	13	*	*	524
Washington	251	*	.	27	*	*	.	14	307
West Virginia	96	.	.	37	*	*	*	.	138
Wisconsin	265	*	.	45	*	*	*	15	344
Wyoming	22	.	.	*	.	.	.	*	25
American Samoa	*	.	.	.	.	.	.	.	*
Guam	12	.	.	*	.	.	.	.	17
Puerto Rico	308	.	.	21	*	*	*	.	343
Saipan	*	.	.	.	.	.	.	.	*
Virgin Islands	*	.	.	.	.	.	.	.	*
Foreign	*	.	.	.	.	.	.	.	*
Unknown	152	.	*	11	*	*	31	*	202
All	17,841	17	17	2,362	284	254	375	262	21,412

The incident cohorts and associated modalities are determined at the time of ESRD initiation without applying the 60-day stable modality rule.

\* Values for cells with fewer than 10 patients are suppressed.

.. Zero values in this cell.

On our website: Incident ESRD Patients for Selected Years, by State (D.g).

Table D.11 (continued)

## Incident ESRD Patients for Selected Years, by Modality, Diabetic Status, &amp; State/Territory

1993 NON-DIABETIC	Center hemo	Center self hemo	Home hemo	CAPD	CCPD	Other PD	Uncertain dialysis	Transplant	All
Alabama	633	*	.	67	10	*	10	41	770
Alaska	23	.	.	*	.	.	.	*	26
Arizona	453	.	*	52	*	*	*	26	549
Arkansas	310	.	.	52	10	*	*	13	392
California	3,222	*	*	368	34	14	341	254	4,240
Colorado	235	.	.	32	*	.	*	13	295
Connecticut	326	.	*	87	14	*	*	28	463
Delaware	102	.	.	15	*	.	*	*	132
District of Columbia	216	.	*	*	*	*	*	*	238
Florida	2,253	.	*	190	57	22	23	55	2,607
Georgia	1,034	.	*	138	27	20	19	38	1,277
Hawaii	125	.	.	*	.	.	*	*	142
Idaho	57	.	.	17	*	*	.	*	82
Illinois	1,782	*	*	181	20	13	122	108	2,232
Indiana	662	.	.	145	*	*	15	50	884
Iowa	271	.	.	31	*	*	*	19	331
Kansas	210	.	.	34	*	*	.	12	262
Kentucky	435	.	.	55	10	*	*	34	545
Louisiana	776	.	.	72	*	*	*	20	885
Maine	90	.	.	*	*	*	.	*	106
Maryland	754	.	*	87	10	*	10	35	902
Massachusetts	716	*	.	78	*	*	12	42	860
Michigan	1,162	*	*	198	*	*	*	77	1,455
Minnesota	424	.	.	13	.	*	*	69	512
Mississippi	446	*	.	45	12	.	*	*	513
Missouri	658	.	*	74	*	*	*	29	782
Montana	67	.	.	*	.	*	.	10	88
Nebraska	103	.	.	48	*	*	*	*	166
Nevada	156	*	.	19	*	*	*	*	189
New Hampshire	81	.	.	14	.	*	*	*	108
New Jersey	1,195	*	.	177	50	22	41	45	1,531
New Mexico	121	.	.	18	*	.	*	11	158
New York	2,388	*	*	356	36	37	24	120	2,965
North Carolina	931	*	*	136	84	50	18	29	1,252
North Dakota	68	.	.	*	.	.	.	*	76
Ohio	1,249	.	*	192	21	46	15	86	1,612
Oklahoma	307	.	.	47	*	23	*	21	402
Oregon	254	.	.	48	*	*	29	*	351
Pennsylvania	1,554	*	.	135	18	34	74	113	1,929
Rhode Island	142	.	.	11	*	.	*	*	162
South Carolina	586	*	*	68	13	*	15	24	718
South Dakota	54	.	.	*	*	.	*	*	67
Tennessee	687	.	.	64	*	*	*	22	790
Texas	2,199	*	*	185	52	19	30	83	2,571
Utah	126	.	.	*	*	*	*	15	154
Vermont	34	.	.	*	*	.	.	*	46
Virginia	881	.	.	104	*	16	*	37	1,053
Washington	421	11	.	31	14	*	*	21	506
West Virginia	210	.	.	44	*	*	*	*	270
Wisconsin	459	*	.	70	*	*	*	39	586
Wyoming	21	.	.	*	*	*	.	*	28
American Samoa	*	.	.	.	.	.	.	.	*
Guam	10	.	.	*	.	.	.	*	17
Puerto Rico	336	.	.	31	*	*	10	*	397
Saipan	*	.	.	.	.	.	.	.	*
Virgin Islands	10	.	.	*	.	.	.	.	11
Foreign	10	.	.	.	.	.	.	41	51
Unknown	354	.	*	21	*	*	330	57	774
All	32,397	35	42	3,912	607	439	1,246	1,840	40,518

The incident cohorts and associated modalities are determined at the time of ESRD initiation without applying the 60-day stable modality rule.

\* Values for cells with fewer than 10 patients are suppressed.

." Zero values in this cell.

On our website: Incident ESRD Patients for Selected Years, by State (D.g).

Table D.11 (continued)

## Incident ESRD Patients for Selected Years, by Modality, Diabetic Status, &amp; State/Territory

1997	Center	Center	Home				Uncertain		
ALL	hemo	self hemo	hemo	CAPD	CCPD	Other PD	dialysis	Transplant	All
Alabama	1,426	.	*	142	38	*	*	32	1,650
Alaska	61	*	.	15	*	.	.	*	81
Arizona	1,236	.	13	89	15	*	*	17	1,379
Arkansas	708	.	*	74	24	*	*	12	826
California	8,020	*	*	701	123	14	108	177	9,150
Colorado	598	.	*	55	18	*	*	21	700
Connecticut	686	*	.	161	34	*	25	14	922
Delaware	194	.	*	13	*	.	*	10	230
District of Columbia	393	.	*	*	.	*	*	*	408
Florida	4,322	*	48	222	176	27	85	82	4,964
Georgia	2,198	*	11	179	46	19	27	26	2,507
Hawaii	392	.	.	20	*	.	*	*	426
Idaho	180	.	*	22	*	.	*	11	224
Illinois	3,308	.	10	228	88	12	56	100	3,802
Indiana	1,304	*	*	195	24	*	33	31	1,594
Iowa	485	*	*	59	23	*	20	19	610
Kansas	546	*	.	86	*	*	*	10	655
Kentucky	864	.	*	64	30	13	29	20	1,021
Louisiana	1,697	.	.	68	36	*	*	26	1,834
Maine	241	.	.	13	*	.	*	*	270
Maryland	1,669	.	14	81	67	*	16	46	1,895
Massachusetts	1,403	.	*	83	34	*	16	37	1,578
Michigan	2,467	.	*	304	39	*	46	63	2,931
Minnesota	850	*	.	44	26	*	12	68	1,003
Mississippi	938	.	*	77	35	14	*	10	1,082
Missouri	1,351	.	*	170	52	*	15	35	1,631
Montana	126	.	*	*	*	*	*	*	146
Nebraska	287	.	*	95	*	.	*	12	409
Nevada	342	*	*	32	*	.	*	20	410
New Hampshire	161	*	*	19	*	.	*	*	199
New Jersey	2,419	.	.	196	158	19	53	50	2,895
New Mexico	410	*	*	41	24	*	*	*	489
New York	5,221	14	13	323	103	59	79	89	5,901
North Carolina	2,133	.	*	208	149	74	32	32	2,629
North Dakota	115	.	.	13	.	.	*	*	138
Ohio	2,827	.	*	283	83	77	71	63	3,407
Oklahoma	801	*	*	70	26	29	10	11	949
Oregon	521	.	*	86	*	.	*	11	632
Pennsylvania	3,685	.	10	157	100	23	86	81	4,142
Rhode Island	249	.	.	17	.	.	*	.	269
South Carolina	1,292	*	*	71	38	*	21	33	1,459
South Dakota	165	.	.	21	*	.	*	*	199
Tennessee	1,525	*	*	59	62	*	18	29	1,703
Texas	5,284	*	*	238	153	*	50	79	5,821
Utah	209	.	*	21	*	.	*	22	262
Vermont	89	.	.	11	*	.	*	*	112
Virginia	2,054	.	*	126	57	24	36	41	2,343
Washington	891	20	*	85	19	*	*	29	1,052
West Virginia	515	.	.	35	31	*	17	11	617
Wisconsin	967	.	.	108	38	*	40	67	1,227
Wyoming	54	.	.	11	*	.	.	*	70
American Samoa	*	.	.	.	.	.	.	.	*
Guam	69	.	.	.	.	.	.	.	69
Puerto Rico	792	.	*	76	75	*	11	*	966
Saipan	17	.	.	.	.	.	.	.	17
Virgin Islands	43	.	.	.	*	.	.	.	44
Foreign	12	.	.	.	.	.	.	25	37
Unknown	*	.	.	.	.	.	*	23	34
All	70,828	53	191	5,581	2,123	486	1,116	1,650	82,028

The incident cohorts and associated modalities are determined at the time of ESRD initiation without applying the 60-day stable modality rule.

\* Values for cells with fewer than 10 patients are suppressed.

.. Zero values in this cell.

On our website: Incident ESRD Patients for Selected Years, by State (D.g).

Table D.11 (continued)

## Incident ESRD Patients for Selected Years, by Modality, Diabetic Status, &amp; State/Territory

1997 DIABETIC	Center hemo	Center self hemo	Home hemo	CAPD	CCPD	Other PD	Uncertain dialysis	Transplant	All
Alabama	641	.	.	51	16	*	*	*	717
Alaska	37	.	.	*	.	.	.	*	44
Arizona	667	.	*	44	*	.	*	*	731
Arkansas	274	.	.	32	*	*	*	*	319
California	3,829	*	*	309	51	*	46	17	4,262
Colorado	280	.	.	27	*	*	*	*	319
Connecticut	300	*	.	90	14	*	*	*	418
Delaware	76	.	.	*	.	.	*	*	86
District of Columbia	144	.	.	*	.	.	*	*	149
Florida	1,646	*	15	106	64	18	31	*	1,891
Georgia	919	*	*	72	20	*	13	*	1,042
Hawaii	232	.	.	12	.	.	*	.	248
Idaho	77	.	.	14	*	.	*	*	100
Illinois	1,265	.	*	89	39	*	19	30	1,451
Indiana	541	*	*	86	10	*	13	*	659
Iowa	200	.	.	27	10	.	*	*	249
Kansas	223	.	.	34	*	*	*	*	264
Kentucky	379	.	.	28	10	*	13	*	437
Louisiana	755	.	.	19	*	.	*	*	794
Maine	90	.	.	*	*	.	*	*	103
Maryland	722	.	*	27	23	*	*	11	790
Massachusetts	536	.	*	33	18	.	*	*	598
Michigan	1,097	.	*	152	13	*	25	18	1,308
Minnesota	333	*	.	16	10	*	*	19	387
Mississippi	399	.	*	25	11	*	.	*	442
Missouri	615	.	*	87	22	*	*	*	741
Montana	61	.	.	*	.	*	*	*	68
Nebraska	124	.	*	33	*	.	*	*	175
Nevada	154	.	.	20	*	.	*	*	184
New Hampshire	62	.	.	*	*	.	*	*	76
New Jersey	992	.	.	80	64	*	21	12	1,175
New Mexico	248	.	.	17	10	*	.	.	277
New York	2,117	*	*	134	46	16	29	12	2,365
North Carolina	932	.	.	96	55	32	20	*	1,139
North Dakota	52	.	.	*	.	.	.	*	58
Ohio	1,263	.	*	155	32	43	32	11	1,537
Oklahoma	387	*	*	35	10	17	*	*	456
Oregon	204	.	*	45	*	.	.	*	255
Pennsylvania	1,494	.	*	72	35	*	37	10	1,659
Rhode Island	97	.	.	*	.	.	*	.	107
South Carolina	541	*	*	22	17	*	*	*	599
South Dakota	72	.	.	11	*	.	.	*	91
Tennessee	606	*	*	24	35	*	*	*	682
Texas	2,782	*	*	131	64	*	20	*	3,014
Utah	102	.	*	15	*	.	*	*	127
Vermont	42	.	.	*	.	.	.	*	51
Virginia	874	.	*	66	14	*	16	*	984
Washington	395	*	*	38	*	.	*	10	458
West Virginia	232	.	.	21	15	*	11	*	283
Wisconsin	420	.	.	58	15	*	12	23	531
Wyoming	29	.	.	*	*	.	.	.	36
American Samoa	*	.	.	.	.	.	.	.	*
Guam	47	.	.	.	.	.	.	.	47
Puerto Rico	454	.	*	43	38	*	*	.	544
Saipan	13	.	.	.	.	.	.	.	13
Virgin Islands	19	.	.	.	*	.	.	.	20
Foreign	*	.	.	.	.	.	.	.	*
Unknown	.	.	.	.	.	.	.	.	.
All	31,099	29	67	2,547	839	216	465	305	35,567

The incident cohorts and associated modalities are determined at the time of ESRD initiation without applying the 60-day stable modality rule.

\* Values for cells with fewer than 10 patients are suppressed.

." Zero values in this cell.

On our website: Incident ESRD Patients for Selected Years, by State (D.g).



Table D.11 (continued)

**Incident ESRD Patients for Selected Years, by Modality, Diabetic Status, & State/Territory**

1997 NON-DIABETIC	Center hemo	Center self hemo	Home hemo	CAPD	CCPD	Other PD	Uncertain dialysis	Transplant	All
Alabama	785	.	*	91	22	*	*	29	933
Alaska	24	.	.	*	*	.	.	*	37
Arizona	569	.	.	45	10	*	*	14	648
Arkansas	434	.	*	42	18	*	*	*	507
California	4,191	.	*	392	72	*	62	160	4,888
Colorado	318	.	*	28	13	.	*	17	381
Connecticut	386	.	.	71	20	.	17	10	504
Delaware	118	.	*	*	*	.	*	*	144
District of Columbia	249	.	*	*	.	*	*	*	259
Florida	2,676	.	33	116	112	*	54	73	3,073
Georgia	1,279	.	*	107	26	10	14	22	1,465
Hawaii	160	.	.	*	*	.	*	*	178
Idaho	103	.	*	*	*	.	.	*	124
Illinois	2,043	.	*	139	49	*	37	70	2,351
Indiana	763	.	*	109	14	.	20	27	935
Iowa	285	*	*	32	13	*	12	15	361
Kansas	323	*	.	52	*	*	*	*	391
Kentucky	485	.	*	36	20	*	16	18	584
Louisiana	942	.	.	49	27	*	*	18	1,040
Maine	151	.	.	*	*	.	*	*	167
Maryland	947	.	11	54	44	.	14	35	1,105
Massachusetts	867	.	.	50	16	*	10	34	980
Michigan	1,370	.	*	152	26	*	21	45	1,623
Minnesota	517	.	.	28	16	.	*	49	616
Mississippi	539	.	*	52	24	*	*	*	640
Missouri	736	.	*	83	30	*	10	27	890
Montana	65	.	*	*	*	.	.	*	78
Nebraska	163	.	.	62	*	.	*	*	234
Nevada	188	*	*	12	*	.	*	15	226
New Hampshire	99	*	*	12	*	.	*	*	123
New Jersey	1,427	.	.	116	94	13	32	38	1,720
New Mexico	162	*	*	24	14	.	*	*	212
New York	3,104	*	*	189	57	43	50	77	3,536
North Carolina	1,201	.	*	112	94	42	12	28	1,490
North Dakota	63	.	.	*	.	.	*	*	80
Ohio	1,564	.	*	128	51	34	39	52	1,870
Oklahoma	414	.	.	35	16	12	*	*	493
Oregon	317	.	*	41	*	.	*	*	377
Pennsylvania	2,191	.	*	85	65	16	49	71	2,483
Rhode Island	152	.	.	10	.	.	.	.	162
South Carolina	751	.	.	49	21	.	13	26	860
South Dakota	93	.	.	10	*	.	*	*	108
Tennessee	919	.	*	35	27	*	*	26	1,021
Texas	2,502	.	*	107	89	*	30	73	2,807
Utah	107	.	.	*	*	.	.	16	135
Vermont	47	.	.	*	*	.	*	*	61
Virginia	1,180	.	*	60	43	16	20	36	1,359
Washington	496	11	*	47	16	*	.	19	594
West Virginia	283	.	.	14	16	*	*	*	334
Wisconsin	547	.	.	50	23	*	28	44	696
Wyoming	25	.	.	*	*	.	.	*	34
American Samoa	*	.	.	.	.	.	.	.	*
Guam	22	.	.	.	.	.	.	.	22
Puerto Rico	338	.	.	33	37	*	*	*	422
Saipan	*	.	.	.	.	.	.	.	*
Virgin Islands	24	.	.	.	.	.	.	.	24
Foreign	*	.	.	.	.	.	.	25	34
Unknown	*	.	.	.	.	.	*	23	34
All	39,729	24	124	3,034	1,284	270	651	1,345	46,461

The incident cohorts and associated modalities are determined at the time of ESRD initiation without applying the 60-day stable modality rule.

\* Values for cells with fewer than 10 patients are suppressed.

.. Zero values in this cell.

On our website: Incident ESRD Patients for Selected Years, by State (D.g).

Table D.11 (continued)

## Incident ESRD Patients for Selected Years, by Modality, Diabetic Status, &amp; State/Territory

2001	Center	Center	Home				Uncertain		
ALL	hemo	self hemo	hemo	CAPD	CCPD	Other PD	dialysis	Transplant	All
Alabama	1,612	.	*	80	55	*	*	28	1,778
Alaska	75	.	.	15	.	.	.	*	94
Arizona	1,579	.	*	88	42	*	*	34	1,761
Arkansas	771	.	.	82	16	*	.	14	885
California	10,265	.	*	662	157	10	37	242	11,375
Colorado	735	.	.	41	*	.	*	27	812
Connecticut	785	.	*	113	38	.	*	32	973
Delaware	247	.	.	13	*	.	.	*	269
District of Columbia	427	.	.	*	*	.	*	*	444
Florida	5,269	*	64	231	112	28	22	131	5,858
Georgia	2,862	.	*	197	85	*	*	50	3,214
Hawaii	438	.	.	17	*	.	*	12	475
Idaho	226	.	*	44	.	.	.	*	276
Illinois	3,732	.	67	231	99	*	24	143	4,300
Indiana	1,660	*	*	158	27	*	*	48	1,900
Iowa	631	.	*	45	11	.	*	42	738
Kansas	569	.	*	70	11	.	*	12	665
Kentucky	1,119	.	*	86	17	12	*	22	1,264
Louisiana	2,062	.	*	74	34	*	*	35	2,211
Maine	258	.	*	11	*	.	.	15	292
Maryland	1,966	.	*	87	64	*	10	85	2,219
Massachusetts	1,447	.	*	71	64	.	*	56	1,646
Michigan	3,112	*	.	257	33	*	*	88	3,508
Minnesota	1,061	.	.	31	20	*	*	116	1,233
Mississippi	1,099	*	*	76	31	*	*	11	1,229
Missouri	1,688	.	*	127	56	*	*	39	1,923
Montana	181	.	.	19	*	.	.	10	211
Nebraska	453	.	.	59	*	*	*	10	534
Nevada	537	.	.	38	12	.	.	21	608
New Hampshire	189	.	.	30	*	.	*	*	234
New Jersey	2,899	.	*	103	94	*	*	111	3,215
New Mexico	493	.	.	36	16	.	*	13	559
New York	6,067	*	*	195	119	61	22	157	6,632
North Carolina	2,660	*	*	152	126	70	*	52	3,070
North Dakota	160	.	.	14	*	.	.	17	193
Ohio	3,644	*	*	261	96	*	11	108	4,140
Oklahoma	1,002	.	.	56	45	16	*	*	1,132
Oregon	709	.	*	54	14	.	*	34	813
Pennsylvania	4,102	.	*	88	110	*	11	110	4,432
Rhode Island	291	.	.	*	*	*	*	11	317
South Carolina	1,496	*	.	70	77	.	*	33	1,680
South Dakota	188	.	.	*	11	.	.	14	220
Tennessee	1,653	*	.	49	92	*	*	31	1,833
Texas	6,537	.	12	233	202	12	30	120	7,146
Utah	358	.	.	15	15	.	*	14	403
Vermont	95	.	.	*	*	*	.	*	103
Virginia	2,254	*	*	117	40	*	*	71	2,495
Washington	1,166	14	*	66	16	*	*	35	1,305
West Virginia	626	.	*	30	57	*	*	11	728
Wisconsin	1,279	.	*	62	48	14	*	65	1,472
Wyoming	73	*	.	*	*	.	.	*	88
American Samoa	16	.	.	.	.	.	*	.	17
Guam	83	.	.	*	.	.	*	.	86
Puerto Rico	1,040	.	*	25	68	*	*	*	1,150
Saipan	*	.	.	.	.	.	.	.	*
Virgin Islands	40	.	.	.	*	.	.	*	42
Foreign	15	.	.	.	.	.	.	19	34
Unknown	12	.	.	*	*	.	29	*	52
All	86,022	38	229	4,711	2,280	299	304	2,412	96,295

The incident cohorts and associated modalities are determined at the time of ESRD initiation without applying the 60-day stable modality rule.

\* Values for cells with fewer than 10 patients are suppressed.

." Zero values in this cell.

On our website: Incident ESRD Patients for Selected Years, by State (D.g).

Table D.11 (continued)

## Incident ESRD Patients for Selected Years, by Modality, Diabetic Status, &amp; State/Territory

2001 DIABETIC	Center hemo	Center self hemo	Home hemo	CAPD	CCPD	Other PD	Uncertain dialysis	Transplant	All
Alabama	756	.	*	33	22	.	.	*	817
Alaska	40	.	.	*	.	.	.	.	48
Arizona	850	.	*	44	19	.	*	*	924
Arkansas	315	.	.	40	*	*	.	*	367
California	4,845	.	*	279	49	*	*	32	5,217
Colorado	351	.	.	23	*	.	.	*	383
Connecticut	320	.	*	44	15	.	.	*	382
Delaware	103	.	.	*	.	.	.	*	106
District of Columbia	157	.	.	.	*	.	.	.	159
Florida	2,201	.	18	96	45	11	10	24	2,405
Georgia	1,207	.	*	79	33	*	*	11	1,338
Hawaii	280	.	.	10	*	.	*	*	297
Idaho	87	.	.	18	.	.	.	.	105
Illinois	1,533	.	26	91	37	*	*	19	1,709
Indiana	716	*	*	59	10	*	*	*	795
Iowa	259	.	*	21	*	.	.	*	297
Kansas	269	.	.	30	*	.	*	*	304
Kentucky	532	.	*	40	*	*	*	*	588
Louisiana	935	.	*	24	10	*	.	*	974
Maine	119	.	.	*	*	.	.	*	124
Maryland	822	.	.	37	25	*	*	*	892
Massachusetts	636	.	.	28	29	.	*	*	703
Michigan	1,435	.	.	118	12	*	.	24	1,593
Minnesota	394	.	.	*	*	*	.	31	441
Mississippi	486	*	*	34	12	*	.	.	536
Missouri	702	.	*	60	29	*	*	*	804
Montana	98	.	.	11	.	.	.	*	113
Nebraska	208	.	.	28	*	.	.	*	241
Nevada	254	.	.	17	*	.	.	*	277
New Hampshire	67	.	.	11	*	.	*	*	82
New Jersey	1,211	.	.	41	33	*	.	15	1,301
New Mexico	315	.	.	22	*	.	.	*	350
New York	2,599	*	*	95	44	19	*	21	2,786
North Carolina	1,162	.	*	74	38	41	.	*	1,324
North Dakota	75	.	.	*	.	.	.	*	84
Ohio	1,754	*	*	132	39	*	.	10	1,942
Oklahoma	488	.	.	30	24	*	.	*	551
Oregon	284	.	*	27	*	.	*	*	325
Pennsylvania	1,843	.	*	37	39	*	*	*	1,934
Rhode Island	106	.	.	*	*	*	.	*	114
South Carolina	696	.	.	29	33	.	*	*	762
South Dakota	87	.	.	*	*	.	.	*	100
Tennessee	733	.	.	18	42	*	.	*	798
Texas	3,469	.	*	114	83	*	12	14	3,704
Utah	191	.	.	*	*	.	.	*	205
Vermont	40	.	.	*	*	.	.	*	43
Virginia	1,018	*	*	52	13	*	.	*	1,097
Washington	507	*	.	30	*	*	*	*	557
West Virginia	321	.	*	15	28	*	*	*	370
Wisconsin	535	.	.	19	11	*	*	13	583
Wyoming	30	*	.	*	.	.	.	*	35
American Samoa	11	.	.	.	.	.	*	.	12
Guam	56	.	.	*	.	.	.	.	57
Puerto Rico	655	.	*	14	43	*	.	*	719
Saipan	*	.	.	.	.	.	.	.	*
Virgin Islands	22	.	.	.	.	.	.	.	22
Foreign	*	.	.	.	.	.	.	.	*
Unknown	.	.	.	.	.	.	*	.	*
All	39,201	14	87	2,069	888	131	59	364	42,813

The incident cohorts and associated modalities are determined at the time of ESRD initiation without applying the 60-day stable modality rule.

\* Values for cells with fewer than 10 patients are suppressed.

.. Zero values in this cell.

On our website: Incident ESRD Patients for Selected Years, by State (D.g).

Table D.11 (continued)

## Incident ESRD Patients for Selected Years, by Modality, Diabetic Status, &amp; State/Territory

2001 NON-DIABETIC	Center hemo	Center self hemo	Home hemo	CAPD	CCPD	Other PD	Uncertain dialysis	Transplant	All
Alabama	856	.	.	47	33	*	*	23	961
Alaska	35	.	.	*	.	.	.	*	46
Arizona	729	.	*	44	23	*	*	29	837
Arkansas	456	.	.	42	*	*	.	10	518
California	5,420	.	*	383	108	*	28	210	6,158
Colorado	384	.	.	18	*	.	*	20	429
Connecticut	465	.	*	69	23	.	*	30	591
Delaware	144	.	.	11	*	.	.	*	163
District of Columbia	270	.	.	*	.	.	*	*	285
Florida	3,068	*	46	135	67	17	12	107	3,453
Georgia	1,655	.	*	118	52	*	*	39	1,876
Hawaii	158	.	.	*	*	.	.	11	178
Idaho	139	.	*	26	.	.	.	*	171
Illinois	2,199	.	41	140	62	*	22	124	2,591
Indiana	944	.	.	99	17	*	*	42	1,105
Iowa	372	.	*	24	*	.	*	35	441
Kansas	300	.	*	40	*	.	*	11	361
Kentucky	587	.	*	46	10	*	*	21	676
Louisiana	1,127	.	.	50	24	.	*	32	1,237
Maine	139	.	*	*	*	.	.	13	168
Maryland	1,144	.	*	50	39	*	*	79	1,327
Massachusetts	811	.	*	43	35	.	*	47	943
Michigan	1,677	*	.	139	21	*	*	64	1,915
Minnesota	667	.	.	23	13	.	*	85	792
Mississippi	613	*	.	42	19	*	*	11	693
Missouri	986	.	*	67	27	*	*	33	1,119
Montana	83	.	.	*	*	.	.	*	98
Nebraska	245	.	.	31	*	*	*	*	293
Nevada	283	.	.	21	10	.	.	17	331
New Hampshire	122	.	.	19	*	.	.	*	152
New Jersey	1,688	.	*	62	61	*	*	96	1,914
New Mexico	178	.	.	14	*	.	*	*	209
New York	3,468	*	*	100	75	42	18	136	3,846
North Carolina	1,498	*	*	78	88	29	*	44	1,746
North Dakota	85	.	.	*	*	.	.	14	109
Ohio	1,890	*	*	129	57	*	11	98	2,198
Oklahoma	514	.	.	26	21	*	*	*	581
Oregon	425	.	.	27	10	.	.	26	488
Pennsylvania	2,259	.	*	51	71	*	10	101	2,498
Rhode Island	185	.	.	*	*	.	*	*	203
South Carolina	800	*	.	41	44	.	*	30	918
South Dakota	101	.	.	*	*	.	.	10	120
Tennessee	920	*	.	31	50	*	*	27	1,035
Texas	3,068	.	*	119	119	*	18	106	3,442
Utah	167	.	.	*	12	.	*	*	198
Vermont	55	.	.	*	*	*	.	*	60
Virginia	1,236	.	*	65	27	*	*	64	1,398
Washington	659	*	*	36	14	*	*	27	748
West Virginia	305	.	.	15	29	*	.	*	358
Wisconsin	744	.	*	43	37	10	*	52	889
Wyoming	43	.	.	*	*	.	.	*	53
American Samoa	*	.	.	.	.	.	.	.	*
Guam	27	.	.	*	.	.	*	.	29
Puerto Rico	385	.	.	11	25	.	*	*	431
Saipan	.	.	.	.	.	.	.	.	.
Virgin Islands	18	.	.	.	*	.	.	*	20
Foreign	*	.	.	.	.	.	.	19	27
Unknown	12	.	.	*	*	.	28	*	51
All	46,821	24	142	2,642	1,392	168	245	2,048	53,482

The incident cohorts and associated modalities are determined at the time of ESRD initiation without applying the 60-day stable modality rule.

\* Values for cells with fewer than 10 patients are suppressed.

." Zero values in this cell.

On our website: Incident ESRD Patients for Selected Years, by State (D.g).

Table D.12

**Prevalent ESRD Patients for Selected Years, by Modality, Diabetic Status, & State/Territory**  
*point prevalent patients on December 31*

1993	Center	Center	Home				Uncertain		
ALL	hemo	self hemo	hemo	CAPD	CCPD	Other PD	dialysis	Transplant	All
Alabama	2,963	.	*	363	107	*	45	1,258	4,747
Alaska	105	*	.	23	*	*	*	99	233
Arizona	2,396	.	12	262	109	*	42	931	3,757
Arkansas	1,187	.	*	236	35	*	17	579	2,068
California	15,142	59	28	1,901	436	21	944	7,197	25,728
Colorado	1,209	*	*	222	54	*	32	815	2,340
Connecticut	1,423	.	11	442	93	*	16	803	2,795
Delaware	490	.	*	87	*	.	11	191	786
District of Columbia	1,025	.	*	34	*	*	*	198	1,268
Florida	8,941	.	31	957	291	23	117	3,275	13,635
Georgia	5,204	.	21	625	160	18	57	1,633	7,718
Hawaii	922	.	.	60	*	.	17	254	1,255
Idaho	280	.	*	96	*	*	12	252	649
Illinois	6,619	*	19	888	178	*	234	3,375	11,323
Indiana	2,576	.	*	756	39	*	47	1,395	4,821
Iowa	1,027	*	22	222	41	*	24	859	2,205
Kansas	969	.	*	284	33	.	*	577	1,872
Kentucky	1,617	.	*	290	68	*	29	1,048	3,064
Louisiana	3,442	*	*	399	93	*	24	894	4,858
Maine	364	.	.	55	18	.	*	329	773
Maryland	3,284	*	17	433	56	*	48	1,321	5,171
Massachusetts	2,832	.	*	353	66	14	50	1,506	4,825
Michigan	4,620	*	17	1,204	89	16	58	2,572	8,578
Minnesota	1,754	.	.	176	15	*	37	1,596	3,581
Mississippi	2,301	.	46	192	41	*	14	482	3,083
Missouri	2,684	*	23	500	130	*	21	1,431	4,798
Montana	266	.	.	63	*	*	*	233	575
Nebraska	448	.	*	272	26	*	*	464	1,222
Nevada	517	.	*	104	23	*	13	360	1,027
New Hampshire	309	.	.	65	26	*	*	254	664
New Jersey	4,692	12	48	848	263	50	226	1,575	7,714
New Mexico	852	*	.	210	26	*	23	396	1,510
New York	10,395	14	48	1,742	249	54	137	3,530	16,169
North Carolina	4,702	*	18	887	305	50	78	1,429	7,470
North Dakota	232	.	.	31	*	.	11	232	512
Ohio	5,287	.	10	996	158	51	99	3,224	9,825
Oklahoma	1,422	*	*	276	18	17	18	771	2,530
Oregon	999	.	*	297	39	*	59	725	2,126
Pennsylvania	6,901	*	14	815	141	61	173	3,431	11,540
Rhode Island	550	.	.	48	13	.	*	216	831
South Carolina	2,874	*	13	311	137	*	90	885	4,320
South Dakota	287	.	*	21	*	.	*	199	528
Tennessee	2,996	*	55	380	75	*	40	1,367	4,918
Texas	11,596	*	20	1,054	381	68	383	4,124	17,629
Utah	473	.	.	55	*	*	16	561	1,113
Vermont	148	.	.	38	*	*	.	133	326
Virginia	4,048	*	18	635	77	12	54	1,497	6,342
Washington	1,873	13	72	226	50	27	35	1,279	3,575
West Virginia	680	*	*	205	28	*	15	484	1,420
Wisconsin	1,798	13	13	430	53	*	72	1,863	4,244
Wyoming	84	.	*	41	.	.	*	110	240
American Samoa	30	.	.	*	.	.	.	.	31
Guam	75	*	.	27	.	.	*	16	121
Puerto Rico	1,945	.	27	274	56	*	40	401	2,751
Saipan	21	.	.	.	.	.	*	*	24
Virgin Islands	60	.	.	*	*	.	.	*	76
Foreign	47	.	*	*	.	.	*	186	251
Unknown	1,203	.	*	84	10	11	830	1,871	4,013
All	143,186	149	672	21,512	4,355	626	4,373	66,695	241,568

The point prevalent cohorts and associated modalities are determined on December 31 of each calendar year without applying the 60-day stable modality rule.

\* Values for cells with fewer than 10 patients are suppressed.

." Zero values in this cell.

On our website: by Modality, Diabetic Status, & State/Territory (D.h).

Table D.12 (continued)

**Prevalent ESRD Patients for Selected Years, by Modality, Diabetic Status, & State/Territory***point prevalent patients on December 31*

1993 DIABETIC	Center hemo	Center self hemo	Home hemo	CAPD	CCPD	Other PD	Uncertain dialysis	Transplant	All
Alabama	797	.	.	85	28	*	*	136	1,053
Alaska	32	*	.	*	.	*	.	15	57
Arizona	1,101	.	*	105	33	*	18	184	1,445
Arkansas	306	.	.	82	*	*	*	97	491
California	5,328	*	*	572	98	*	302	1,147	7,465
Colorado	473	*	.	85	18	*	*	214	800
Connecticut	405	.	*	139	30	*	*	139	725
Delaware	144	.	.	17	*	.	*	35	199
District of Columbia	285	.	.	*	.	.	.	19	311
Florida	2,424	.	*	246	73	12	29	602	3,390
Georgia	1,516	.	*	154	35	*	10	274	1,997
Hawaii	437	.	.	27	.	.	12	24	500
Idaho	87	.	.	39	*	.	*	68	200
Illinois	1,680	*	*	245	56	*	40	578	2,605
Indiana	735	.	*	221	10	*	12	246	1,226
Iowa	299	*	*	73	14	.	*	189	587
Kansas	298	.	*	111	*	.	.	99	517
Kentucky	493	.	*	76	22	*	*	152	748
Louisiana	1,050	.	.	107	26	.	*	150	1,340
Maine	103	.	.	16	*	.	.	54	176
Maryland	1,006	.	*	104	16	*	14	216	1,364
Massachusetts	748	.	*	78	22	*	11	271	1,135
Michigan	1,439	*	*	408	21	*	13	527	2,418
Minnesota	590	.	.	67	*	*	*	457	1,127
Mississippi	631	.	*	60	12	*	*	58	775
Missouri	823	.	*	165	39	*	*	264	1,302
Montana	100	.	.	26	*	.	.	73	202
Nebraska	157	.	*	92	*	*	.	146	408
Nevada	154	.	*	32	*	.	*	77	276
New Hampshire	70	.	.	20	*	*	*	59	154
New Jersey	1,279	.	*	247	75	12	64	236	1,920
New Mexico	440	.	.	103	11	*	*	72	636
New York	2,872	*	10	478	68	20	23	487	3,961
North Carolina	1,404	.	*	249	88	15	15	230	2,005
North Dakota	99	.	.	13	*	.	*	64	184
Ohio	1,663	.	*	296	45	23	23	682	2,735
Oklahoma	572	.	*	112	*	13	*	155	866
Oregon	282	.	.	91	12	*	19	163	568
Pennsylvania	2,027	*	*	261	55	17	30	625	3,022
Rhode Island	142	.	.	*	*	.	*	31	183
South Carolina	883	*	*	78	26	*	27	136	1,154
South Dakota	143	.	*	*	*	.	*	52	205
Tennessee	832	*	*	105	16	*	*	210	1,178
Texas	4,240	.	*	343	90	18	146	784	5,624
Utah	180	.	.	21	.	.	*	190	395
Vermont	43	.	.	*	*	.	.	31	84
Virginia	1,146	.	*	176	23	*	12	234	1,597
Washington	582	*	19	77	22	*	*	330	1,045
West Virginia	198	.	.	64	*	*	*	103	376
Wisconsin	584	*	*	143	13	.	25	517	1,287
Wyoming	30	.	.	11	.	.	.	31	72
American Samoa	11	.	.	.	.	.	.	.	11
Guam	35	*	.	14	.	.	.	*	52
Puerto Rico	797	.	10	89	20	*	*	30	958
Saipan	*	.	.	.	.	.	*	.	*
Virgin Islands	11	.	.	*	*	.	.	.	13
Foreign	*	.	.	*	.	.	.	*	12
Unknown	341	.	*	26	*	*	71	15	459
All	44,561	31	127	6,491	1,179	199	1,035	11,980	65,603

The point prevalent cohorts and associated modalities are determined on December 31 of each calendar year without applying the 60-day stable modality rule.

\* Values for cells with fewer than 10 patients are suppressed.

." Zero values in this cell.

On our website: by Modality, Diabetic Status, &amp; State/Territory (D.h).

Table D.12 (continued)

**Prevalent ESRD Patients for Selected Years, by Modality, Diabetic Status, & State/Territory***point prevalent patients on December 31*

1993 NON-DIABETIC	Center hemo	Center self hemo	Home hemo	CAPD	CCPD	Other PD	Uncertain dialysis	Transplant	All
Alabama	2,166	.	*	278	79	*	40	1,122	3,694
Alaska	73	.	.	15	*	*	*	84	176
Arizona	1,295	.	11	157	76	*	24	747	2,312
Arkansas	881	.	*	154	32	*	15	482	1,577
California	9,814	50	23	1,329	338	17	642	6,050	18,263
Colorado	736	.	*	137	36	*	25	601	1,540
Connecticut	1,018	.	*	303	63	*	*	664	2,070
Delaware	346	.	*	70	*	.	10	156	587
District of Columbia	740	.	*	27	*	*	*	179	957
Florida	6,517	.	27	711	218	11	88	2,673	10,245
Georgia	3,688	.	18	471	125	13	47	1,359	5,721
Hawaii	485	.	.	33	*	.	*	230	755
Idaho	193	.	*	57	*	*	*	184	449
Illinois	4,939	*	16	643	122	*	194	2,797	8,718
Indiana	1,841	.	*	535	29	*	35	1,149	3,595
Iowa	728	*	18	149	27	*	17	670	1,618
Kansas	671	.	*	173	27	.	*	478	1,355
Kentucky	1,124	.	*	214	46	*	26	896	2,316
Louisiana	2,392	*	*	292	67	*	17	744	3,518
Maine	261	.	.	39	15	.	*	275	597
Maryland	2,278	*	12	329	40	*	34	1,105	3,807
Massachusetts	2,084	.	*	275	44	10	39	1,235	3,690
Michigan	3,181	*	16	796	68	*	45	2,045	6,160
Minnesota	1,164	.	.	109	13	.	29	1,139	2,454
Mississippi	1,670	.	41	132	29	*	*	424	2,308
Missouri	1,861	*	17	335	91	*	17	1,167	3,496
Montana	166	.	.	37	*	*	*	160	373
Nebraska	291	.	*	180	17	.	*	318	814
Nevada	363	.	*	72	16	*	12	283	751
New Hampshire	239	.	.	45	23	*	*	195	510
New Jersey	3,413	12	41	601	188	38	162	1,339	5,794
New Mexico	412	*	.	107	15	*	14	324	874
New York	7,523	11	38	1,264	181	34	114	3,043	12,208
North Carolina	3,298	*	14	638	217	35	63	1,199	5,465
North Dakota	133	.	.	18	*	.	*	168	328
Ohio	3,624	.	*	700	113	28	76	2,542	7,090
Oklahoma	850	*	*	164	12	*	12	616	1,664
Oregon	717	.	*	206	27	*	40	562	1,558
Pennsylvania	4,874	*	10	554	86	44	143	2,806	8,518
Rhode Island	408	.	.	40	12	.	*	185	648
South Carolina	1,991	*	12	233	111	*	63	749	3,166
South Dakota	144	.	*	16	*	.	*	147	323
Tennessee	2,164	*	51	275	59	*	31	1,157	3,740
Texas	7,356	*	17	711	291	50	237	3,340	12,005
Utah	293	.	.	34	*	*	12	371	718
Vermont	105	.	.	29	*	*	.	102	242
Virginia	2,902	*	15	459	54	*	42	1,263	4,745
Washington	1,291	*	53	149	28	23	28	949	2,530
West Virginia	482	*	*	141	21	*	13	381	1,044
Wisconsin	1,214	10	11	287	40	*	47	1,346	2,957
Wyoming	54	.	*	30	.	.	*	79	168
American Samoa	19	.	.	*	.	.	.	.	20
Guam	40	.	.	13	.	.	*	15	69
Puerto Rico	1,148	.	17	185	36	*	32	371	1,793
Saipan	14	.	.	.	.	.	.	*	16
Virgin Islands	49	.	.	*	.	.	.	*	63
Foreign	40	.	*	*	.	.	*	185	239
Unknown	862	.	*	58	*	*	759	1,856	3,554
All	98,625	118	545	15,021	3,176	427	3,338	54,715	175,965

The point prevalent cohorts and associated modalities are determined on December 31 of each calendar year without applying the 60-day stable modality rule.

\* Values for cells with fewer than 10 patients are suppressed.

." Zero values in this cell.

On our website: by Modality, Diabetic Status, &amp; State/Territory (D.h).

Table D.12 (continued)

**Prevalent ESRD Patients for Selected Years, by Modality, Diabetic Status, & State/Territory***point prevalent patients on December 31*

1997	Center	Center	Home				Uncertain		
ALL	hemo	self hemo	hemo	CAPD	CCPD	Other PD	dialysis	Transplant	All
Alabama	4,371	.	16	375	206	*	38	1,648	6,660
Alaska	318	*	*	36	*	*	*	166	539
Arizona	3,489	15	46	207	175	*	76	1,267	5,280
Arkansas	1,891	*	17	204	61	*	22	831	3,030
California	21,881	29	36	1,878	1,083	37	1,140	9,669	35,753
Colorado	1,708	.	14	157	118	*	37	1,111	3,150
Connecticut	2,127	*	*	329	245	*	55	1,008	3,778
Delaware	664	.	*	36	46	.	15	294	1,057
District of Columbia	1,338	.	*	28	18	*	13	234	1,639
Florida	12,048	26	315	701	601	22	308	4,479	18,500
Georgia	7,627	*	115	515	329	15	104	2,130	10,840
Hawaii	1,295	.	*	80	23	*	21	333	1,758
Idaho	424	.	10	89	36	.	12	355	926
Illinois	9,160	*	97	744	439	11	247	4,598	15,299
Indiana	3,719	.	22	598	218	*	116	1,814	6,496
Iowa	1,332	.	24	161	87	.	35	1,110	2,749
Kansas	1,425	.	17	263	86	.	19	721	2,531
Kentucky	2,244	.	*	199	128	16	91	1,344	4,026
Louisiana	5,076	.	17	232	194	*	37	1,288	6,847
Maine	613	.	.	36	40	*	18	393	1,101
Maryland	4,786	15	46	369	227	11	86	1,919	7,459
Massachusetts	3,724	*	44	229	222	10	77	2,015	6,324
Michigan	7,022	.	37	1,008	266	16	144	3,475	11,968
Minnesota	2,313	.	*	131	90	.	64	2,183	4,786
Mississippi	3,368	.	74	190	91	*	26	696	4,452
Missouri	3,732	*	62	426	260	*	50	1,870	6,403
Montana	397	*	*	39	13	*	*	331	791
Nebraska	715	*	*	252	77	*	23	565	1,637
Nevada	888	.	*	88	46	.	28	508	1,565
New Hampshire	407	.	*	71	33	.	14	322	849
New Jersey	6,571	69	58	537	408	41	279	2,355	10,318
New Mexico	1,305	*	*	153	97	*	13	556	2,131
New York	14,848	207	134	1,266	581	76	286	4,914	22,312
North Carolina	7,084	*	41	698	458	96	95	2,022	10,495
North Dakota	284	*	.	37	*	.	*	266	604
Ohio	7,628	*	52	937	444	33	213	4,054	13,364
Oklahoma	2,128	*	*	227	101	17	33	959	3,479
Oregon	1,349	*	48	230	96	.	52	988	2,764
Pennsylvania	9,933	38	51	559	387	14	266	4,480	15,728
Rhode Island	689	.	.	41	26	.	14	316	1,086
South Carolina	4,180	*	27	246	257	*	96	1,268	6,079
South Dakota	452	.	*	43	27	.	12	251	787
Tennessee	4,453	*	37	268	221	*	64	1,738	6,789
Texas	16,559	24	68	832	765	33	525	5,594	24,400
Utah	605	.	*	59	26	*	11	738	1,444
Vermont	231	.	*	31	23	.	*	200	495
Virginia	5,925	*	54	436	298	17	137	2,109	8,977
Washington	2,554	16	120	249	119	*	34	1,655	4,749
West Virginia	1,102	*	*	119	95	*	37	634	2,003
Wisconsin	2,567	.	23	322	179	*	97	2,460	5,652
Wyoming	151	*	*	26	11	*	*	135	337
American Samoa	40	.	.	.	.	.	.	.	40
Guam	177	.	.	19	.	.	.	17	213
Puerto Rico	2,374	*	11	340	294	13	30	479	3,543
Saipan	48	.	.	.	.	.	.	*	52
Virgin Islands	113	.	.	*	*	.	*	11	133
Foreign	61	*	*	*	.	.	*	288	361
Unknown	1,355	.	*	86	24	11	1,111	1,683	4,279
All	204,868	485	1,831	17,439	10,411	566	6,356	88,851	330,807

The point prevalent cohorts and associated modalities are determined on December 31 of each calendar year without applying the 60-day stable modality rule.

\* Values for cells with fewer than 10 patients are suppressed.

." Zero values in this cell.

On our website: by Modality, Diabetic Status, &amp; State/Territory (D.h).



Table D.12 (continued)

**Prevalent ESRD Patients for Selected Years, by Modality, Diabetic Status, & State/Territory***point prevalent patients on December 31*

<b>1997</b>	<b>Center</b>	<b>Center</b>	<b>Home</b>				<b>Uncertain</b>			
<b>DIABETIC</b>	<b>hemo</b>	<b>self hemo</b>	<b>hemo</b>	<b>CAPD</b>	<b>CCPD</b>	<b>Other PD</b>	<b>dialysis</b>	<b>Transplant</b>	<b>All</b>	
Alabama	1,555	.	*	110	65	.	*	11	241	1,985
Alaska	127	.	.	13	.	.	*	.	35	177
Arizona	1,829	*	24	89	65	*	*	34	302	2,348
Arkansas	644	.	*	83	18	.	.	*	170	926
California	9,053	*	*	670	308	13	453	1,664	12,175	
Colorado	741	.	*	60	37	*	*	.	309	1,159
Connecticut	788	*	.	131	81	*	*	20	176	1,200
Delaware	236	.	*	*	*	.	.	*	62	317
District of Columbia	418	.	*	*	*	.	.	*	29	459
Florida	3,959	*	122	221	202	11	88	834	5,442	
Georgia	2,731	*	36	168	107	*	*	30	378	3,455
Hawaii	691	.	*	37	*	*	*	.	35	781
Idaho	168	.	*	42	12	.	.	*	92	323
Illinois	3,061	.	33	236	153	*	*	63	901	4,452
Indiana	1,312	.	*	207	66	*	*	40	339	1,973
Iowa	496	.	*	67	28	.	.	12	259	866
Kansas	527	.	*	79	36	.	*	.	154	808
Kentucky	807	.	*	75	42	*	*	34	219	1,184
Louisiana	1,906	.	*	58	41	*	*	.	260	2,279
Maine	203	.	.	10	10	*	*	.	77	309
Maryland	1,670	*	15	103	59	*	*	22	408	2,282
Massachusetts	1,271	.	*	59	77	*	*	28	366	1,814
Michigan	2,728	.	*	385	71	*	*	57	805	4,060
Minnesota	792	.	.	45	32	.	.	17	636	1,522
Mississippi	1,194	.	11	59	29	*	.	.	98	1,394
Missouri	1,427	.	27	164	84	.	*	*	373	2,083
Montana	174	*	.	15	*	*	*	.	113	310
Nebraska	280	.	*	89	33	*	*	.	196	606
Nevada	321	.	*	39	23	.	.	14	109	507
New Hampshire	135	.	.	25	10	.	.	*	68	244
New Jersey	2,335	22	*	182	133	*	103	443	3,233	
New Mexico	763	.	.	73	46	.	*	.	140	1,031
New York	5,148	47	24	410	174	23	61	796	6,683	
North Carolina	2,667	.	10	200	144	41	40	408	3,510	
North Dakota	120	.	.	16	*	.	*	.	72	215
Ohio	2,897	*	17	390	166	15	70	885	4,442	
Oklahoma	979	*	*	91	43	10	10	211	1,347	
Oregon	501	.	10	100	31	.	21	237	900	
Pennsylvania	3,533	*	15	184	128	*	*	73	886	4,834
Rhode Island	233	.	.	12	*	.	*	.	52	314
South Carolina	1,490	*	*	59	78	*	*	41	217	1,891
South Dakota	215	.	.	20	13	.	*	.	74	324
Tennessee	1,570	.	*	77	100	*	.	22	291	2,065
Texas	7,379	10	*	327	284	*	225	1,105	9,344	
Utah	270	.	*	27	*	.	*	.	249	558
Vermont	93	.	.	12	*	.	*	.	52	164
Virginia	2,093	.	13	139	75	*	38	355	2,721	
Washington	981	*	31	95	21	.	12	417	1,566	
West Virginia	439	*	*	39	43	*	20	125	672	
Wisconsin	939	.	*	123	53	*	33	702	1,856	
Wyoming	72	.	*	12	*	*	*	.	38	133
American Samoa	17	.	.	.	.	.	.	.	.	17
Guam	106	.	.	*	.	.	.	*	.	115
Puerto Rico	1,068	.	*	133	130	*	*	41	1,386	
Saipan	28	.	.	.	.	.	.	.	.	28
Virgin Islands	44	.	.	.	*	.	*	.	.	46
Foreign	14	*	.	.	.	.	*	*	.	23
Unknown	327	.	*	15	*	*	51	14	411	
All	77,565	123	485	6,096	3,401	205	1,868	17,526	107,269	

The point prevalent cohorts and associated modalities are determined on December 31 of each calendar year without applying the 60-day stable modality rule.

\* Values for cells with fewer than 10 patients are suppressed.

." Zero values in this cell.

On our website: by Modality, Diabetic Status, &amp; State/Territory (D.h).

Table D.12 (continued)

**Prevalent ESRD Patients for Selected Years, by Modality, Diabetic Status, & State/Territory***point prevalent patients on December 31*

1997	Center	Center	Home				Uncertain		
NON-DIABETIC	hemo	self hemo	hemo	CAPD	CCPD	Other PD	dialysis	Transplant	All
Alabama	2,816	.	14	265	141	*	27	1,407	4,675
Alaska	191	*	*	23	*	*	*	131	362
Arizona	1,660	14	22	118	110	*	42	965	2,932
Arkansas	1,247	*	13	121	43	*	15	661	2,104
California	12,828	20	31	1,208	775	24	687	8,005	23,578
Colorado	967	.	13	97	81	*	28	802	1,991
Connecticut	1,339	.	*	198	164	*	35	832	2,578
Delaware	428	.	*	28	40	.	11	232	740
District of Columbia	920	.	*	23	16	*	10	205	1,180
Florida	8,089	21	193	480	399	11	220	3,645	13,058
Georgia	4,896	*	79	347	222	12	74	1,752	7,385
Hawaii	604	.	*	43	16	.	13	298	977
Idaho	256	.	*	47	24	.	*	263	603
Illinois	6,099	*	64	508	286	*	184	3,697	10,847
Indiana	2,407	.	17	391	152	*	76	1,475	4,523
Iowa	836	.	20	94	59	.	23	851	1,883
Kansas	898	.	13	184	50	.	11	567	1,723
Kentucky	1,437	.	*	124	86	10	57	1,125	2,842
Louisiana	3,170	.	13	174	153	*	28	1,028	4,568
Maine	410	.	.	26	30	.	10	316	792
Maryland	3,116	14	31	266	168	*	64	1,511	5,177
Massachusetts	2,453	*	36	170	145	*	49	1,649	4,510
Michigan	4,294	.	28	623	195	11	87	2,670	7,908
Minnesota	1,521	.	*	86	58	.	47	1,547	3,264
Mississippi	2,174	.	63	131	62	*	26	598	3,058
Missouri	2,305	*	35	262	176	*	42	1,497	4,320
Montana	223	.	*	24	*	*	*	218	481
Nebraska	435	*	*	163	44	.	17	369	1,031
Nevada	567	.	*	49	23	.	14	399	1,058
New Hampshire	272	.	*	46	23	.	*	254	605
New Jersey	4,236	47	52	355	275	32	176	1,912	7,085
New Mexico	542	*	*	80	51	*	*	416	1,100
New York	9,700	160	110	856	407	53	225	4,118	15,629
North Carolina	4,417	*	31	498	314	55	55	1,614	6,985
North Dakota	164	*	.	21	*	.	*	194	389
Ohio	4,731	*	35	547	278	18	143	3,169	8,922
Oklahoma	1,149	*	*	136	58	*	23	748	2,132
Oregon	848	*	38	130	65	.	31	751	1,864
Pennsylvania	6,400	29	36	375	259	*	193	3,594	10,894
Rhode Island	456	.	.	29	18	.	*	264	772
South Carolina	2,690	*	23	187	179	*	55	1,051	4,188
South Dakota	237	.	*	23	14	.	10	177	463
Tennessee	2,883	*	32	191	121	*	42	1,447	4,724
Texas	9,180	14	59	505	481	28	300	4,489	15,056
Utah	335	.	*	32	19	*	*	489	886
Vermont	138	.	*	19	18	.	*	148	331
Virginia	3,832	*	41	297	223	*	99	1,754	6,256
Washington	1,573	*	89	154	98	*	22	1,238	3,183
West Virginia	663	*	*	80	52	*	17	509	1,331
Wisconsin	1,628	.	18	199	126	*	64	1,758	3,796
Wyoming	79	*	*	14	*	.	*	97	204
American Samoa	23	.	.	.	.	.	.	.	23
Guam	71	.	.	11	.	.	.	16	98
Puerto Rico	1,306	*	10	207	164	*	23	438	2,157
Saipan	20	.	.	.	.	.	.	*	24
Virgin Islands	69	.	.	*	*	.	*	11	87
Foreign	47	.	*	*	.	.	*	281	338
Unknown	1,028	.	*	71	22	10	1,060	1,669	3,868
All	127,303	362	1,346	11,343	7,010	361	4,488	71,325	223,538

The point prevalent cohorts and associated modalities are determined on December 31 of each calendar year without applying the 60-day stable modality rule.

\* Values for cells with fewer than 10 patients are suppressed.

." Zero values in this cell.

On our website: by Modality, Diabetic Status, &amp; State/Territory (D.h).

Table D.12 (continued)

**Prevalent ESRD Patients for Selected Years, by Modality, Diabetic Status, & State/Territory***point prevalent patients on December 31*

2001	Center	Center	Home				Uncertain		
All	hemo	self hemo	hemo	CAPD	CCPD	Other PD	dialysis	Transplant	All
Alabama	5,457	*	18	219	289	*	21	2,008	8,022
Alaska	233	*	*	50	16	.	*	209	515
Arizona	4,615	*	24	185	187	.	34	1,739	6,789
Arkansas	2,166	63	*	217	92	*	22	970	3,541
California	29,861	24	46	1,578	1,280	24	481	12,479	45,773
Colorado	2,278	.	14	93	86	*	19	1,436	3,929
Connecticut	2,592	.	*	248	254	*	13	1,256	4,373
Delaware	810	.	*	18	36	.	*	410	1,281
District of Columbia	1,608	.	*	20	22	.	11	301	1,964
Florida	14,971	*	112	464	718	19	125	5,752	22,167
Georgia	9,984	*	66	402	413	14	68	2,729	13,678
Hawaii	1,585	.	*	70	42	.	16	454	2,171
Idaho	555	*	*	111	39	.	*	462	1,177
Illinois	11,621	*	109	530	506	*	105	5,781	18,660
Indiana	5,035	.	21	416	255	*	31	2,338	8,099
Iowa	1,812	*	17	126	72	.	26	1,397	3,451
Kansas	1,797	.	19	147	98	*	27	890	2,981
Kentucky	3,103	.	12	153	147	*	33	1,546	5,001
Louisiana	6,497	.	*	163	204	*	35	1,602	8,515
Maine	749	.	*	23	52	.	*	514	1,346
Maryland	5,777	.	22	279	271	13	49	2,869	9,280
Massachusetts	4,193	*	19	174	210	19	32	2,565	7,213
Michigan	9,080	*	14	684	333	*	53	4,390	14,560
Minnesota	2,810	.	*	83	95	*	28	2,858	5,877
Mississippi	4,070	*	33	179	210	*	14	918	5,437
Missouri	4,968	*	37	308	285	*	16	2,286	7,903
Montana	505	.	10	35	29	.	*	410	997
Nebraska	1,114	.	*	123	87	*	14	708	2,052
Nevada	1,271	.	*	80	73	*	32	664	2,124
New Hampshire	558	.	*	64	31	.	*	421	1,081
New Jersey	8,577	12	24	269	403	10	107	3,266	12,668
New Mexico	1,807	.	*	102	123	.	10	683	2,730
New York	19,176	181	79	682	623	49	135	6,667	27,592
North Carolina	9,398	.	27	349	492	119	55	2,637	13,077
North Dakota	352	.	.	26	*	.	*	362	751
Ohio	10,481	*	33	643	535	*	130	4,861	16,693
Oklahoma	2,798	*	*	165	181	12	28	1,157	4,346
Oregon	1,841	.	17	129	119	*	32	1,325	3,465
Pennsylvania	11,729	*	25	334	435	17	106	5,725	18,375
Rhode Island	783	.	.	21	22	*	11	439	1,277
South Carolina	5,219	*	*	161	356	*	60	1,642	7,451
South Dakota	572	.	*	29	36	.	*	334	974
Tennessee	5,498	.	26	204	333	11	33	2,119	8,224
Texas	22,199	*	51	610	1,007	34	383	7,242	31,528
Utah	871	.	*	58	28	*	*	957	1,924
Vermont	353	.	*	*	15	.	*	228	607
Virginia	7,387	.	51	321	355	11	54	2,818	10,997
Washington	3,395	*	71	166	134	10	29	2,252	6,066
West Virginia	1,479	*	*	79	171	*	*	832	2,572
Wisconsin	3,395	.	20	185	181	*	45	3,021	6,853
Wyoming	220	.	*	11	20	.	.	171	426
American Samoa	66	.	.	.	.	.	*	.	67
Guam	248	.	.	14	.	.	*	19	285
Puerto Rico	2,924	.	*	92	306	*	14	759	4,104
Saipan	70	.	.	.	.	.	*	10	85
Virgin Islands	134	.	.	*	*	.	.	21	160
Foreign	43	.	.	*	*	.	*	343	397
Unknown	565	.	*	27	21	.	197	1,614	2,430
All	263,255	333	1,122	11,927	12,341	445	2,792	113,866	406,081

The point prevalent cohorts and associated modalities are determined on December 31 of each calendar year without applying the 60-day stable modality rule.

\* Values for cells with fewer than 10 patients are suppressed.

." Zero values in this cell.

On our website: by Modality, Diabetic Status, &amp; State/Territory (D.h).

Table D.12 (continued)

**Prevalent ESRD Patients for Selected Years, by Modality, Diabetic Status, & State/Territory***point prevalent patients on December 31*

2001 DIABETIC	Center hemo	Center self hemo	Home hemo	CAPD	CCPD	Other PD	Uncertain dialysis	Transplant	All
Alabama	2,281	.	*	61	78	*	14	332	2,772
Alaska	101	.	*	17	*	.	.	43	170
Arizona	2,525	.	*	73	69	.	13	444	3,132
Arkansas	811	25	*	93	29	.	*	211	1,179
California	13,346	*	*	565	406	14	173	2,282	16,800
Colorado	1,057	.	*	40	23	*	*	387	1,521
Connecticut	991	.	*	103	85	.	.	217	1,398
Delaware	330	.	.	*	*	.	.	101	440
District of Columbia	502	.	*	*	*	.	*	52	571
Florida	5,654	*	28	149	224	*	39	1,150	7,253
Georgia	3,895	.	17	114	114	*	21	543	4,709
Hawaii	926	.	.	33	24	.	*	70	1,061
Idaho	248	*	*	41	19	.	.	122	433
Illinois	4,316	.	38	174	173	*	18	1,163	5,885
Indiana	2,003	.	10	137	88	*	*	456	2,703
Iowa	718	.	*	56	23	.	*	343	1,149
Kansas	794	.	*	62	29	*	*	193	1,087
Kentucky	1,306	.	*	56	50	*	*	280	1,707
Louisiana	2,569	.	*	48	50	*	*	345	3,019
Maine	320	.	.	*	14	.	*	103	443
Maryland	2,117	.	*	86	83	*	*	608	2,911
Massachusetts	1,574	.	*	56	67	*	*	485	2,196
Michigan	3,796	.	*	272	119	*	14	1,046	5,254
Minnesota	1,038	.	.	23	23	*	*	805	1,899
Mississippi	1,549	.	*	67	77	*	*	148	1,854
Missouri	2,007	.	*	122	99	.	*	473	2,712
Montana	246	.	*	18	13	.	*	147	426
Nebraska	475	.	.	51	32	.	*	227	788
Nevada	538	.	*	36	27	*	13	149	766
New Hampshire	186	.	.	24	*	.	.	93	312
New Jersey	3,345	*	*	87	124	*	34	646	4,244
New Mexico	1,114	.	.	52	61	.	*	213	1,446
New York	7,275	51	17	235	189	20	34	1,138	8,959
North Carolina	3,774	.	*	111	133	50	17	575	4,668
North Dakota	145	.	.	12	*	.	*	101	264
Ohio	4,522	*	*	276	200	*	42	1,119	6,172
Oklahoma	1,348	.	.	79	82	*	13	266	1,794
Oregon	683	.	*	57	48	*	15	316	1,126
Pennsylvania	4,592	.	*	121	131	*	20	1,165	6,039
Rhode Island	276	.	.	*	*	*	*	84	378
South Carolina	2,165	.	.	54	118	*	28	307	2,673
South Dakota	294	.	.	13	10	.	*	95	413
Tennessee	2,127	.	*	51	134	*	*	372	2,700
Texas	10,758	*	15	239	364	*	169	1,594	13,143
Utah	417	.	*	25	*	.	.	302	753
Vermont	146	.	*	.	*	.	.	53	207
Virginia	2,940	.	13	110	115	*	14	525	3,719
Washington	1,408	*	18	53	44	*	*	546	2,083
West Virginia	709	*	*	37	85	.	*	182	1,017
Wisconsin	1,260	.	*	59	55	*	23	841	2,241
Wyoming	99	.	*	*	*	.	.	52	167
American Samoa	38	.	.	.	.	.	*	.	39
Guam	150	.	.	*	.	.	*	*	157
Puerto Rico	1,504	.	*	39	145	*	*	89	1,785
Saipan	55	.	.	.	.	.	*	*	59
Virgin Islands	62	.	.	*	*	.	.	*	67
Foreign	11	.	.	.	*	.	.	*	20
Unknown	59	.	.	*	*	.	*	13	80
All	109,495	95	270	4,319	4,145	170	844	23,625	142,963

The point prevalent cohorts and associated modalities are determined on December 31 of each calendar year without applying the 60-day stable modality rule.

\* Values for cells with fewer than 10 patients are suppressed.

." Zero values in this cell.

On our website: by Modality, Diabetic Status, &amp; State/Territory (D.h).

Table D.12 (continued)

**Prevalent ESRD Patients for Selected Years, by Modality, Diabetic Status, & State/Territory***point prevalent patients on December 31*

2001 NON-DIABETIC	Center hemo	Center self hemo	Home hemo	CAPD	CCPD	Other PD	Uncertain dialysis	Transplant	All
Alabama	3,176	*	14	158	211	*	*	1,676	5,250
Alaska	132	*	*	33	*	.	*	166	345
Arizona	2,090	*	16	112	118	.	21	1,295	3,657
Arkansas	1,355	38	*	124	63	*	13	759	2,362
California	16,515	19	37	1,013	874	10	308	10,197	28,973
Colorado	1,221	.	*	53	63	*	11	1,049	2,408
Connecticut	1,601	.	*	145	169	*	13	1,039	2,975
Delaware	480	.	*	17	28	.	*	309	841
District of Columbia	1,106	.	*	13	15	.	*	249	1,393
Florida	9,317	*	84	315	494	11	86	4,602	14,914
Georgia	6,089	*	49	288	299	*	47	2,186	8,969
Hawaii	659	.	*	37	18	.	*	384	1,110
Idaho	307	.	*	70	20	.	*	340	744
Illinois	7,305	*	71	356	333	*	87	4,618	12,775
Indiana	3,032	.	11	279	167	*	23	1,882	5,396
Iowa	1,094	*	12	70	49	.	22	1,054	2,302
Kansas	1,003	.	16	85	69	*	23	697	1,894
Kentucky	1,797	.	*	97	97	*	25	1,266	3,294
Louisiana	3,928	.	*	115	154	*	31	1,257	5,496
Maine	429	.	*	18	38	.	*	411	903
Maryland	3,660	.	15	193	188	10	42	2,261	6,369
Massachusetts	2,619	*	14	118	143	11	31	2,080	5,017
Michigan	5,284	*	*	412	214	*	39	3,344	9,306
Minnesota	1,772	.	*	60	72	.	19	2,053	3,978
Mississippi	2,521	*	29	112	133	*	*	770	3,583
Missouri	2,961	*	29	186	186	*	13	1,813	5,191
Montana	259	.	*	17	16	.	*	263	571
Nebraska	639	.	*	72	55	*	11	481	1,264
Nevada	733	.	*	44	46	.	19	515	1,358
New Hampshire	372	.	*	40	22	.	*	328	769
New Jersey	5,232	10	23	182	279	*	73	2,620	8,424
New Mexico	693	.	*	50	62	.	*	470	1,284
New York	11,901	130	62	447	434	29	101	5,529	18,633
North Carolina	5,624	.	19	238	359	69	38	2,062	8,409
North Dakota	207	.	.	14	*	.	.	261	487
Ohio	5,959	*	26	367	335	*	88	3,742	10,521
Oklahoma	1,450	*	*	86	99	*	15	891	2,552
Oregon	1,158	.	11	72	71	*	17	1,009	2,339
Pennsylvania	7,137	*	21	213	304	11	86	4,560	12,336
Rhode Island	507	.	.	15	14	.	*	355	899
South Carolina	3,054	*	*	107	238	*	32	1,335	4,778
South Dakota	278	.	*	16	26	.	*	239	561
Tennessee	3,371	.	25	153	199	*	24	1,747	5,524
Texas	11,441	*	36	371	643	31	214	5,648	18,385
Utah	454	.	*	33	20	*	*	655	1,171
Vermont	207	.	*	*	*	.	*	175	400
Virginia	4,447	.	38	211	240	*	40	2,293	7,278
Washington	1,987	*	53	113	90	*	23	1,706	3,983
West Virginia	770	.	*	42	86	*	*	650	1,555
Wisconsin	2,135	.	18	126	126	*	22	2,180	4,612
Wyoming	121	.	*	*	12	.	.	119	259
American Samoa	28	.	.	.	.	.	.	.	28
Guam	98	.	.	10	.	.	*	18	128
Puerto Rico	1,420	.	*	53	161	*	*	670	2,319
Saipan	15	.	.	.	.	.	*	*	26
Virgin Islands	72	.	.	*	*	.	.	18	93
Foreign	32	.	.	*	.	.	*	335	377
Unknown	506	.	*	25	20	.	192	1,601	2,350
All	153,760	238	852	7,608	8,196	275	1,948	90,241	263,118

The point prevalent cohorts and associated modalities are determined on December 31 of each calendar year without applying the 60-day stable modality rule.

\* Values for cells with fewer than 10 patients are suppressed.

." Zero values in this cell.

On our website: by Modality, Diabetic Status, &amp; State/Territory (D.h).

Table D.13

**Prevalent Dialysis Patients, by Modality, Age, & State/Territory***point prevalent dialysis patients on December 31, 1999-2001 combined, by state, age, & modality*

	Center hemo	Center self hemo	Home hemo	CAPD	CCPD	Other PD	Uncertain dialysis	All
Alabama	15,598	*	52	730	801	20	84	17,290
Alaska	673	*	10	129	36	.	*	852
Arizona	13,116	27	84	591	490	12	155	14,475
Arkansas	6,396	64	24	611	237	13	76	7,421
California	83,585	70	282	4,749	3,584	75	2,125	94,470
Colorado	6,555	*	37	294	294	*	68	7,255
Connecticut	7,486	*	22	853	751	13	108	9,234
Delaware	2,326	.	*	54	119	.	28	2,533
District of Columbia	4,616	.	*	76	76	*	40	4,818
Florida	42,888	29	389	1,464	2,042	63	649	47,524
Georgia	28,153	*	310	1,287	1,137	36	326	31,251
Hawaii	4,482	*	32	229	111	.	101	4,956
Idaho	1,571	*	24	294	126	.	23	2,041
Illinois	33,571	*	348	1,750	1,516	29	420	37,640
Indiana	14,431	.	49	1,321	737	21	188	16,747
Iowa	5,091	*	57	417	254	.	101	5,922
Kansas	5,218	.	50	529	274	*	113	6,191
Kentucky	8,659	.	23	429	423	35	200	9,769
Louisiana	18,782	*	47	511	608	11	106	20,066
Maine	2,158	.	12	78	127	*	33	2,410
Maryland	16,827	*	53	893	729	29	202	18,734
Massachusetts	12,309	*	67	563	581	46	196	13,763
Michigan	25,810	*	48	2,225	918	11	268	29,283
Minnesota	7,995	.	11	251	301	*	135	8,697
Mississippi	11,772	10	126	524	528	25	62	13,047
Missouri	14,210	*	147	886	839	*	89	16,181
Montana	1,437	.	26	110	68	*	11	1,655
Nebraska	2,977	*	11	480	320	*	46	3,836
Nevada	3,543	.	13	233	182	*	122	4,096
New Hampshire	1,597	*	*	169	87	.	31	1,892
New Jersey	24,766	61	86	932	1,250	35	519	27,649
New Mexico	5,076	*	13	309	365	.	53	5,817
New York	55,163	567	259	2,444	1,885	146	507	60,971
North Carolina	26,423	*	70	1,253	1,491	335	248	29,822
North Dakota	942	.	.	96	35	.	33	1,106
Ohio	29,729	*	71	2,133	1,541	29	445	33,956
Oklahoma	7,981	*	12	537	492	50	97	9,172
Oregon	5,137	.	58	450	345	*	131	6,125
Pennsylvania	34,410	15	83	1,170	1,286	41	540	37,545
Rhode Island	2,323	.	.	85	52	*	41	2,502
South Carolina	15,038	*	42	537	1,003	11	238	16,874
South Dakota	1,676	.	*	95	97	.	22	1,893
Tennessee	16,007	.	87	641	905	34	121	17,795
Texas	62,585	27	202	1,962	2,743	102	1,354	68,975
Utah	2,383	.	15	181	71	*	42	2,694
Vermont	965	.	16	28	53	.	21	1,083
Virginia	21,301	*	140	1,082	1,013	28	259	23,825
Washington	9,564	30	247	553	400	23	97	10,914
West Virginia	4,122	*	13	260	440	*	66	4,910
Wisconsin	9,694	.	61	638	507	12	244	11,156
Wyoming	635	*	11	46	54	.	*	750
American Samoa	171	.	*	.	.	.	*	174
Guam	695	.	*	41	*	.	*	747
Puerto Rico	8,406	.	20	426	995	*	60	9,913
Saipan	211	.	.	.	.	.	12	223
Virgin Islands	394	.	.	*	*	.	.	408
Foreign	157	.	.	*	*	.	29	196
Unknown	1,984	.	17	114	74	*	828	3,020
All	751,770	958	3,904	38,757	35,405	1,344	12,126	844,264

The point prevalent cohorts and associated modalities are determined on December 31 of each calendar year without applying the 60-day stable modality rule.

The age of the point prevalent patient is determined as of December 31 of each calendar year.

Patients with unknown age are excluded.

\* Values for cells with fewer than 10 patients are suppressed.

." Zero values in this cell.

Table D.13 (continued)

**Prevalent Dialysis Patients, by Modality, Age, & State/Territory***point prevalent dialysis patients on December 31, 1999-2001 combined, by state, age, & modality*

<b>UNDER 65</b>	<b>Center hemo</b>	<b>Center self hemo</b>	<b>Home hemo</b>	<b>CAPD</b>	<b>CCPD</b>	<b>Other PD</b>	<b>Uncertain dialysis</b>	<b>All</b>
Alabama	9,091	*	41	568	587	17	41	10,350
Alaska	491	*	*	94	27	.	*	623
Arizona	7,117	25	49	443	340	*	72	8,055
Arkansas	3,751	29	11	449	178	*	46	4,471
California	45,004	51	174	3,454	2,644	47	1,171	52,545
Colorado	3,647	*	29	211	208	*	53	4,152
Connecticut	3,472	.	17	571	499	10	48	4,617
Delaware	1,274	.	*	40	86	.	17	1,423
District of Columbia	2,873	.	*	61	64	*	20	3,026
Florida	22,287	21	214	1,072	1,389	39	279	25,301
Georgia	17,598	*	209	1,047	912	34	205	20,007
Hawaii	2,433	.	14	180	74	.	42	2,743
Idaho	826	*	17	181	101	.	13	1,140
Illinois	17,829	*	148	1,213	992	24	250	20,459
Indiana	7,353	.	29	932	471	17	88	8,890
Iowa	2,260	*	39	211	151	.	60	2,723
Kansas	2,654	.	41	346	167	*	69	3,282
Kentucky	4,705	.	21	328	308	27	95	5,484
Louisiana	11,454	.	29	406	450	10	78	12,427
Maine	950	.	12	45	92	.	18	1,117
Maryland	9,439	*	43	625	493	26	134	10,761
Massachusetts	5,305	*	38	377	367	17	87	6,192
Michigan	13,827	*	35	1,559	623	*	152	16,207
Minnesota	3,477	.	10	121	179	*	91	3,881
Mississippi	7,275	*	118	405	411	25	40	8,283
Missouri	7,355	*	75	599	573	*	52	8,660
Montana	751	.	14	67	42	*	*	883
Nebraska	1,530	*	11	251	162	*	31	1,987
Nevada	1,999	.	12	166	128	*	87	2,394
New Hampshire	748	*	*	109	55	.	12	929
New Jersey	12,379	50	74	606	809	21	286	14,225
New Mexico	2,833	*	*	236	259	.	31	3,369
New York	29,640	412	171	1,766	1,325	99	333	33,746
North Carolina	15,744	*	62	957	1,093	246	137	18,241
North Dakota	464	.	.	32	23	.	21	540
Ohio	15,014	*	45	1,480	1,004	24	248	17,817
Oklahoma	4,551	*	10	383	367	32	60	5,405
Oregon	2,669	.	29	293	235	*	82	3,312
Pennsylvania	16,452	*	71	800	915	31	260	18,536
Rhode Island	943	.	.	61	40	*	12	1,057
South Carolina	9,145	*	41	458	761	10	154	10,573
South Dakota	816	.	.	69	66	.	12	963
Tennessee	9,306	.	80	471	663	30	68	10,618
Texas	37,387	20	142	1,500	2,013	79	927	42,068
Utah	1,352	.	10	111	46	*	22	1,543
Vermont	404	.	13	21	24	.	*	467
Virginia	11,899	*	100	831	698	19	153	13,702
Washington	5,266	*	144	363	276	17	71	6,145
West Virginia	1,985	*	*	171	284	*	28	2,480
Wisconsin	4,469	.	47	313	265	*	129	5,227
Wyoming	352	.	*	33	45	.	.	439
American Samoa	129	.	*	.	.	.	*	132
Guam	463	.	.	38	.	.	*	509
Puerto Rico	5,364	.	16	319	813	*	40	6,556
Saipan	158	.	.	.	.	.	*	166
Virgin Islands	259	.	.	*	*	.	.	268
Foreign	67	.	.	*	*	.	17	90
Unknown	1,675	.	15	103	62	*	625	2,483
All	409,960	670	2,572	27,553	24,869	969	7,096	473,689

The point prevalent cohorts and associated modalities are determined on December 31 of each calendar year without applying the 60-day stable modality rule.

The age of the point prevalent patient is determined as of December 31 of each calendar year.

Patients with unknown age are excluded.

\* Values for cells with fewer than 10 patients are suppressed.

." Zero values in this cell.

Table D.13 (continued)

**Prevalent Dialysis Patients, by Modality, Age, & State/Territory***point prevalent dialysis patients on December 31, 1999-2001 combined, by state, age, & modality*

65+	Center hemo	Center self hemo	Home hemo	CAPD	CCPD	Other PD	Uncertain dialysis	All
Alabama	6,507	.	11	162	214	*	43	6,940
Alaska	182	*	*	35	*	.	*	229
Arizona	5,999	*	35	148	150	*	83	6,420
Arkansas	2,645	35	13	162	59	*	30	2,950
California	38,581	19	108	1,295	940	28	954	41,925
Colorado	2,908	.	*	83	86	*	15	3,103
Connecticut	4,014	*	*	282	252	*	60	4,617
Delaware	1,052	.	.	14	33	.	11	1,110
District of Columbia	1,743	.	*	15	12	.	20	1,792
Florida	20,601	*	175	392	653	24	370	22,223
Georgia	10,555	.	101	240	225	*	121	11,244
Hawaii	2,049	*	18	49	37	.	59	2,213
Idaho	745	*	*	113	25	.	10	901
Illinois	15,742	*	200	537	524	*	170	17,181
Indiana	7,078	.	20	389	266	*	100	7,857
Iowa	2,831	.	18	206	103	.	41	3,199
Kansas	2,564	.	*	183	107	*	44	2,909
Kentucky	3,954	.	*	101	115	*	105	4,285
Louisiana	7,328	*	18	105	158	*	28	7,639
Maine	1,208	.	.	33	35	*	15	1,293
Maryland	7,388	.	10	268	236	*	68	7,973
Massachusetts	7,004	.	29	186	214	29	109	7,571
Michigan	11,983	*	13	666	295	*	116	13,076
Minnesota	4,518	.	*	130	122	*	44	4,816
Mississippi	4,497	*	*	119	117	.	22	4,764
Missouri	6,855	*	72	287	266	*	37	7,521
Montana	686	.	12	43	26	*	*	772
Nebraska	1,447	.	.	229	158	.	15	1,849
Nevada	1,544	.	*	67	54	*	35	1,702
New Hampshire	849	.	*	60	32	.	19	963
New Jersey	12,387	11	12	326	441	14	233	13,424
New Mexico	2,243	.	*	73	106	.	22	2,448
New York	25,523	155	88	678	560	47	174	27,225
North Carolina	10,679	.	*	296	398	89	111	11,581
North Dakota	478	.	.	64	12	.	12	566
Ohio	14,715	*	26	653	537	*	197	16,139
Oklahoma	3,430	*	*	154	125	18	37	3,767
Oregon	2,468	.	29	157	110	.	49	2,813
Pennsylvania	17,957	*	12	370	371	10	280	19,008
Rhode Island	1,380	.	.	24	12	.	29	1,445
South Carolina	5,892	*	.	79	242	*	84	6,300
South Dakota	860	.	*	26	31	.	10	930
Tennessee	6,701	.	*	170	242	*	53	7,177
Texas	25,198	*	60	462	730	23	427	26,907
Utah	1,031	.	*	70	25	.	20	1,151
Vermont	561	.	*	*	29	.	16	616
Virginia	9,402	.	40	251	315	*	106	10,123
Washington	4,298	22	103	190	124	*	26	4,769
West Virginia	2,137	*	*	89	156	*	38	2,430
Wisconsin	5,225	.	14	325	242	*	115	5,929
Wyoming	283	*	*	13	*	.	*	311
American Samoa	42	.	.	.	.	.	.	42
Guam	232	.	*	*	.	.	*	238
Puerto Rico	3,042	.	*	107	182	*	20	3,357
Saipan	53	.	.	.	.	.	*	57
Virgin Islands	135	.	.	*	*	.	.	140
Foreign	90	.	.	*	.	.	12	106
Unknown	259	.	*	*	10	.	202	482
All	341,758	288	1,332	11,202	10,534	375	5,029	370,518

The point prevalent cohorts and associated modalities are determined on December 31 of each calendar year without applying the 60-day stable modality rule. The age of the point prevalent patient is determined as of December 31 of each calendar year.

Patients with unknown age are excluded.

\* Values for cells with fewer than 10 patients are suppressed.

." Zero values in this cell.