

Reference section C

Patient characteristics



table c.1 · 349
number of medical evidence forms filed
by year of Medicare certification

table c.2 · 350
counts of new ESRD patients, with or without a medical evidence form: by data source
incident patients

table c.3 · 351
counts of new ESRD patients, with or without a medical evidence form: by race
incident patients

table c.4 · 352
patient counts
incident patients, by demographic characteristics & primary diagnosis

table c.5 · 353
patient counts: by ESRD network
incident patients by race & ethnicity, May 1995-April 2003 combined

table c.6 · 354
percent distribution of patients: by race & Hispanic ethnicity
incident patients

table c.7 · 354
percent distribution of diabetic patients: by race & Hispanic ethnicity
incident patients

table c.8 · 354
patient counts: by employment status
incident patients

table c.9 · 355
patient counts: by insurance payor
incident patients

table c.10 · 355
percent distribution of patients: by comorbidity
incident patients

table c.11 · 355
percent distribution of patients: by primary diagnosis
incident patients

table c.12 · 356
lab values & physical characteristics: pediatric & adult patients
incident patients

table c.13 · 357
lab values & physical characteristics
incident patients, by demographic characteristics, May 1995-April 2003 combined

table c.14 · 358
lab values & physical characteristics: by year & age
incident patients, May 1995-April 2003 combined

table c.15 · 359
lab values & physical characteristics: by race
incident patients, by gender, age, & diabetic status, May 1995-April 2003 combined

table c.16 · 362
lab values & physical characteristics: by Hispanic ethnicity
incident patients, by gender, age, & diabetic status



Table C.1
Number of Medical Evidence Forms Filed
by year of Medicare certification

	Form version		For this patient, this form is			All
	Pre-1995	1995 revision	1st form filed	2nd form filed	3rd or later	
1960-1972	355	.	355	.	.	355
1973	1,478	.	1,474	*	.	1,478
1974	3,518	.	3,511	*	.	3,518
1975	4,317	.	4,302	15	.	4,317
1976	5,014	.	5,003	11	.	5,014
1977	7,215	.	7,190	25	.	7,215
1978	7,473	.	7,442	29	*	7,473
1979	8,270	.	8,212	58	.	8,270
1980	10,472	.	10,431	41	.	10,472
1981	14,927	.	14,834	91	*	14,927
1982	20,724	.	20,571	151	*	20,724
1983	22,414	.	22,257	153	*	22,414
1984	25,100	.	24,823	270	*	25,100
1985	28,022	.	27,723	291	*	28,022
1986	30,889	.	30,486	398	*	30,889
1987	34,511	.	34,087	411	13	34,511
1988	37,211	.	36,743	448	20	37,211
1989	43,498	.	42,790	685	23	43,498
1990	49,548	.	48,665	861	22	49,548
1991	54,068	.	53,000	1,032	36	54,068
1992	60,240	.	59,225	978	37	60,240
1993	61,251	.	59,688	1,479	84	61,251
1994	71,207	.	68,261	2,655	291	71,207
1995	19,294	52,764	68,591	3,232	235	72,058
1996	681	79,066	76,237	3,218	292	79,747
1997	660	84,252	81,367	3,297	248	84,912
1998	889	89,638	86,649	3,563	315	90,527
1999	1,165	93,067	90,280	3,627	325	94,232
2000	1,052	96,550	93,344	3,931	327	97,602
2001	951	101,271	97,625	4,232	365	102,222
2002	978	103,018	99,284	4,372	340	103,996
2003 [^]	522	71,707	69,024	2,924	281	72,229
All	627,914	771,333	1,353,474	42,489	3,284	1,399,247

[^]Incomplete data

*Values for cells with 10 or fewer patients are suppressed.

." Zero values in this cell.

Table C.2
Counts of New ESRD Patients, with or without a Medical Evidence Form: by Data Source
incident patients

	Percent with no CMS-2728	Version of CMS 2728			Source of data of 1st ESRD service			*Patient ever been Medicare entitled?		Pts used in remaining C tables	All
		No form	Pre-1995	1995 Revision	CMS-2728	Medicare claims	§Tx record	No	Yes		
1960-1972	31.4	527	1,013	136	650	861	165	.	.		1,676
1973	5.6	202	3,388	36	3,228	307	91	.	.		3,626
1974	4.2	187	4,203	20	4,038	296	76	.	.		4,410
1975	3.7	204	5,228	30	5,008	359	95	.	.		5,462
1976	4.1	256	5,977	36	5,662	517	90	.	.		6,269
1977	58.6	12,580	8,751	135	6,094	15,102	270	.	.		21,466
1978	39.5	5,794	8,794	86	6,294	8,078	302	554	14,120		14,674
1979	39.5	6,534	9,869	121	6,879	9,321	324	619	15,905		16,524
1980	35.0	6,250	11,451	142	8,198	9,331	314	776	17,067		17,843
1981	17.1	3,394	16,394	116	13,241	6,357	306	929	18,975		19,904
1982	7.0	1,575	20,772	90	19,024	2,988	425	1,095	21,342		22,437
1983	11.5	2,995	22,887	81	20,029	5,536	398	1,199	24,764		25,963
1984	6.3	1,714	25,512	110	20,620	6,257	459	1,223	26,113		27,336
1985	5.7	1,717	28,425	152	21,125	8,643	526	1,354	28,940		30,294
1986	6.3	2,081	30,789	190	20,860	11,510	690	1,393	31,667		33,060
1987	5.8	2,112	34,041	248	22,863	12,660	878	1,780	34,621		36,401
1988	7.8	3,170	37,287	332	24,823	14,679	1,287	2,653	38,136		40,789
1989	5.8	2,677	42,888	345	28,310	16,322	1,278	2,822	43,088		45,910
1990	6.3	3,140	46,493	391	32,843	15,897	1,284	3,310	46,714		50,024
1991	4.9	2,697	52,184	423	31,644	22,460	1,200	3,567	51,737		55,304
1992	4.4	2,684	57,274	546	35,234	24,101	1,169	4,148	56,356		60,504
1993	8.7	5,531	57,294	752	35,489	26,794	1,294	5,109	58,468		63,577
1994	5.7	3,972	64,430	1,338	40,435	28,162	1,143	6,789	62,951	405	69,740
1995	3.5	2,536	13,885	55,798	45,876	25,084	1,259	8,009	64,210	54,871	72,219
1996	1.7	1,318	425	76,381	50,927	25,850	1,347	8,787	69,337	74,762	78,124
1997	1.4	1,130	410	80,871	54,794	26,238	1,379	10,085	72,326	79,690	82,411
1998	1.3	1,141	523	85,671	59,666	26,211	1,458	11,050	76,285	84,747	87,335
1999	1.6	1,483	1,119	89,289	63,334	26,831	1,726	12,312	79,579	88,200	91,891
2000	1.5	1,422	924	91,936	64,817	27,574	1,891	12,678	81,604	91,165	94,282
2001	1.3	1,248	832	94,381	66,207	28,316	1,938	12,760	83,701	94,163	96,461
2002	1.3	1,340	927	98,105	66,806	31,547	2,019	13,690	86,682	97,905	100,372
2003^	6.4	4,161	403	60,241	43,170	20,215	1,420	.	.	60,241	64,805
All	6.1	87,772	614,792	738,529	928,188	484,404	28,501	236,405	204,688	726,149	1,441,093

^ Incomplete data.

*Medicare entitlement determined from the payor sequence. See Appendix A.

§ Represents CMS & UNOS transplant records.

." Zero values in this cell.



Table C.3
Counts of New ESRD Patients, with or without a Medical Evidence Form: by Race
incident patients

	White	Black	Nat. Am/ Alas. Native	Asian	Pacific Islander	Mid-East/ Arabian	Indian Sub- continent	Other/ Multiracial	Unknown	No CMS- 2728	All
1960-1972	764	204	15	14	*	.	.	*	143	527	1,676
1973	450	155	*	*	*	.	.	*	2,805	202	3,626
1974	577	195	*	13	*	.	.	.	3,426	187	4,410
1975	685	258	11	*	*	.	*	*	4,286	204	5,462
1976	799	294	15	12	*	.	.	*	4,885	256	6,269
1977	1,637	607	25	28	12	*	.	*	6,570	12,580	21,466
1978	1,636	629	39	34	12	.	.	*	6,529	5,794	14,674
1979	2,077	786	51	36	19	.	.	*	7,015	6,534	16,524
1980	2,674	1,010	88	45	16	*	.	*	7,754	6,250	17,843
1981	6,090	2,477	161	126	22	.	.	*	7,631	3,394	19,904
1982	13,966	5,603	210	310	27	*	.	*	737	1,575	22,437
1983	15,133	6,496	282	326	160	*	*	*	564	2,995	25,963
1984	16,802	7,207	292	380	318	.	*	13	609	1,714	27,336
1985	18,726	7,981	303	496	297	*	*	*	761	1,717	30,294
1986	20,135	8,384	382	511	385	*	.	13	1,167	2,081	33,060
1987	21,932	9,325	439	563	485	*	*	14	1,527	2,112	36,401
1988	23,873	10,244	523	676	353	*	*	20	1,926	3,170	40,789
1989	27,628	12,116	584	841	*	*	*	32	2,019	2,677	45,910
1990	30,257	12,814	633	1,013	*	*	*	34	2,121	3,140	50,024
1991	33,681	14,698	701	1,134	*	*	*	44	2,338	2,697	55,304
1992	36,675	16,326	861	1,293	*	*	*	74	2,576	2,684	60,504
1993	36,281	16,275	753	1,419	*	*	*	82	3,211	5,531	63,577
1994	40,655	18,685	850	1,848	*	14	*	115	3,582	3,972	69,740
1995	42,757	20,693	1,019	1,685	472	222	166	1,602	1,067	2,536	72,219
1996	47,643	22,194	1,236	1,838	706	191	188	2,327	483	1,318	78,124
1997	50,721	23,640	911	1,798	701	233	218	2,670	389	1,130	82,411
1998	53,631	25,221	1,135	2,239	801	282	242	2,302	341	1,141	87,335
1999	56,572	25,601	1,423	2,343	752	302	264	2,367	784	1,483	91,891
2000	57,725	25,792	2,128	2,404	800	377	295	2,793	546	1,422	94,282
2001	60,707	26,948	1,104	2,273	868	348	332	2,372	261	1,248	96,461
2002	63,208	28,078	1,090	2,367	933	386	297	2,385	288	1,340	100,372
2003^	38,294	17,390	637	1,457	590	203	226	1,718	129	4,161	64,805
All	824,391	368,326	17,917	29,534	8,793	2,595	2,263	21,032	78,470	87,772	1,441,093

^Incomplete data

*Values for cells with 10 or fewer patients are suppressed.

." Zero values in this cell.

Table C.4
Patient Counts

incident patients by demographic characteristics & primary diagnosis

	1995 [^]	1996	1997	1998	1999	2000	2001	2002	2003 [^]	All
0-4	104	179	187	177	189	197	204	173	69	1,479
5-9	96	137	122	130	145	144	135	158	54	1,121
10-14	168	259	271	263	274	286	288	323	134	2,266
15-19	370	551	511	523	571	544	554	557	243	4,424
20-24	635	1,000	947	992	945	951	1,010	996	429	7,905
25-29	1,033	1,651	1,652	1,687	1,631	1,662	1,618	1,512	603	13,049
30-34	1,582	2,414	2,373	2,396	2,397	2,395	2,315	2,367	949	19,188
35-39	2,066	3,290	3,280	3,302	3,362	3,233	3,281	3,299	1,352	26,465
40-44	2,608	4,231	4,203	4,520	4,504	4,604	4,594	4,629	1,937	35,830
45-49	3,160	5,236	5,365	5,773	5,931	6,067	6,158	6,466	2,828	46,984
50-54	3,526	5,729	6,288	6,881	7,475	7,693	8,289	8,280	3,673	57,834
55-59	4,180	6,782	7,330	7,949	8,393	8,745	9,025	9,570	4,138	66,112
60-64	5,210	8,403	8,810	9,391	9,535	9,837	10,233	10,423	4,576	76,418
65-69	6,116	10,313	10,780	10,985	11,000	11,208	11,462	11,632	4,792	88,288
70-74	6,107	10,689	11,243	11,532	11,814	12,130	12,310	12,510	5,113	93,448
75-79	4,767	8,388	9,260	10,060	10,946	11,228	11,356	12,060	5,060	83,125
80-84	2,768	4,918	5,666	6,043	6,653	7,219	7,460	8,597	3,591	52,915
85+	1,201	2,210	2,581	3,067	3,523	3,792	4,088	4,553	1,991	27,006
Unknown	*	*	*	.	*	*	*	.	.	*
0-19	738	1,126	1,091	1,093	1,179	1,171	1,181	1,211	500	9,290
20-44	7,924	12,586	12,455	12,897	12,839	12,845	12,818	12,803	5,270	102,437
45-64	16,076	26,150	27,793	29,994	31,334	32,342	33,705	34,739	15,215	247,348
65-74	12,223	21,002	22,023	22,517	22,814	23,338	23,772	24,142	9,905	181,736
75+	8,736	15,516	17,507	19,170	21,122	22,239	22,904	25,210	10,642	163,046
Unknown	*	*	*	.	*	*	*	.	.	*
Male	23,988	40,789	43,093	45,603	47,710	49,137	50,763	53,298	22,487	376,868
Female	21,713	35,592	37,778	40,068	41,579	42,796	43,618	44,807	19,045	326,996
Unknown	*	.	.	.	*
White	28,009	47,390	50,496	53,337	56,162	57,217	60,219	62,716	26,291	441,837
Black	13,655	22,104	23,510	25,065	25,387	25,479	26,673	27,715	11,882	201,470
Native American/Alaskan Native	686	1,235	910	1,129	1,412	2,119	1,096	1,079	427	10,093
Asian	1,057	1,834	1,795	2,229	2,333	2,388	2,247	2,347	980	17,210
Pacific Islander	384	706	699	800	750	796	862	925	410	6,332
Middle-eastern/Arabian	183	189	233	282	300	377	341	384	147	2,436
Indian Subcontinent	128	187	217	239	262	293	330	293	166	2,115
Other/multiracial	1,297	2,315	2,658	2,292	2,345	2,770	2,356	2,364	1,137	19,534
Unknown	302	421	353	298	338	497	257	282	92	2,840
Hispanic-Mexican	3,371	5,493	5,446	5,773	6,435	8,065	6,868	7,469	3,288	52,208
Hispanic-other	2,008	3,297	3,637	3,903	4,119	4,244	4,651	4,916	2,203	32,978
Non-Hispanic	40,035	67,287	71,545	75,683	78,381	79,305	82,703	85,503	35,945	616,387
Unknown	287	304	243	312	354	322	159	217	96	2,294
Diabetes	19,368	33,065	35,563	37,946	39,690	41,529	42,707	43,922	18,733	312,523
Hypertension	11,775	19,963	21,479	22,645	23,896	24,351	25,316	26,708	11,623	187,756
Glomerulonephritis	5,168	8,226	8,211	8,386	8,356	8,093	7,979	8,164	3,285	65,868
Cystic kidney	1,228	1,872	1,969	2,124	2,157	2,149	2,207	2,210	955	16,871
Other urologic	888	1,581	1,505	1,555	1,656	1,613	1,593	1,682	662	12,735
Other cause	5,442	8,536	8,864	9,487	9,980	10,352	10,667	11,488	4,690	79,506
Unknown cause	1,733	3,035	3,158	3,436	3,457	3,812	3,907	3,931	1,584	28,053
Missing	99	103	122	92	97	37	*	.	.	555
All	45,701	76,381	80,871	85,671	89,289	91,936	94,381	98,105	41,532	703,867

The age of the incident patient is determined as of the date of ESRD initiation.
Data are based on first & new Medical Evidence forms filed between May 1995 & June 2003.
[^]Incomplete data
*Values for cells with 10 or fewer patients are suppressed.
"." Zero values in this cell.



Table C.5

Patient Counts: by ESRD Network

incident patients, by race & ethnicity, May 1995-April 2003 combined

	White	Black	Nat. Am/ Alas. Native	Asian	Pacific Islander	Mid-East/ Arabian	Indian Subcontinent	Other/ Multiracial	Unk.	All
1 CT, MA, ME, NH, RI, VT	22,040	3,574	59	350	31	67	57	472	534	27,184
2 NY	28,391	16,032	225	1,361	102	203	421	2,280	793	49,808
3 NJ, PR, VI	16,222	8,067	39	409	93	122	217	7,700	89	32,958
4 DE, PA	26,836	8,975	63	172	33	44	67	515	88	36,793
5 MD, DC, VA, WV	20,427	20,396	246	651	72	153	113	487	59	42,604
6 NC, SC, GA	23,388	31,156	646	505	39	43	75	537	23	56,412
7 FL	29,354	12,557	61	305	90	90	129	599	37	43,222
8 AL, MS, TN	17,687	18,179	126	123	22	31	25	81	28	36,302
9 IN, KY, OH	39,240	13,193	57	182	29	58	56	360	252	53,427
10 IL	19,553	10,750	51	484	112	103	157	404	118	31,732
11 MI, MN, ND, SD, WI	34,414	10,781	1,233	524	62	250	74	285	14	47,637
12 IA, KS, MO, NE	20,879	5,497	1,501	519	37	41	31	219	*	28,728
13 AR, LA, OK	15,573	13,062	1,222	150	31	20	28	260	*	30,356
14 TX	34,996	12,996	307	903	105	122	122	1,980	291	51,822
15 AZ, CO, NV, NM, UT, WY	22,747	2,217	2,968	470	246	65	51	756	114	29,634
16 AK, ID, MT, OR, WA	15,914	1,258	685	693	411	35	36	116	*	19,149
17 N. CA, HI, Guam, AS	17,590	4,685	354	5,119	3,536	252	229	1,197	281	33,243
18 S. CA	36,586	8,095	250	4,290	1,281	737	227	1,286	104	52,856
Unknown										
All	441,837	201,470	10,093	17,210	6,332	2,436	2,115	19,534	2,840	703,867

	Hispanic Mexican	Hispanic Other	Non- Hispanic	Unk.	All
1 CT, MA, ME, NH, RI, VT	93	1,312	24,752	1,027	27,184
2 NY	294	5,515	43,506	493	49,808
3 NJ, PR, VI	168	10,425	22,211	154	32,958
4 DE, PA	56	625	36,031	81	36,793
5 MD, DC, VA, WV	824	514	41,245	21	42,604
6 NC, SC, GA	901	330	55,181	.	56,412
7 FL	393	4,569	38,253	*	43,222
8 AL, MS, TN	67	62	36,173	.	36,302
9 IN, KY, OH	456	270	52,607	94	53,427
10 IL	1,808	713	29,167	44	31,732
11 MI, MN, ND, SD, WI	849	398	46,384	*	47,637
12 IA, KS, MO, NE	2,166	78	26,432	52	28,728
13 AR, LA, OK	258	204	29,891	*	30,356
14 TX	17,570	1,681	32,561	*	51,822
15 AZ, CO, NV, NM, UT, WY	4,997	1,815	22,813	*	29,634
16 AK, ID, MT, OR, WA	862	185	18,102	.	19,149
17 N. CA, HI, Guam, AS	5,013	1,253	26,689	288	33,243
18 S. CA	15,433	3,029	34,389	*	52,856
Unknown					
All	52,208	32,978	616,387	2,294	703,867

Data are based on first & new Medical Evidence forms filed between May 1995 & June 2003.

*Values for cells with 10 or fewer patients are suppressed.

"." Zero values in this cell.

Table C.6
Percent Distribution of Patients: by Race & Hispanic Ethnicity
incident patients

	1995 [^]	1996	1997	1998	1999	2000	2001	2002	2003 [^]
White	61.3	62.0	62.4	62.3	62.9	62.2	63.8	63.9	63.3
Black	29.9	28.9	29.1	29.3	28.4	27.7	28.3	28.3	28.6
Native American/Alaskan Native	1.5	1.6	1.1	1.3	1.6	2.3	1.2	1.1	1.0
Asian	2.3	2.4	2.2	2.6	2.6	2.6	2.4	2.4	2.4
Pacific Islander	0.8	0.9	0.9	0.9	0.8	0.9	0.9	0.9	1.0
Mid-East/Arabian	0.4	0.2	0.3	0.3	0.3	0.4	0.4	0.4	0.4
Indian Subcontinent	0.3	0.2	0.3	0.3	0.3	0.3	0.3	0.3	0.4
Other/multiracial	2.8	3.0	3.3	2.7	2.6	3.0	2.5	2.4	2.7
Unknown	0.7	0.6	0.4	0.3	0.4	0.5	0.3	0.3	0.2
Hispanic-Mexican	7.4	7.2	6.7	6.7	7.2	8.8	7.3	7.6	7.9
Hispanic-Other	4.4	4.3	4.5	4.6	4.6	4.6	4.9	5.0	5.3
Non-Hispanic	87.6	88.1	88.5	88.3	87.8	86.3	87.6	87.2	86.5
Unknown	0.6	0.4	0.3	0.4	0.4	0.4	0.2	0.2	0.2

Table C.7
Percent Distribution of Diabetic Patients: by Race & Hispanic Ethnicity
incident patients

	1995 [^]	1996	1997	1998	1999	2000	2001	2002	2003 [^]
White	41.9	42.4	43.2	43.5	43.8	44.7	44.6	44.2	44.7
Black	41.1	42.5	42.7	43.5	42.8	43.3	44.0	43.7	43.4
Native American/Alaskan Native	64.1	63.7	74.2	71.1	67.2	59.9	72.3	71.7	74.0
Asian	38.6	43.2	43.5	44.1	45.3	44.8	44.4	43.9	44.9
Pacific Islander	61.2	53.5	56.5	58.3	58.0	60.2	62.4	57.7	59.3
Mid-East/Arabian	41.0	29.1	42.1	38.7	43.0	37.1	38.7	42.4	46.9
Indian Subcontinent	37.5	44.9	46.1	47.7	45.4	46.1	51.8	54.3	44.0
Other/multiracial	52.4	56.1	55.8	55.0	57.4	57.1	56.2	54.4	54.8
Unknown	47.7	42.5	48.2	42.3	50.6	53.5	47.5	46.8	53.3
Hispanic-Mexican	60.8	61.7	65.1	63.2	63.2	61.4	63.9	63.1	63.6
Hispanic-Other	50.0	52.0	53.3	54.2	55.9	55.6	55.6	54.4	56.1
Non-Hispanic	40.5	41.4	41.9	42.3	42.3	43.0	43.1	42.6	42.7
Unknown	40.8	44.7	42.8	43.9	44.4	36.6	44.7	35.5	46.9
All	42.4	43.3	44.0	44.3	44.5	45.2	45.2	44.8	45.1

Table C.8
Patient Counts: by Employment Status
incident patients

	1995 [^]	1996	1997	1998	1999	2000	2001	2002	2003 [^]
Current employment status									
Employed full-time	3,705	6,094	6,765	7,356	7,781	7,889	8,241	8,232	3,490
Employed part-time	871	1,361	1,423	1,578	1,557	1,556	1,629	1,670	680
Homemaker	2,778	4,464	4,624	4,811	4,816	4,914	4,766	4,787	1,951
Retired due to age/preference	14,482	25,797	29,290	32,555	34,552	35,448	36,576	38,952	16,209
Retired due to disability	7,590	12,898	14,011	15,798	16,215	16,854	17,239	18,157	7,749
Medical leave of absence	1,396	2,027	2,046	2,281	2,505	2,956	3,038	3,175	1,389
Student	576	892	876	914	965	929	960	1,026	423
Unemployed	9,955	15,538	15,831	16,436	16,997	17,443	18,033	17,805	7,984
Missing	4,348	7,310	6,005	3,942	3,901	3,947	3,899	4,301	1,657
Previous employment status									
Employed full-time	6,804	10,549	11,280	12,405	12,929	13,295	13,744	13,070	5,370
Employed part-time	1,205	1,818	1,827	2,110	2,031	2,015	2,051	2,086	868
Homemaker	2,791	4,479	4,601	4,817	4,805	4,934	4,792	4,641	1,851
Retired due to age/preference	13,331	24,068	27,513	30,861	32,561	33,519	34,720	35,998	14,500
Retired due to disability	6,172	10,894	12,027	13,635	14,119	14,765	15,014	15,370	6,460
Medical leave of absence	254	408	391	442	459	509	478	501	204
Student	560	875	887	895	938	929	942	976	418
Unemployed	7,692	12,241	12,567	12,927	13,785	13,927	14,416	13,670	6,034
Missing	6,892	11,049	9,778	7,579	7,662	8,043	8,224	11,793	5,827
All	45,701	76,381	80,871	85,671	89,289	91,936	94,381	98,105	41,532



Table C.9
Patient Counts: by Insurance Payor
incident patients

	1995 [^]	1996	1997	1998	1999	2000	2001	2002	2003 [^]
Medicaid coverage	11,423	18,212	19,074	20,159	20,298	21,919	22,269	23,965	10,281
Medicare coverage	21,969	38,386	41,230	42,804	44,963	50,233	51,966	54,964	22,934
DVA coverage	818	1,014	912	952	1,399	1,078	1,183	1,262	546
Employer group health insurance	11,452	18,248	19,352	19,899	20,478	21,039	22,152	22,892	9,404
Other medical insurance	13,261	24,193	26,090	27,518	28,515	29,519	30,860	31,478	13,171
No medical insurance	3,468	5,519	5,762	5,950	7,075	7,331	7,257	7,322	3,280
Patient applying for Medicare	38,324	64,368	67,651	70,616	72,597	72,767	73,905	75,946	32,121
All	45,701	76,381	80,871	85,671	89,289	91,936	94,381	98,105	41,532

Table C.10
Percent Distribution of Patients: by Comorbidity
incident patients

	1995 [^]	1996	1997	1998	1999	2000	2001	2002	2003 [^]
Congestive heart failure	30.8	32.1	33.0	32.4	31.6	32.1	31.5	31.1	31.9
Ischemic heart disease/CAD	21.3	22.5	23.6	23.8	23.9	24.6	24.9	25.0	24.9
Myocardial infarction	8.4	8.8	8.9	8.8	8.8	9.0	8.9	8.7	8.3
Cardiac arrest	1.0	1.0	0.9	0.8	0.8	0.8	0.7	0.7	0.7
Cardiac dysrhythmia	5.3	5.5	5.8	5.7	5.8	6.0	6.0	6.2	6.2
Pericarditis	1.5	1.2	1.0	0.9	0.8	0.7	0.6	0.5	0.5
Cerebrovascular disease, CVA/TIA	8.0	8.7	9.1	9.0	9.1	9.1	9.4	9.4	9.0
Peripheral vascular disease	13.9	14.2	14.8	14.6	14.2	14.4	14.2	13.9	13.9
History of hypertension	67.3	70.3	72.4	73.9	75.2	76.6	78.3	79.3	80.4
Diabetes: primary/contributing	36.0	38.0	39.3	40.4	41.5	43.3	44.7	45.4	46.3
Diabetes: on insulin	22.1	23.1	23.5	23.4	23.2	23.1	22.6	22.3	22.7
Chronic obstructive pulmonary disease	6.6	6.8	7.0	7.0	7.1	7.3	7.5	7.6	7.8
Current smoker/tobacco use	5.9	5.7	5.4	5.2	5.0	4.9	5.0	5.0	4.9
Malignant neoplasm/cancer	4.8	4.7	4.9	5.1	5.4	5.6	5.9	6.2	5.8
Alcohol dependence	1.8	1.8	1.6	1.4	1.4	1.4	1.3	1.3	1.2
Drug dependence	1.1	1.1	1.0	1.0	1.0	1.0	1.0	0.9	0.9
AIDS	4.3	4.2	2.5	0.5	1.7	1.8	0.4	0.4	0.4
HIV	4.6	4.6	2.8	0.9	2.0	2.2	0.7	0.8	0.7
Inability to ambulate	4.9	4.5	4.2	3.9	3.9	4.1	3.9	4.2	4.0
Inability to transfer	1.7	1.6	1.6	1.5	1.4	1.5	1.5	1.5	1.4
All	45,701	76,381	80,871	85,671	89,289	91,936	94,381	98,105	41,532

Table C.11
Percent Distribution of Patients: by Primary Diagnosis
incident patients

	1995 [^]	1996	1997	1998	1999	2000	2001	2002	2003 [^]
Diabetes	42.4	43.3	44.0	44.3	44.5	45.2	45.2	44.8	45.1
Hypertension	25.8	26.1	26.6	26.4	26.8	26.5	26.8	27.2	28.0
Glomerulonephritis	11.3	10.8	10.2	9.8	9.4	8.8	8.5	8.3	7.9
Cystic kidney	2.7	2.5	2.4	2.5	2.4	2.3	2.3	2.3	2.3
Other urologic	1.9	2.1	1.9	1.8	1.9	1.8	1.7	1.7	1.6
Other cause	11.9	11.2	11.0	11.1	11.2	11.3	11.3	11.7	11.3
Unknown cause	4.0	4.1	4.1	4.1	4.0	4.2	4.1	4.0	3.8
All patients	45,701	76,381	80,871	85,671	89,289	91,936	94,381	98,105	41,532

The age of the incident patient is determined as of the date of ESRD initiation.
 Data are based on first & new Medical Evidence forms filed between May 1995 & June 2003.
[^]Incomplete data
 On our website: Patient Comorbidities in detail (C.c-m).



Table C.12
Lab Values & Physical Characteristics: Pediatric & Adult Patients
incident patients

	1995 [^] # of forms w/value	Mean values	1996 # of forms w/value	Mean values	1997 # of forms w/value	Mean values	1998 # of forms w/value	Mean values	1999 # of forms w/value	Mean values	2000 # of forms w/value	Mean values	2001 # of forms w/value	Mean values	2002 # of forms w/value	Mean values	2003 [^] # of forms w/value	Mean values
~0-19																		
Serum albumin (g/dl)	644	3.5	994	3.4	937	3.3	931	3.3	956	3.3	960	3.2	957	3.2	1,009	3.2	418	3.3
BUN (mg/dl)	676	94.0	1,050	92.0	1,005	90.3	1,025	91.2	1,096	85.3	1,082	87.1	1,106	84.0	1,128	86.4	472	89.5
Serum creatinine (mg/dl)	717	9.8	1,099	9.6	1,066	9.4	1,062	9.2	1,163	8.7	1,153	8.8	1,163	8.6	1,190	8.7	485	9.0
Hematocrit (%)	703	27.0	1,078	26.6	1,025	27.3	1,024	27.3	1,122	27.9	1,101	28.3	1,120	28.5	1,150	28.6	482	28.8
Hemoglobin (g/dl)	644	9.2	1,019	9.1	983	9.2	972	9.3	1,070	9.4	1,061	9.6	1,103	9.7	1,118	9.7	467	9.7
Height (cm)	701	146.0	1,061	146.1	1,017	145.8	1,021	146.3	1,110	146.9	1,114	144.1	1,125	145.6	1,154	145.2	477	149.1
Weight (kg)	737	44.8	1,122	45.2	1,088	44.6	1,090	46.6	1,177	47.8	1,170	48.2	1,181	49.1	1,209	48.2	499	51.0
Body mass index (kg/m ²)	635	20.8	951	21.0	915	21.5	927	21.7	1,044	21.9	1,078	22.0	1,092	22.1	1,126	22.1	467	22.3
eGFR (ml/min) by Schwartz form.	674	9.6	1,033	9.8	989	9.9	978	9.9	1,081	10.4	1,083	10.5	1,086	10.7	1,117	10.8	453	10.8
~20+																		
Serum albumin (g/dl)	37,065	3.2	61,570	3.2	64,020	3.2	64,172	3.2	64,513	3.1	66,703	3.1	68,071	3.1	71,106	3.1	30,191	3.1
BUN (mg/dl)	41,259	94.0	70,219	94.2	74,662	92.4	79,150	90.3	82,493	88.8	85,231	87.8	87,453	86.6	91,598	85.6	38,848	85.3
Serum creatinine (mg/dl)	44,756	8.6	74,919	8.4	79,420	8.1	84,195	7.9	87,689	7.6	90,282	7.4	92,723	7.3	96,445	7.1	40,841	7.0
Hematocrit (%)	41,908	27.9	70,672	28.1	74,952	28.5	79,545	28.8	82,478	29.2	84,567	29.5	86,650	29.7	90,656	30.0	38,466	30.1
Hemoglobin (g/dl)	38,037	9.4	64,724	9.4	69,389	9.5	74,118	9.6	77,560	9.7	80,797	9.8	83,679	9.9	88,274	10.0	37,589	10.0
Height (cm)	43,938	167.1	73,487	167.3	77,913	167.3	82,509	167.2	86,677	167.0	90,427	167.0	92,969	167.4	96,737	167.6	40,988	167.7
Weight (kg)	44,894	66.7	75,122	67.9	79,625	68.5	84,390	69.6	87,841	72.3	90,389	75.9	92,907	76.8	96,640	77.3	40,917	77.9
Body mass index (kg/m ²)	39,300	25.3	66,516	25.5	70,763	25.7	75,698	26.1	82,153	26.6	88,955	27.1	91,982	27.3	96,059	27.5	40,761	27.7
eGFR (ml/min) by MDRD form.	44,615	7.5	74,705	7.7	79,130	8.1	83,807	8.5	87,201	8.8	89,636	9.1	91,988	9.4	95,593	9.7	40,455	9.8

The age of the incident patient is determined as of the date of ESRD initiation.
 Data are based on first & new Medical Evidence forms filed between May 1995 & June 2003.
[^]Incomplete data.

Table C.13

Lab Values & Physical Characteristics

incident patients, by demographic characteristics, May 1995-April 2003 combined

	Patients	Serum albumin (g/dl)	BUN (mg/dl)	Creat. (mg/dl)	Hct %	Hgb (g/dl)	Height (cm)	Weight (kg)	Body mass index (kg/m ²)	eGFR (ml/min 1.73m ²) by MDRD formula
0-4	1,479	3.1	73.7	5.0	29.2	9.9	99.3	20.7	19.2	10.6
5-9	1,121	3.3	88.3	7.0	27.9	9.5	117.5	25.4	18.1	10.8
10-14	2,266	3.4	89.6	9.1	27.9	9.5	146.0	43.5	20.5	11.2
15-19	4,424	3.3	92.5	10.9	27.2	9.2	164.5	63.4	23.8	9.6
20-24	7,905	3.2	93.7	11.6	26.7	9.1	167.7	69.3	25.3	7.1
25-29	13,049	3.2	90.3	10.9	27.3	9.2	168.0	71.7	26.0	7.2
30-34	19,188	3.2	88.9	10.4	27.8	9.4	168.7	73.2	26.4	7.4
35-39	26,465	3.1	88.9	10.0	28.1	9.4	169.4	74.9	26.8	7.5
40-44	35,830	3.1	89.3	9.6	28.3	9.5	169.6	76.9	27.4	7.7
45-49	46,984	3.1	89.3	9.0	28.6	9.6	169.3	78.3	27.9	7.9
50-54	57,834	3.1	89.3	8.4	28.9	9.6	168.8	79.3	28.4	8.2
55-59	66,112	3.1	89.1	7.9	29.0	9.7	168.0	77.9	28.2	8.5
60-64	76,418	3.1	89.1	7.5	29.1	9.7	167.3	75.8	27.7	8.7
65-69	88,288	3.2	89.1	7.1	29.4	9.8	166.8	73.0	26.9	9.0
70-74	93,448	3.2	89.0	6.8	29.6	9.8	166.4	70.3	26.0	9.3
75-79	83,125	3.2	89.3	6.5	29.8	9.9	166.0	67.6	25.1	9.6
80-84	52,915	3.2	89.3	6.3	29.9	10.0	165.3	64.8	24.3	9.9
85+	27,006	3.1	89.1	6.0	30.0	10.0	163.8	61.7	23.4	10.1
Unknown	*	3.3	75.1	6.8	28.7	9.6	167.8	64.8	24.6	11.0
0-19	9,290	3.3	88.5	9.0	27.8	9.4	145.9	47.2	21.7	10.3
20-44	102,437	3.1	89.6	10.1	27.9	9.4	169.0	74.4	26.7	7.5
45-64	247,348	3.1	89.2	8.1	28.9	9.6	168.2	77.7	28.0	8.4
65-74	181,736	3.2	89.0	7.0	29.5	9.8	166.6	71.6	26.4	9.1
75+	163,046	3.2	89.2	6.4	29.9	9.9	165.4	65.7	24.6	9.8
Unknown	*	3.3	75.1	6.8	28.7	9.6	167.8	64.8	24.6	11.0
Male	376,868	3.2	93.2	8.3	29.3	9.8	173.1	76.3	26.0	9.1
Female	326,996	3.1	84.6	7.0	28.9	9.6	160.0	68.1	27.2	8.4
Unknown	*	2.5	59.7	4.5	29.3	9.6	180.2	79.2	24.6	15.4
White	441,837	3.2	89.9	7.1	29.6	9.9	167.3	72.3	26.4	9.1
Black	201,470	3.1	87.5	8.9	28.1	9.3	167.8	75.0	27.2	8.5
Native American/Alaskan Native	10,093	2.9	85.5	7.7	28.5	9.5	165.8	73.3	27.3	8.2
Asian	17,210	3.2	91.1	8.2	28.9	9.7	159.9	58.5	23.4	7.8
Pacific Islander	6,332	3.0	94.5	8.7	28.9	9.6	161.6	65.6	25.5	7.2
Middle-eastern/Arabian	2,436	3.3	91.5	7.9	29.6	9.8	163.4	68.1	25.7	8.5
Indian Subcontinent	2,115	3.2	85.7	7.9	29.1	9.7	163.2	62.4	23.9	8.5
Other/multiracial	19,534	3.1	90.0	8.4	28.4	9.5	162.0	67.4	26.0	7.8
Unknown	2,840	3.1	92.7	8.1	28.4	9.5	163.7	69.3	26.0	8.2
Hispanic-Mexican	52,208	3.1	87.1	7.7	28.6	9.6	162.2	66.8	26.3	8.5
Hispanic-Other	32,978	3.1	89.3	8.2	28.7	9.6	162.8	68.3	26.0	8.0
Non-Hispanic	616,387	3.2	89.4	7.7	29.2	9.7	167.6	73.2	26.6	8.9
Unknown	2,294	3.2	96.5	7.9	29.1	9.8	166.0	70.5	25.9	8.4
Diabetic	312,523	3.1	87.4	6.9	29.2	9.7	166.6	75.9	27.9	9.5
Non-diabetic	391,344	3.2	90.6	8.4	29.0	9.7	167.3	69.7	25.5	8.2
Year of first service										
1995 [^]	45,701	3.2	94.0	8.7	27.9	9.4	166.8	66.3	25.2	7.5
1996	76,381	3.2	94.1	8.5	28.1	9.4	167.0	67.5	25.4	7.8
1997	80,871	3.2	92.4	8.1	28.5	9.5	167.0	68.2	25.7	8.1
1998	85,671	3.2	90.3	7.9	28.8	9.6	166.9	69.3	26.0	8.5
1999	89,289	3.1	88.7	7.6	29.2	9.7	166.7	72.0	26.5	8.8
2000	91,936	3.1	87.8	7.4	29.5	9.8	166.7	75.6	27.0	9.1
2001	94,381	3.1	86.6	7.3	29.7	9.9	167.2	76.5	27.2	9.4
2002	98,105	3.1	85.6	7.1	30.0	10.0	167.3	77.0	27.5	9.7
2003 [^]	41,532	3.1	85.4	7.1	30.1	10.0	167.4	77.5	27.6	9.8

Data are based on first & new Medical Evidence forms filed between May 1995 & June 2003.

The age of the incident patient is determined as of the date of ESRD initiation.

[^]Incomplete data.

*Values for cells with 10 or fewer patients are suppressed.

Table C.14
Lab Values & Physical Characteristics: by Year & Age
incident patients, May 1995-April 2003 combined

		Patients	Serum						Body mass GFR (ml/min 1.73m ²)		
			albumin (g/dl)	BUN (mg/dl)	Creat. (mg/dl)	Hct %	Hgb (g/dl)	Height (cm)	Weight (kg)	index (kg/m ²)	by MDRD formula
1995 [^]	0-19	738	3.5	94.0	9.8	27.0	9.2	146.0	44.8	20.8	9.6
	20-44	7,924	3.2	94.4	11.0	26.9	9.1	169.1	68.0	25.2	6.7
	45-64	16,076	3.2	93.8	8.9	27.7	9.3	168.0	70.7	26.5	7.3
	65-74	12,223	3.3	93.2	7.8	28.3	9.5	166.3	65.4	25.1	7.8
	75+	8,736	3.3	95.0	7.3	28.6	9.6	164.9	60.0	23.5	8.2
	Unknown	*	3.8	66.8	8.0	30.3	10.1	171.0	59.2	24.1	8.1
1996	0-19	1,126	3.4	92.0	9.6	26.6	9.1	146.1	45.2	21.0	9.8
	20-44	12,586	3.2	94.1	10.8	27.0	9.1	169.1	69.5	25.6	6.8
	45-64	26,150	3.2	94.0	8.8	27.9	9.4	168.1	72.0	26.8	7.5
	65-74	21,002	3.2	94.1	7.6	28.5	9.6	166.7	66.7	25.3	8.1
	75+	15,516	3.2	94.5	7.1	28.8	9.7	165.3	61.1	23.6	8.4
	Unknown	*	2.7	.	8.5	.	9.6	.	71.0	.	.
1997	0-19	1,091	3.3	90.3	9.4	27.3	9.2	145.8	44.6	21.5	9.9
	20-44	12,455	3.2	92.6	10.5	27.4	9.2	169.1	70.1	25.9	7.1
	45-64	27,793	3.2	92.4	8.5	28.2	9.4	168.2	72.9	27.0	7.8
	65-74	22,023	3.2	91.7	7.3	28.9	9.6	166.7	67.3	25.4	8.5
	75+	17,507	3.2	93.3	6.8	29.1	9.7	165.3	62.0	23.9	8.9
	Unknown	*	2.6	78.5	5.5	23.8	8.0	169.0	70.5	24.6	11.1
1998	0-19	1,093	3.3	91.2	9.2	27.3	9.3	146.3	46.6	21.7	9.9
	20-44	12,897	3.2	91.3	10.3	27.6	9.3	169.1	71.3	26.2	7.3
	45-64	29,994	3.2	89.9	8.2	28.6	9.5	168.1	74.1	27.3	8.1
	65-74	22,517	3.2	90.1	7.1	29.2	9.7	166.5	68.6	25.9	8.8
	75+	19,170	3.2	90.6	6.6	29.5	9.8	165.3	62.8	24.2	9.3
	Unknown										
1999	0-19	1,179	3.3	85.3	8.7	27.9	9.4	146.9	47.8	21.9	10.4
	20-44	12,839	3.1	89.3	10.1	28.0	9.4	168.7	74.3	26.8	7.5
	45-64	31,334	3.1	88.8	8.1	29.0	9.6	167.9	77.1	28.0	8.4
	65-74	22,814	3.2	88.0	6.9	29.6	9.8	166.4	71.3	26.4	9.2
	75+	21,122	3.2	89.2	6.4	29.9	9.9	165.2	65.1	24.4	9.7
	Unknown	*	3.5	72.0	9.6	29.4	9.9	168.0	71.0	25.2	5.1
2000	0-19	1,171	3.2	87.1	8.8	28.3	9.6	144.1	48.2	22.0	10.5
	20-44	12,845	3.1	87.4	9.8	28.4	9.5	168.6	78.0	27.3	7.8
	45-64	32,342	3.1	88.0	7.8	29.2	9.7	167.9	81.1	28.6	8.7
	65-74	23,338	3.1	87.7	6.7	29.8	9.9	166.4	75.1	27.1	9.6
	75+	22,239	3.1	87.8	6.1	30.2	10.0	165.3	68.0	24.8	10.1
	Unknown	*	3.4	60.0	3.2	26.9	9.2	160.0	74.0	28.9	22.5
2001	0-19	1,181	3.2	84.0	8.6	28.5	9.7	145.6	49.1	22.1	10.7
	20-45	12,818	3.1	86.2	9.7	28.6	9.6	169.1	79.2	27.5	7.9
	45-65	33,705	3.1	86.8	7.7	29.5	9.8	168.5	82.0	28.8	8.9
	65-75	23,772	3.1	86.7	6.6	30.0	10.0	166.8	75.8	27.2	9.8
	75+	22,904	3.1	86.4	6.0	30.4	10.1	165.6	68.9	25.1	10.4
	Unknown	*	.	120.0	4.1	33.0	10.9	160.0	54.0	21.1	11.0
2002	0-19	1,211	3.2	86.4	8.7	28.6	9.7	145.2	48.2	22.1	10.8
	20-45	12,803	3.1	85.2	9.5	28.7	9.7	169.4	79.8	27.7	8.1
	45-65	34,739	3.1	85.8	7.6	29.7	9.9	168.6	82.8	29.1	9.2
	65-75	24,142	3.1	85.3	6.4	30.3	10.1	166.9	76.6	27.5	10.1
	75+	25,210	3.1	85.9	5.9	30.7	10.2	165.8	69.4	25.3	10.7
	Unknown										
2003 [^]	0-19	500	3.3	89.5	9.0	28.8	9.7	149.1	51.0	22.3	10.8
	20-45	5,270	3.1	85.7	9.5	28.8	9.6	169.3	80.3	27.9	8.1
	45-65	15,215	3.1	85.3	7.5	29.9	9.9	168.7	83.1	29.2	9.3
	65-75	9,905	3.1	85.5	6.4	30.5	10.1	167.0	77.2	27.7	10.2
	75+	10,642	3.1	85.1	5.8	30.8	10.2	165.9	69.9	25.4	10.9
	Unknown										

[^]Incomplete data. Data are based on first & new Medical Evidence forms filed between May 1995 & June 2003. The age of the incident patient is determined as of the date of ESRD initiation. *Values for cells with 10 or fewer patients are suppressed. "." Zero values in this cell.



Table C.15

Lab Values & Physical Characteristics: by Race

incident patients, by gender, age, & diabetic status, May 1995-April 2003 combined

WHITE			Serum					Body mass			~eGFR (ml/min) by MDRD formula		
Age group	Diabetic status	Patients	albumin (g/dl)	BUN (mg/dl)	Creat. (mg/dl)	Hct %	Hgb (g/dl)	Height (cm)	Weight (kg)	index (kg/m ²)			
Male	0-19	Diabetic	47	3.2	88.8	8.0	28.0	9.6	169.3	67.7	24.4	11.1	
		Non-diabetic	3,232	3.5	90.0	8.5	28.7	9.8	144.9	44.7	20.9	11.1	
	20-44	Diabetic	11,901	3.0	84.2	8.0	29.1	9.9	173.9	75.6	25.8	9.4	
		Non-diabetic	18,357	3.5	98.4	10.9	28.5	9.7	174.5	78.4	26.4	7.1	
	45-64	Diabetic	41,534	3.1	92.0	7.3	29.5	9.9	174.1	84.3	28.4	9.7	
		Non-diabetic	36,365	3.3	94.4	8.8	29.6	10.0	175.0	80.1	26.8	8.0	
	65-74	Diabetic	29,413	3.2	94.1	6.5	30.0	10.0	173.3	78.9	26.9	10.6	
		Non-diabetic	38,128	3.3	93.1	7.6	29.9	10.0	173.9	73.8	25.1	9.0	
	75+	Diabetic	18,410	3.2	94.5	5.9	30.2	10.1	172.2	74.0	25.5	11.4	
		Non-diabetic	49,946	3.2	93.2	6.8	30.2	10.1	172.3	69.4	24.0	9.9	
	Female	0-19	Diabetic	54	3.0	77.8	6.5	27.5	9.1	159.3	62.2	25.3	11.1
			Non-diabetic	2,425	3.3	85.9	8.2	27.8	9.4	140.3	42.3	21.1	10.1
		20-44	Diabetic	8,270	3.0	78.3	6.8	28.8	9.6	161.4	66.0	26.2	8.7
			Non-diabetic	12,267	3.4	84.8	8.8	28.3	9.5	161.2	66.1	26.3	6.9
45-64		Diabetic	35,342	3.0	85.3	6.1	29.3	9.7	160.6	76.8	30.4	8.9	
		Non-diabetic	25,653	3.3	85.7	7.5	29.4	9.8	161.3	68.6	27.1	7.3	
65-74		Diabetic	29,829	3.1	87.2	5.7	29.7	9.9	159.6	71.8	28.7	9.3	
		Non-diabetic	25,931	3.2	85.2	6.6	29.7	9.9	160.0	63.3	25.5	8.1	
75+		Diabetic	18,588	3.1	87.0	5.2	30.0	10.0	158.4	66.1	26.8	10.0	
		Non-diabetic	36,138	3.2	85.8	6.0	30.0	10.0	158.4	58.7	24.0	8.8	
BLACK													
Male		0-19	Diabetic	30	3.1	78.3	7.6	28.1	9.4	166.9	71.2	26.3	12.5
			Non-diabetic	1,261	3.2	89.0	10.9	27.7	9.3	155.6	55.4	22.7	10.0
		20-44	Diabetic	4,479	2.9	83.7	9.6	28.0	9.3	175.1	84.9	28.3	9.4
	Non-diabetic		19,262	3.1	98.8	13.1	27.5	9.3	175.6	80.2	26.7	6.9	
	45-64	Diabetic	19,191	3.1	89.0	8.7	28.4	9.4	175.5	84.8	28.0	9.6	
		Non-diabetic	23,663	3.2	95.7	10.8	28.1	9.4	175.2	78.5	26.1	7.9	
	65-74	Diabetic	8,767	3.2	90.3	7.8	29.0	9.6	173.5	79.0	26.8	10.4	
		Non-diabetic	9,362	3.2	91.9	8.9	28.4	9.4	173.6	73.1	24.9	9.2	
	75+	Diabetic	3,938	3.2	88.7	7.0	29.2	9.7	171.8	73.4	25.5	11.3	
		Non-diabetic	7,443	3.2	89.7	7.8	29.0	9.6	172.2	67.4	23.5	10.3	
	Female	0-19	Diabetic	28	2.8	66.8	8.3	26.9	8.8	161.0	60.7	26.4	9.7
			Non-diabetic	1,103	3.0	85.5	10.1	26.1	8.7	151.8	56.1	24.1	9.3
		20-44	Diabetic	4,367	2.8	73.4	8.2	27.2	9.0	163.2	79.0	30.2	8.5
			Non-diabetic	13,596	3.0	84.4	10.7	26.4	8.8	162.9	71.9	27.8	6.6
45-64		Diabetic	23,358	3.0	80.7	7.5	28.2	9.3	162.4	79.2	30.5	8.5	
		Non-diabetic	17,590	3.2	85.7	9.1	27.7	9.2	162.3	72.2	28.0	7.2	
65-74		Diabetic	15,011	3.1	81.5	6.9	28.6	9.4	161.4	72.6	28.4	8.9	
		Non-diabetic	10,726	3.2	83.5	7.7	28.3	9.3	161.0	66.2	26.2	8.2	
75+		Diabetic	7,681	3.1	81.6	6.4	28.8	9.5	160.3	67.1	26.7	9.6	
		Non-diabetic	10,608	3.1	83.0	6.9	28.6	9.4	159.7	60.3	24.3	8.9	
NATIVE AMERICAN/ALASKAN NATIVE													
Male		0-19	Diabetic
			Non-diabetic	61	3.3	81.6	9.8	27.2	9.2	149.0	49.2	22.2	11.0
		20-44	Diabetic	403	2.7	82.5	8.7	28.1	9.4	174.0	84.4	28.5	8.4
	Non-diabetic		430	3.3	99.0	12.1	27.4	9.3	174.4	80.3	27.1	6.6	
	45-64	Diabetic	1,720	2.9	87.7	8.3	28.6	9.6	173.6	83.3	28.0	8.5	
		Non-diabetic	591	3.1	94.6	9.8	27.8	9.3	174.4	78.1	26.5	7.4	
	65-74	Diabetic	614	3.0	88.4	7.0	29.5	9.9	171.7	75.9	26.5	9.8	
		Non-diabetic	342	3.1	94.5	8.2	29.0	9.7	173.5	70.6	24.5	8.5	
	75+	Diabetic	232	3.2	93.0	6.2	29.4	9.7	170.8	69.2	24.9	10.8	
		Non-diabetic	320	3.2	90.6	7.1	29.5	9.8	171.5	64.6	23.9	9.8	
	Female	0-19	Diabetic	*	2.8	84.0	5.3	23.4	8.4	170.2	80.0	27.6	8.5
			Non-diabetic	53	3.0	78.9	8.8	26.3	9.0	141.7	48.2	21.6	9.4
		20-44	Diabetic	362	2.6	74.9	7.6	27.4	9.0	162.1	77.3	30.0	7.9
			Non-diabetic	353	3.2	85.1	9.9	26.7	8.9	160.5	68.4	27.1	6.2
45-64		Diabetic	2,057	2.8	80.1	6.9	28.6	9.5	160.0	73.9	29.3	7.8	
		Non-diabetic	499	3.2	85.8	8.1	27.9	9.4	160.6	68.1	27.1	6.8	
65-74		Diabetic	1,019	2.9	80.8	6.2	28.8	9.5	158.8	67.0	27.0	8.5	
		Non-diabetic	327	3.1	82.7	7.0	28.3	9.4	159.6	62.9	25.8	7.6	
75+		Diabetic	397	2.9	83.8	5.8	29.3	9.7	157.9	62.7	25.9	8.8	
		Non-diabetic	312	3.1	84.1	6.1	28.7	9.5	157.6	54.0	23.4	8.6	

Data are based on first & new Medical Evidence forms filed between May 1995 & June 2003.

The age of the incident patient is determined as of the date of ESRD initiation.

~Schwartz formula used for ages 0-18; Levey formula for ages 19+.

*Values for cells with 10 or fewer patients are suppressed.

"." Zero values in this cell.

Table C.15 (continued)

Lab Values & Physical Characteristics: by Race

incident patients, by gender, age, & diabetic status, May 1995-April 2003 combined

ASIAN			Patients	Serum	BUN	Creat.	Hct	Hgb	Height	Weight	Body mass index	~eGFR (ml/min)	
Age group	Diabetic status	(g/dl)		(mg/dl)	(mg/dl)	%	(g/dl)	(cm)	(kg)	(kg/m ²)	by MDRD formula		
Male	0-19	Diabetic	*	3.4	71.0	20.2	21.6	7.3	172.4	60.0	20.2	3.6	
		Non-diabetic	119	3.2	99.7	10.7	28.3	9.5	145.1	44.1	20.1	9.8	
	20-44	Diabetic	225	3.0	90.6	9.3	29.2	9.7	168.8	74.9	26.6	7.8	
		Non-diabetic	1,101	3.4	106.9	12.6	26.7	9.1	166.8	63.5	23.3	6.0	
	45-64	Diabetic	1,663	3.0	92.8	8.4	29.1	9.8	167.1	68.9	25.2	8.2	
		Non-diabetic	1,509	3.4	98.1	10.2	28.8	9.7	166.0	64.8	24.1	6.8	
	65-74	Diabetic	1,080	3.1	91.7	7.5	29.8	10.0	165.3	63.5	23.7	9.1	
		Non-diabetic	1,055	3.3	92.2	8.3	29.4	9.9	164.5	60.0	22.6	8.1	
	75+	Diabetic	813	3.1	91.7	6.6	30.1	10.1	163.3	59.9	23.0	10.1	
		Non-diabetic	1,285	3.2	91.7	7.3	29.7	10.0	163.4	56.9	21.9	9.1	
	Female	~0-19	Diabetic
			Non-diabetic	111	3.1	97.9	8.7	25.8	8.7	145.6	41.7	19.9	9.4
		20-44	Diabetic	174	2.8	83.9	8.1	28.2	9.4	157.0	60.8	25.8	7.2
			Non-diabetic	1,038	3.3	91.6	9.9	27.0	9.0	156.2	52.5	22.3	5.9
45-64		Diabetic	1,304	3.0	85.9	7.3	28.9	9.6	155.9	60.4	25.3	7.4	
		Non-diabetic	1,349	3.4	91.3	8.7	28.3	9.4	155.2	53.9	23.1	6.1	
65-74		Diabetic	1,320	3.1	83.6	6.4	29.3	9.8	154.3	54.8	23.6	8.1	
		Non-diabetic	971	3.3	84.7	7.3	29.3	9.8	153.5	52.1	22.7	7.2	
75+		Diabetic	980	3.0	86.0	5.9	29.6	9.9	153.0	52.0	22.6	8.9	
		Non-diabetic	1,111	3.2	85.3	6.5	29.8	9.9	151.7	48.0	21.3	7.9	
PACIFIC ISLANDER													
Male		0-19	Diabetic
			Non-diabetic	28	3.0	117.1	11.5	26.8	8.9	148.4	46.8	22.3	9.7
		~20-44	Diabetic	154	2.9	100.2	11.1	28.6	9.5	170.6	89.1	31.0	6.7
	Non-diabetic		311	3.3	109.4	13.3	28.0	9.5	169.6	73.9	26.1	5.6	
	45-64	Diabetic	944	2.9	100.9	9.5	28.9	9.7	169.2	78.0	27.7	7.2	
		Non-diabetic	429	3.3	106.1	11.0	28.9	9.7	167.9	70.7	25.7	6.3	
	65-74	Diabetic	457	3.0	93.5	7.7	29.9	10.0	167.0	67.4	24.9	8.9	
		Non-diabetic	296	3.2	95.7	8.8	28.9	9.7	164.9	61.4	23.2	7.6	
	75+	Diabetic	194	3.1	94.8	6.7	30.1	10.0	164.5	60.9	23.2	10.3	
		Non-diabetic	281	3.2	92.6	7.6	29.6	9.9	162.7	56.0	21.8	9.1	
	Female	0-19	Diabetic	*	3.1	78.0	4.4	40.4	13.5	152.0	39.0	16.9	10.4
			Non-diabetic	46	2.9	95.7	9.4	26.2	8.7	146.6	44.6	21.4	9.2
		20-44	Diabetic	150	2.6	85.5	9.0	27.1	9.0	160.2	74.7	30.1	6.4
			Non-diabetic	316	3.3	97.0	11.1	27.0	9.0	158.4	62.0	24.9	5.4
45-64		Diabetic	920	2.8	89.9	8.0	28.6	9.5	157.6	67.9	27.7	6.7	
		Non-diabetic	395	3.3	92.4	9.2	28.1	9.4	156.4	59.0	24.7	5.7	
65-74		Diabetic	603	3.0	87.2	6.8	29.3	9.7	156.1	59.5	24.7	7.6	
		Non-diabetic	266	3.3	88.8	7.9	29.4	9.8	154.2	54.3	23.2	6.3	
75+		Diabetic	280	3.0	85.4	5.8	29.8	10.0	153.9	54.2	23.1	8.7	
		Non-diabetic	261	3.2	84.9	6.5	29.8	9.9	151.3	48.6	21.7	7.9	
MID-EAST/ARABIAN													
Male		0-19	Diabetic
			Non-diabetic	53	3.5	101.7	9.5	27.9	9.4	129.9	37.9	20.1	9.3
		20-44	Diabetic	34	3.2	97.5	8.5	28.1	9.3	168.9	80.5	27.2	8.5
	Non-diabetic		230	3.6	102.0	11.7	29.0	9.8	169.2	71.1	25.3	6.8	
	45-64	Diabetic	304	3.2	90.4	7.4	29.8	9.9	168.4	74.9	26.8	9.4	
		Non-diabetic	293	3.5	100.0	9.6	29.5	9.8	170.5	75.3	26.1	7.1	
	65-74	Diabetic	196	3.2	90.3	6.6	29.8	10.0	169.1	73.7	26.1	10.4	
		Non-diabetic	189	3.3	84.1	7.8	30.0	9.9	167.9	68.3	24.5	9.1	
	75+	Diabetic	95	3.2	98.3	6.4	30.9	10.2	166.1	69.2	25.0	11.3	
		Non-diabetic	178	3.3	91.8	7.0	30.3	10.0	167.9	65.4	23.3	9.7	
	Female	0-19	Diabetic
			Non-diabetic	33	3.5	85.3	8.8	27.1	9.1	134.0	38.4	19.9	8.7
		20-44	Diabetic	15	3.2	92.2	8.6	28.5	9.5	157.3	71.3	28.9	7.0
			Non-diabetic	113	3.6	89.3	10.1	27.5	9.4	159.1	58.8	23.3	5.5
45-64		Diabetic	111	3.0	84.9	6.3	29.6	9.7	156.6	69.3	28.7	8.5	
		Non-diabetic	161	3.5	90.1	8.4	28.5	9.4	157.7	64.1	26.4	6.2	
65-74		Diabetic	146	3.2	83.6	5.6	29.7	9.7	155.8	66.4	27.8	9.7	
		Non-diabetic	117	3.2	86.0	6.6	30.5	10.0	156.3	63.8	26.7	8.2	
75+		Diabetic	69	3.2	80.2	5.0	30.1	9.9	154.0	59.6	25.4	10.4	
		Non-diabetic	99	3.2	89.7	6.1	31.0	10.2	156.3	62.2	25.5	8.5	

Data are based on first & new Medical Evidence forms filed between May 1995 & June 2003.

The age of the incident patient is determined as of the date of ESRD initiation.

~Schwartz formula used for ages 0-18; Levey formula for ages 19+.

*Values for cells with 10 or fewer patients are suppressed.

." Zero values in this cell.



Table C.15 (continued)

Lab Values & Physical Characteristics: by Race

incident patients, by gender, age, & diabetic status, May 1995-April 2003 combined

INDIAN SUBCONTINENT			Serum					Body mass		~eGFR (ml/min)			
Age group	Diabetic status	Patients	albumin (g/dl)	BUN (mg/dl)	Creat. (mg/dl)	Hct %	Hgb (g/dl)	Height (cm)	Weight (kg)	index (kg/m ²)	by MDRD formula		
Male	0-19	Diabetic		
		Non-diabetic	34	3.4	98.0	9.0	30.4	10.3	148.3	44.3	20.0	10.7	
	20-44	Diabetic	39	3.0	88.0	8.5	27.7	9.4	170.2	73.0	25.9	8.5	
		Non-diabetic	195	3.4	91.5	11.3	28.8	9.7	171.3	67.1	23.1	6.8	
	45-64	Diabetic	390	3.1	86.2	7.6	29.3	9.8	167.7	67.9	24.6	9.2	
		Non-diabetic	270	3.5	92.3	10.0	28.2	9.5	168.4	64.9	22.9	6.9	
	65-74	Diabetic	178	3.2	85.5	6.6	29.9	9.9	166.6	66.4	24.0	10.5	
		Non-diabetic	125	3.3	86.3	7.6	29.5	10.0	165.9	61.9	23.3	9.5	
	75+	Diabetic	44	3.2	82.6	6.2	31.2	10.4	168.2	63.6	22.5	10.4	
		Non-diabetic	92	3.2	88.7	6.6	31.0	10.1	164.4	58.4	22.1	10.2	
	Female	0-19	Diabetic	
			Non-diabetic	13	3.5	78.4	7.3	31.5	10.5	132.6	35.5	19.8	9.5
		20-44	Diabetic	27	2.6	82.9	8.2	28.0	9.3	157.4	61.1	25.2	6.9
			Non-diabetic	123	3.4	86.1	9.1	27.2	9.0	157.6	56.3	23.0	6.7
45-64		Diabetic	170	3.0	80.0	6.7	28.8	9.5	156.9	61.2	26.2	8.1	
		Non-diabetic	117	3.3	85.9	7.9	28.3	9.4	155.5	55.2	24.0	6.8	
65-74		Diabetic	109	3.2	76.5	5.7	30.3	10.0	155.2	58.9	25.2	9.7	
		Non-diabetic	90	3.2	77.3	6.6	28.9	9.6	153.2	55.0	24.2	8.2	
75+		Diabetic	46	2.8	70.3	5.0	30.3	9.9	153.9	55.2	24.0	10.9	
		Non-diabetic	53	3.0	81.1	5.9	27.7	9.3	154.6	54.9	23.1	8.5	
OTHER/MULTIRACIAL													
Male		0-19	Diabetic	*	2.4	114.5	9.5	28.6	9.3	170.2	81.0	28.3	8.1
			Non-diabetic	271	3.5	97.0	10.0	27.5	9.3	142.8	46.7	21.7	9.5
		20-44	Diabetic	618	2.9	87.4	9.0	27.9	9.4	169.2	75.8	26.5	8.3
	Non-diabetic		1,578	3.3	102.8	12.8	27.1	9.2	169.0	72.2	25.6	5.9	
	45-64	Diabetic	3,187	3.0	90.1	8.2	28.7	9.6	167.6	74.7	26.9	8.5	
		Non-diabetic	1,716	3.3	98.0	10.4	28.1	9.5	168.1	72.6	26.1	6.8	
	65-74	Diabetic	1,433	3.1	91.1	7.3	29.1	9.8	166.8	69.8	25.7	9.3	
		Non-diabetic	818	3.3	95.8	8.8	28.9	9.7	166.0	68.2	24.8	7.7	
	75+	Diabetic	609	3.2	89.7	6.7	29.5	9.8	165.8	67.3	25.0	10.0	
		Non-diabetic	749	3.2	92.4	7.8	29.5	9.8	165.4	63.0	23.3	8.8	
	Female	0-19	Diabetic	*	3.8	81.0	12.1	32.1	11.0	153.5	54.0	22.8	4.9
			Non-diabetic	214	3.2	89.4	8.6	26.1	8.8	134.2	42.1	21.6	9.1
		20-44	Diabetic	396	2.8	78.6	7.5	27.5	9.2	159.2	67.9	27.1	7.9
			Non-diabetic	1,047	3.3	90.3	10.1	26.6	8.9	158.0	62.6	25.5	6.0
45-64		Diabetic	2,403	2.9	82.3	6.8	28.4	9.4	156.7	67.6	28.0	7.8	
		Non-diabetic	1,072	3.3	91.8	9.0	27.9	9.3	156.5	63.8	26.2	6.1	
65-74		Diabetic	1,551	3.0	82.6	6.3	28.8	9.6	155.7	62.9	26.4	8.2	
		Non-diabetic	585	3.2	84.3	7.5	29.0	9.7	155.4	58.5	24.6	7.1	
75+		Diabetic	677	3.1	83.3	5.9	29.4	9.8	154.8	58.9	25.0	8.9	
		Non-diabetic	606	3.2	85.4	6.7	29.4	9.8	153.9	54.3	23.4	7.9	
UNKNOWN													
Male		0-19	Diabetic	
			Non-diabetic	42	3.5	80.9	8.8	27.4	9.5	139.0	43.2	20.1	10.1
		20-44	Diabetic	88	3.0	86.1	9.1	28.0	9.6	170.8	76.8	26.6	8.3
	Non-diabetic		210	3.2	102.9	12.4	27.3	9.2	168.9	73.2	26.0	6.4	
	45-64	Diabetic	344	3.1	98.4	8.6	28.3	9.4	169.7	77.7	27.3	8.4	
		Non-diabetic	247	3.2	100.8	9.9	28.1	9.5	169.4	74.4	26.5	7.4	
	65-74	Diabetic	175	3.1	96.5	7.1	29.2	9.8	169.1	72.0	25.2	9.9	
		Non-diabetic	185	3.3	99.3	8.1	29.1	9.8	171.8	70.5	24.5	8.7	
	75+	Diabetic	90	3.1	95.7	6.6	30.6	10.1	168.6	68.7	24.4	10.4	
		Non-diabetic	175	3.1	93.0	7.0	29.6	9.9	167.8	68.3	23.7	9.5	
	Female	0-19	Diabetic	
			Non-diabetic	24	3.4	88.0	9.2	27.4	9.6	135.2	44.8	24.4	8.2
		20-44	Diabetic	59	2.8	78.7	7.3	27.9	9.4	162.0	73.0	28.1	8.0
			Non-diabetic	149	3.2	92.0	10.1	26.1	8.9	159.5	63.3	25.4	6.3
45-64		Diabetic	290	3.0	83.2	6.7	28.3	9.4	157.3	71.6	29.1	8.1	
		Non-diabetic	197	3.2	89.6	8.4	28.3	9.3	159.9	67.0	26.3	6.7	
65-74		Diabetic	224	3.0	85.4	6.4	28.9	9.7	157.1	65.9	26.9	8.8	
		Non-diabetic	98	3.3	86.0	7.2	29.2	9.6	159.5	61.2	24.2	7.3	
75+		Diabetic	89	3.1	86.7	5.8	28.9	9.7	157.0	60.4	24.7	9.0	
		Non-diabetic	154	3.1	91.2	6.6	28.9	9.7	155.7	59.6	24.6	8.1	

Data are based on first & new Medical Evidence forms filed between May 1995 & June 2003.

The age of the incident patient is determined as of the date of ESRD initiation.

~Schwartz formula used for ages 0-18; Levey formula for ages 19+.

*Values for cells with 10 or fewer patients are suppressed.

"." Zero values in this cell.

Table C.16
Lab Values & Physical Characteristics: by Hispanic Ethnicity
incident patients, by gender, age, & diabetic status

HISPANIC-MEXICAN			Serum							Body mass		~eGFR (ml/min)	
	Age group	Diabetes	Patients	albumin (g/dl)	BUN (mg/dl)	Creat. (mg/dl)	Hct %	Hgb (g/dl)	Height (cm)	Weight (kg)	index (kg/m ²)	by MDRD formula	
Male	0-19	Diabetic	*	3.3	96.0	11.2	27.8	8.8	158.4	62.7	25.6	8.1	
		Non-diabetic	672	3.4	97.1	10.1	27.3	9.3	144.6	45.1	21.0	9.8	
	20-44	Diabetic	1,680	2.8	84.5	8.7	28.1	9.5	170.3	75.9	27.1	8.7	
		Non-diabetic	3,431	3.3	103.6	12.7	27.0	9.2	169.1	71.7	25.8	6.1	
	45-64	Diabetic	8,585	3.0	88.2	7.9	28.7	9.6	169.0	74.5	27.0	8.9	
		Non-diabetic	3,184	3.2	94.3	9.7	28.1	9.5	169.1	72.6	26.3	7.5	
	65-74	Diabetic	4,031	3.1	88.7	7.0	29.2	9.8	168.0	69.2	25.8	9.9	
		Non-diabetic	1,986	3.2	91.2	8.1	29.2	9.8	168.6	66.7	24.8	8.6	
	75+	Diabetic	1,740	3.1	89.8	6.1	29.7	10.0	166.7	65.6	24.8	11.0	
		Non-diabetic	1,787	3.2	88.2	7.0	29.8	9.9	167.0	62.9	23.7	9.8	
	Female	0-19	Diabetic	*	2.9	74.1	5.9	28.9	9.5	159.0	58.0	23.1	12.7
			Non-diabetic	550	3.2	87.9	8.9	26.4	8.9	138.4	41.4	21.0	9.5
		20-44	Diabetic	1,214	2.7	77.9	7.3	27.7	9.3	158.5	67.5	27.8	8.1
			Non-diabetic	2,315	3.3	90.1	9.9	26.6	8.9	157.5	62.1	26.0	6.2
45-64		Diabetic	7,961	2.9	80.8	6.5	28.6	9.5	156.7	67.2	28.4	8.4	
		Non-diabetic	2,242	3.2	86.2	8.2	28.2	9.4	157.1	64.5	27.0	6.9	
65-74		Diabetic	5,263	3.0	81.3	6.0	29.2	9.7	155.7	62.2	26.7	8.9	
		Non-diabetic	1,574	3.2	83.6	6.9	29.1	9.7	155.9	59.0	25.5	8.0	
75+		Diabetic	2,352	3.0	82.0	5.4	29.5	9.8	154.4	58.4	25.6	9.9	
		Non-diabetic	1,624	3.1	83.2	5.9	29.7	9.9	154.2	53.4	24.0	8.9	
HISPANIC-OTHER													
Male		0-19	Diabetic	*	2.2	82.9	10.7	26.4	8.9	170.5	63.9	25.4	9.9
			Non-diabetic	365	3.4	95.6	9.7	27.7	9.4	145.1	49.1	21.8	9.8
		20-44	Diabetic	1,038	3.0	86.2	8.8	28.4	9.5	169.2	77.4	27.3	8.5
	Non-diabetic		2,454	3.3	100.8	12.4	27.4	9.3	169.8	74.1	26.0	6.2	
	45-64	Diabetic	4,951	3.1	90.2	8.1	28.9	9.7	168.1	75.9	27.0	8.7	
		Non-diabetic	2,822	3.3	98.3	10.2	28.4	9.6	168.8	73.1	26.0	7.0	
	65-74	Diabetic	2,520	3.1	89.8	7.1	29.4	9.8	167.0	71.0	25.8	9.6	
		Non-diabetic	1,612	3.3	93.4	8.6	29.0	9.7	167.3	67.7	24.7	7.9	
	75+	Diabetic	1,128	3.2	90.2	6.6	30.0	10.0	166.9	68.5	24.8	10.2	
		Non-diabetic	1,581	3.2	92.4	7.6	29.6	9.8	166.4	63.9	23.5	8.9	
	Female	0-19	Diabetic	*	4.5	69.0	16.5	34.0	11.7	152.0	44.0	19.0	3.0
			Non-diabetic	297	3.2	88.9	8.8	27.2	9.2	138.7	42.0	21.5	9.1
		20-44	Diabetic	712	2.8	77.7	7.4	27.7	9.2	159.3	67.7	27.2	7.8
			Non-diabetic	1,685	3.3	89.1	9.9	26.8	9.0	158.7	63.0	25.3	6.1
45-64		Diabetic	3,718	2.9	81.6	6.7	28.7	9.5	157.3	69.2	28.3	8.0	
		Non-diabetic	1,903	3.3	90.2	8.7	28.1	9.3	157.5	63.6	26.0	6.3	
65-74		Diabetic	2,593	3.0	83.3	6.3	29.1	9.7	156.6	64.7	26.6	8.4	
		Non-diabetic	1,142	3.2	84.9	7.4	29.0	9.7	156.3	60.4	25.1	7.3	
75+		Diabetic	1,260	3.1	82.5	5.8	29.8	9.9	155.4	60.2	25.4	9.0	
		Non-diabetic	1,188	3.2	84.7	6.5	29.7	9.9	155.4	56.0	23.8	8.1	
NON-HISPANIC													
Male		0-19	Diabetic	66	3.2	84.2	7.6	28.0	9.5	169.5	70.5	25.0	11.9
			Non-diabetic	4,038	3.4	88.9	9.1	28.6	9.7	148.0	47.6	21.5	10.9
		20-44	Diabetic	15,166	3.0	84.3	8.5	28.9	9.7	174.6	78.6	26.5	9.4
	Non-diabetic		35,674	3.3	98.5	12.0	28.1	9.5	175.3	79.5	26.5	7.0	
	45-64	Diabetic	55,552	3.1	91.6	7.8	29.2	9.8	175.2	85.3	28.3	9.6	
		Non-diabetic	58,875	3.3	95.0	9.6	29.0	9.7	175.2	79.4	26.5	7.9	
	65-74	Diabetic	35,631	3.2	93.8	6.8	29.8	9.9	173.7	79.5	26.9	10.5	
		Non-diabetic	46,732	3.2	92.9	7.9	29.6	9.9	173.8	73.6	25.0	9.0	
	75+	Diabetic	21,476	3.2	93.8	6.1	30.1	10.0	172.3	73.9	25.4	11.3	
		Non-diabetic	56,882	3.2	92.8	6.9	30.1	10.0	172.2	69.0	23.9	10.0	
	Female	0-19	Diabetic	78	2.9	74.8	7.1	27.3	9.0	159.9	62.0	25.8	10.3
			Non-diabetic	3,155	3.2	85.8	8.8	27.2	9.2	144.6	47.3	22.2	9.9
		20-44	Diabetic	11,860	2.9	76.7	7.3	28.3	9.4	162.4	71.0	27.7	8.7
			Non-diabetic	24,918	3.2	84.5	9.8	27.3	9.2	162.2	69.0	27.0	6.7
45-64		Diabetic	54,086	3.0	83.9	6.7	28.9	9.6	161.8	78.6	30.5	8.7	
		Non-diabetic	42,730	3.3	85.9	8.2	28.7	9.6	161.7	69.8	27.3	7.2	
65-74		Diabetic	41,778	3.1	85.6	6.1	29.3	9.7	160.5	72.5	28.6	9.2	
		Non-diabetic	36,254	3.2	84.8	6.9	29.3	9.7	160.3	64.0	25.6	8.1	
75+		Diabetic	25,093	3.1	85.8	5.6	29.6	9.8	159.1	66.4	26.7	9.8	
		Non-diabetic	46,334	3.1	85.2	6.2	29.7	9.9	158.7	58.9	24.0	8.8	

The age of the incident patient is determined as of the date of ESRD initiation.
 Data are based on first & new Medical Evidence forms filed between May 1995 & June 2003.
 ~Schwartz-Levey formula used for ages 0-18; Levey formula for ages 19+.
 *Values for cells with 10 or fewer patients are suppressed.

