

Table of Contents

Section D: Treatment Modalities: Medicare Only

- D.1 Percentages & counts of reported ESRD patients: by treatment modality
incident & point prevalent patients
- D.2 Counts of incident ESRD patients, by initial mode of therapy: all dialysis
by demographic characteristics & primary diagnosis
- D.3 Counts of incident ESRD patients, by initial mode of therapy: hemodialysis
by demographic characteristics & primary diagnosis
- D.4 Counts of incident ESRD patients, by initial mode of therapy: peritoneal dialysis
by demographic characteristics & primary diagnosis
- D.5 Counts of incident ESRD patients, by initial mode of therapy: transplant
by demographic characteristics & primary diagnosis
- D.6 Counts of prevalent ESRD patients, by mode of therapy: all dialysis
point prevalent patients on December 31, by demographic characteristics & primary diagnosis
- D.7 Counts of prevalent ESRD patients, by mode of therapy: hemodialysis
point prevalent patients on December 31, by demographic characteristics & primary diagnosis
- D.8 Counts of prevalent ESRD patients, by mode of therapy: peritoneal dialysis
point prevalent patients on December 31, by demographic characteristics & primary diagnosis
- D.9 Counts of prevalent ESRD patients, by mode of therapy: transplant
point prevalent patients on December 31, by demographic characteristics & primary diagnosis
- D.15 Incident ESRD patients, 2004: by modality & state/territory
incident ESRD patients
- D.16 Prevalent ESRD patients, 2004: by modality & state/territory
point prevalent ESRD patients on December 31

Table D.1
Percentages & counts of reported ESRD patients: by treatment modality
incident & point prevalent patients

INCIDENT COUNTS	1980	1981	1982	1983	1984	1985	1986	1987	1988	1989	1990	1991	1992
Modality													
Dialysis	16,788	18,695	20,911	24,167	25,606	28,317	30,906	33,800	37,296	42,129	45,871	50,970	55,415
Functioning graft	277	279	342	315	348	432	586	680	676	756	806	814	834
Dialysis type													
Center hemodialysis	14,746	15,934	17,212	19,960	21,084	22,640	24,912	27,357	30,660	34,839	37,706	41,980	45,921
Center self hemodialysis	60	56
Home hemodialysis	156	142	152	173	203	378	359	372	419	501	529	668	771
CAPD	933	1,805	3,029	3,591	3,751	3,843	3,910	4,190	4,452	4,842	5,560	5,763	5,876
CCPD	*	*	*	11	32	131	211	322	291	432	565	787	1,017
Other PD	637	402	218	241	239	532	691	626	683	742	624	731	708
Uncertain dialysis	313	407	295	191	297	793	823	933	791	773	887	981	1,066
All	17,065	18,974	21,253	24,482	25,954	28,749	31,492	34,480	37,972	42,885	46,677	51,784	56,249
PERCENTAGES													
Modality													
Dialysis	98.4	98.5	98.4	98.7	98.7	98.5	98.1	98.0	98.2	98.2	98.3	98.4	98.5
Functioning graft	1.6	1.5	1.6	1.3	1.3	1.5	1.9	2.0	1.8	1.8	1.7	1.6	1.5
Dialysis type													
Center hemodialysis	87.8	85.2	82.3	82.6	82.3	80.0	80.6	80.9	82.2	82.7	82.2	82.4	82.9
Center self hemodialysis	0.1	0.1
Home hemo	0.9	0.8	0.7	0.7	0.8	1.3	1.2	1.1	1.1	1.2	1.2	1.3	1.4
CAPD	5.6	9.7	14.5	14.9	14.6	13.6	12.7	12.4	11.9	11.5	12.1	11.3	10.6
CCPD	0.0	0.0	0.0	0.0	0.1	0.5	0.7	1.0	0.8	1.0	1.2	1.5	1.8
Other PD	3.8	2.2	1.0	1.0	0.9	1.9	2.2	1.9	1.8	1.8	1.4	1.4	1.3
Uncertain dialysis	1.9	2.2	1.4	0.8	1.2	2.8	2.7	2.8	2.1	1.8	1.9	1.9	1.9
All	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0
POINT PREVALENT COUNTS													
Modality													
Dialysis	47,656	53,561	59,938	69,411	76,354	82,528	88,920	96,680	105,416	116,830	129,073	144,026	158,494
Functioning graft	8,110	9,863	11,746	13,958	16,669	19,889	23,806	27,333	29,888	32,496	35,971	39,815	43,619
Dialysis type													
Center hemodialysis	42,547	48,313	53,128	59,949	63,470	64,496	70,030	77,399	86,427	96,215	106,043	117,190	128,831
Center self hemodialysis	681	763
Home hemodialysis	1,988	1,002	857	1,461	2,694	5,771	5,498	4,722	3,128	3,079	2,898	3,141	3,387
CAPD	811	2,474	4,162	6,490	8,441	10,204	11,113	12,008	13,008	14,488	16,335	18,245	20,011
CCPD	*	15	41	61	188	523	764	973	1,082	1,294	1,913	2,817	3,587
Other PD	1,371	1,042	956	791	703	784	702	617	578	538	429	368	317
Uncertain dialysis	936	715	794	659	858	750	813	961	1,193	1,216	1,455	1,584	1,598
All	55,766	63,424	71,684	83,369	93,023	102,417	112,726	124,013	135,304	149,326	165,044	183,841	202,113
PERCENTAGES													
Modality													
Dialysis	85.5	84.4	83.6	83.3	82.1	80.6	78.9	78.0	77.9	78.2	78.2	78.3	78.4
Functioning graft	14.5	15.6	16.4	16.7	17.9	19.4	21.1	22.0	22.1	21.8	21.8	21.7	21.6
Dialysis type													
Center hemodialysis	89.3	90.2	88.6	86.4	83.1	78.2	78.8	80.1	82.0	82.4	82.2	81.4	81.3
Center self hemodialysis	0.5	0.5
Home hemodialysis	4.2	1.9	1.4	2.1	3.5	7.0	6.2	4.9	3.0	2.6	2.2	2.2	2.1
CAPD	1.7	4.6	6.9	9.4	11.1	12.4	12.5	12.4	12.3	12.4	12.7	12.7	12.6
CCPD	0.0	0.0	0.1	0.1	0.2	0.6	0.9	1.0	1.0	1.1	1.5	2.0	2.3
Other PD	2.9	1.9	1.6	1.1	0.9	0.9	0.8	0.6	0.5	0.5	0.3	0.3	0.2
Uncertain dialysis	2.0	1.3	1.3	0.9	1.1	0.9	0.9	1.0	1.1	1.0	1.1	1.1	1.0
All	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0

The incident cohorts and associated modalities are determined at the time of ESRD initiation without applying the 60-day stable modality rule.

The point prevalent cohorts and associated modalities are determined on December 31 of each calendar year without applying the 60-day stable modality rule.

* Values for cells with 10 or fewer patients are suppressed.

." Zero values in this cell.

Table D.1
Percentages & counts of reported ESRD patients: by treatment modality
incident & point prevalent patients

INCIDENT COUNTS	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004
Modality												
Dialysis	57,793	62,129	63,244	68,442	71,555	75,529	78,825	80,754	82,758	86,741	88,080	87,831
Functioning graft	792	776	856	824	765	757	808	965	1,099	1,224	1,325	1,539
Dialysis type												
Center hemodialysis	48,056	51,978	53,686	58,869	63,553	67,799	70,825	73,314	76,037	80,141	81,553	81,511
Center self hemodialysis	41	24	52	57	48	46	42	30	37	44	39	25
Home hemodialysis	868	883	528	568	292	482	589	210	213	187	211	198
CAPD	6,226	6,171	5,981	5,520	4,902	4,452	4,283	4,054	4,083	3,864	3,970	3,847
CCPD	953	1,180	1,765	2,007	1,835	1,887	1,912	1,907	1,937	1,858	1,753	1,766
Other PD	630	592	450	434	406	378	250	296	268	406	283	268
Uncertain dialysis	1,019	1,301	782	987	519	485	924	943	183	241	271	216
All	58,585	62,905	64,100	69,266	72,320	76,286	79,633	81,719	83,857	87,965	89,405	89,370
PERCENTAGES												
Modality												
Dialysis	98.6	98.8	98.7	98.8	98.9	99.0	99.0	98.8	98.7	98.6	98.5	98.3
Functioning graft	1.4	1.2	1.3	1.2	1.1	1.0	1.0	1.2	1.3	1.4	1.5	1.7
Dialysis type												
Center hemodialysis	83.2	83.7	84.9	86.0	88.8	89.8	89.9	90.8	91.9	92.4	92.6	92.8
Center self hemodialysis	0.1	0.0	0.1	0.1	0.1	0.1	0.1	0.0	0.0	0.1	0.0	0.0
Home hemo	1.5	1.4	0.8	0.8	0.4	0.6	0.7	0.3	0.3	0.2	0.2	0.2
CAPD	10.8	9.9	9.5	8.1	6.9	5.9	5.4	5.0	4.9	4.5	4.5	4.4
CCPD	1.6	1.9	2.8	2.9	2.6	2.5	2.4	2.4	2.3	2.1	2.0	2.0
Other PD	1.1	1.0	0.7	0.6	0.6	0.5	0.3	0.4	0.3	0.5	0.3	0.3
Uncertain dialysis	1.8	2.1	1.2	1.4	0.7	0.6	1.2	1.2	0.2	0.3	0.3	0.2
All	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0
POINT PREVALENT COUNTS												
Modality												
Dialysis	171,562	186,203	198,293	212,295	225,107	238,252	250,997	263,329	274,279	285,728	295,385	302,492
Functioning graft	47,800	50,339	54,846	59,427	63,559	68,018	72,791	76,895	81,115	85,831	88,220	93,875
Dialysis type												
Center hemodialysis	140,186	152,621	164,581	179,243	194,438	209,970	223,269	236,522	248,310	259,681	268,991	276,438
Center self hemodialysis	132	195	513	529	466	361	341	271	323	277	196	193
Home hemodialysis	3,755	3,728	2,974	2,829	2,474	2,378	2,230	2,024	1,756	1,682	1,833	1,829
CAPD	21,634	22,597	21,181	18,864	16,587	14,490	13,054	11,936	11,151	10,355	10,017	9,511
CCPD	4,183	5,228	7,565	9,169	9,905	9,965	10,639	11,237	12,029	12,997	13,614	13,893
Other PD	228	207	165	132	102	88	64	56	46	65	42	39
Uncertain dialysis	1,444	1,627	1,314	1,529	1,135	1,000	1,400	1,283	664	671	692	589
All	219,362	236,542	253,139	271,722	288,666	306,270	323,788	340,224	355,394	371,559	383,605	396,367
PERCENTAGES												
Modality												
Dialysis	78.2	78.7	78.3	78.1	78.0	77.8	77.5	77.4	77.2	76.9	77.0	76.3
Functioning graft	21.8	21.3	21.7	21.9	22.0	22.2	22.5	22.6	22.8	23.1	23.0	23.7
Dialysis type												
Center hemodialysis	81.7	82.0	83.0	84.4	86.4	88.1	89.0	89.8	90.5	90.9	91.1	91.4
Center self hemodialysis	0.1	0.1	0.3	0.2	0.2	0.2	0.1	0.1	0.1	0.1	0.1	0.1
Home hemodialysis	2.2	2.0	1.5	1.3	1.1	1.0	0.9	0.8	0.6	0.6	0.6	0.6
CAPD	12.6	12.1	10.7	8.9	7.4	6.1	5.2	4.5	4.1	3.6	3.4	3.1
CCPD	2.4	2.8	3.8	4.3	4.4	4.2	4.2	4.3	4.4	4.5	4.6	4.6
Other PD	0.1	0.1	0.1	0.1	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0
Uncertain dialysis	0.8	0.9	0.7	0.7	0.5	0.4	0.6	0.5	0.2	0.2	0.2	0.2
All	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0

The incident cohorts and associated modalities are determined at the time of ESRD initiation without applying the 60-day stable modality rule.

The point prevalent cohorts and associated modalities are determined on December 31 of each calendar year without applying the 60-day stable modality rule.

* Values for cells with 10 or fewer patients are suppressed.

." Zero values in this cell.

Table D.2

Counts of incident ESRD patients, by initial mode of therapy: all dialysis

by demographic characteristics & primary diagnosis

	1980	1981	1982	1983	1984	1985	1986	1987	1988	1989	1990	1991	1992
0-4	34	52	75	84	86	109	114	98	104	109	96	124	106
5-9	78	63	82	74	84	73	92	81	69	74	101	74	66
10-14	147	153	188	164	187	168	152	144	154	132	152	164	161
15-19	375	407	380	367	353	359	361	380	391	416	384	355	403
20-29	1,333	1,490	1,660	1,612	1,751	1,781	1,840	1,819	1,910	2,027	2,065	2,119	2,176
30-39	1,991	2,233	2,477	2,541	2,721	2,900	3,186	3,220	3,519	3,826	3,975	4,146	4,432
40-49	2,254	2,547	2,623	2,935	3,075	3,331	3,587	3,991	4,306	4,914	5,290	5,921	6,338
50-59	3,537	4,105	4,365	4,507	4,732	5,057	5,307	5,644	6,340	6,834	7,336	7,896	8,660
60-64	2,008	2,293	2,563	2,984	3,273	3,663	3,771	4,200	4,572	4,930	5,387	5,950	6,170
65-69	2,136	2,212	2,543	3,270	3,438	3,843	4,371	4,995	5,518	6,389	7,081	7,864	8,449
70-79	2,472	2,669	3,315	4,617	4,828	5,662	6,483	7,314	8,154	9,614	10,699	12,526	13,978
80+	423	470	637	1,007	1,072	1,359	1,637	1,909	2,257	2,857	3,302	3,831	4,475
Missing/unknown	.	*	*	*	*	12	*	*	*	*	*	.	*
0-19	634	675	725	689	710	709	719	703	718	731	733	717	736
20-44	4,356	4,927	5,358	5,515	5,891	6,226	6,711	6,959	7,486	8,148	8,644	9,128	9,593
45-64	6,767	7,741	8,330	9,064	9,661	10,506	10,980	11,915	13,161	14,383	15,409	16,904	18,183
65-74	3,699	3,910	4,652	6,101	6,313	7,252	8,133	9,279	10,202	11,811	13,239	15,051	16,475
75+	1,332	1,441	1,843	2,793	3,025	3,612	4,358	4,939	5,727	7,049	7,843	9,170	10,427
Unknown	.	*	*	*	*	12	*	*	*	*	*	.	*
Male	9,372	10,376	11,538	13,318	14,157	15,414	16,955	18,388	20,143	22,864	24,828	27,348	29,501
Female	7,416	8,319	9,373	10,848	11,449	12,903	13,951	15,412	17,153	19,265	21,043	23,622	25,914
Unknown	.	.	.	*
White	11,774	13,090	14,679	16,602	17,598	19,476	21,305	23,057	25,212	28,440	30,999	34,235	37,066
Black	4,552	5,107	5,746	6,949	7,311	8,026	8,641	9,686	10,791	12,382	13,306	14,968	16,322
Native American	95	122	151	227	235	250	314	364	434	482	508	547	654
Asian	94	144	281	315	409	511	580	641	770	800	1,039	1,201	1,342
Other/unknown	273	232	54	74	53	54	66	52	89	25	19	19	31
†Hispanic													
†Non-Hispanic													
Diabetes	2,498	3,790	4,957	5,819	7,019	8,129	9,393	10,648	12,010	14,503	16,618	19,081	21,099
Hypertension	2,956	4,279	5,518	6,040	6,792	7,788	8,480	9,744	11,108	13,020	14,344	15,960	17,759
Glomerulonephritis	2,586	3,383	3,965	4,174	4,492	4,682	4,918	5,296	5,601	5,911	6,234	6,468	6,618
Cystic kidney	731	887	1,021	1,048	1,081	1,148	1,234	1,244	1,279	1,267	1,405	1,452	1,530
Other urologic	420	617	803	904	894	1,112	1,115	1,222	1,231	1,258	1,151	1,338	1,395
Other cause	1,619	2,024	2,568	2,842	3,024	3,263	3,322	3,307	3,535	4,098	4,177	4,574	4,933
Unknown cause	1,468	1,512	1,233	1,251	1,299	1,347	1,567	1,720	1,749	1,515	1,593	1,787	1,848
Missing cause	4,510	2,203	846	2,089	1,005	848	877	619	783	557	349	310	233
All	16,788	18,695	20,911	24,167	25,606	28,317	30,906	33,800	37,296	42,129	45,871	50,970	55,415

The incident cohorts and associated modalities are determined at the time of ESRD initiation without applying the 60-day stable modality rule.

The age of the point prevalent patient is determined as of December 31 of each calendar year.

†CMS began collecting Hispanic ethnicity data in April 1995; cells for years prior to 1996 have therefore been left blank.

* Values for cells with 10 or fewer patients are suppressed.

." Zero values in this cell.

Table D.2

Counts of incident ESRD patients, by initial mode of therapy: all dialysis

by demographic characteristics & primary diagnosis

	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004
0-4	110	96	106	108	97	89	80	104	92	114	99	116
5-9	84	97	76	74	59	66	77	73	74	91	63	77
10-14	164	181	171	180	163	152	145	163	152	198	180	183
15-19	384	358	380	383	315	356	359	328	351	340	360	343
20-29	2,173	2,095	2,008	2,018	1,874	1,899	1,793	1,843	1,871	1,729	1,709	1,678
30-39	4,404	4,535	4,334	4,367	4,127	4,081	4,206	4,054	4,070	4,121	4,018	3,953
40-49	6,636	6,843	7,164	7,517	7,287	7,863	7,813	8,007	7,989	8,287	8,536	8,461
50-59	9,013	10,027	9,948	10,341	11,036	11,852	12,602	12,930	13,584	14,119	14,545	14,625
60-64	6,418	6,644	6,868	7,145	7,339	7,741	7,856	8,134	8,397	8,544	8,981	9,018
65-69	8,764	9,372	9,336	10,174	10,552	10,799	10,846	10,961	11,188	11,423	11,635	11,436
70-79	14,760	16,334	16,843	19,012	20,450	21,512	22,803	23,158	23,452	24,525	24,338	23,922
80+	4,883	5,544	6,008	7,123	8,255	9,119	10,245	10,999	11,538	13,249	13,616	14,018
Missing/unknown	.	*	*	.	*	*	.	*
0-19	742	732	733	745	634	663	661	668	669	743	702	719
20-44	9,596	9,635	9,567	9,715	9,183	9,373	9,285	9,293	9,327	9,264	9,258	9,096
45-64	19,048	20,509	20,755	21,673	22,480	24,063	24,985	25,675	26,584	27,536	28,531	28,639
65-74	17,249	18,700	18,856	20,817	21,689	22,287	22,658	22,919	23,310	23,871	23,742	23,421
75+	11,158	12,550	13,331	15,492	17,568	19,143	21,236	22,199	22,868	25,326	25,847	25,955
Unknown	.	*	*	.	*	*	.	*
Male	30,995	33,016	33,558	36,778	38,248	40,435	42,411	43,385	44,719	47,338	48,027	48,717
Female	26,798	29,113	29,686	31,664	33,307	35,094	36,412	37,369	38,039	39,403	40,045	39,100
Unknown	*	.	.	.	*	14
White	38,331	41,176	41,318	45,064	47,336	49,584	52,418	53,982	55,054	57,608	58,436	58,710
Black	17,167	18,316	18,714	19,793	20,631	22,061	22,221	22,492	23,304	24,450	24,812	23,984
Native American	657	711	747	770	805	901	924	956	952	970	931	992
Asian	1,592	1,803	1,853	2,046	2,072	2,329	2,579	2,557	2,686	2,890	2,902	2,945
Other/unknown	46	123	612	769	711	654	683	767	762	823	999	1,200
†Hispanic				7,461	7,552	8,001	8,712	10,157	9,370	9,930	10,559	10,415
†Non-Hispanic				60,981	64,003	67,528	70,113	70,597	73,388	76,811	77,521	77,416
Diabetes	22,442	24,832	26,914	30,045	31,970	34,089	35,736	37,184	38,151	39,490	40,027	39,779
Hypertension	18,333	19,690	17,594	18,621	19,874	20,925	22,009	22,314	23,099	24,544	25,373	24,753
Glomerulonephritis	6,803	7,248	7,146	7,500	7,241	7,262	7,212	6,932	6,829	7,025	6,755	6,602
Cystic kidney	1,578	1,624	1,632	1,512	1,570	1,623	1,682	1,632	1,672	1,684	1,672	1,631
Other urologic	1,190	1,233	1,760	2,021	2,012	2,036	2,354	2,223	2,305	2,443	2,341	2,325
Other cause	5,174	5,419	5,886	6,087	6,206	6,636	6,801	7,159	7,247	7,951	8,056	8,290
Unknown cause	1,884	1,937	2,169	2,525	2,576	2,823	2,885	3,155	3,331	3,345	3,415	3,262
Missing cause	389	146	143	131	106	135	146	155	124	259	441	1,189
All	57,793	62,129	63,244	68,442	71,555	75,529	78,825	80,754	82,758	86,741	88,080	87,831

The incident cohorts and associated modalities are determined at the time of ESRD initiation without applying the 60-day stable modality rule.

The age of the point prevalent patient is determined as of December 31 of each calendar year.

†CMS began collecting Hispanic ethnicity data in April 1995; cells for years prior to 1996 have therefore been left blank.

* Values for cells with 10 or fewer patients are suppressed.

." Zero values in this cell.

Table D.3

Counts of incident ESRD patients, by initial mode of therapy: hemodialysis

by demographic characteristics & primary diagnosis

	1980	1981	1982	1983	1984	1985	1986	1987	1988	1989	1990	1991	1992
0-4	16	16	21	24	16	29	34	15	20	31	26	30	18
5-9	53	39	46	32	29	27	28	27	21	25	39	27	24
10-14	123	111	106	89	95	93	78	74	71	71	71	64	84
15-19	346	362	305	289	273	267	267	269	291	307	267	270	274
20-29	1,194	1,296	1,405	1,351	1,429	1,471	1,472	1,437	1,495	1,611	1,593	1,669	1,712
30-39	1,733	1,894	2,017	2,008	2,191	2,258	2,495	2,502	2,741	2,961	3,043	3,140	3,412
40-49	2,003	2,232	2,149	2,386	2,508	2,656	2,831	3,176	3,447	3,851	4,160	4,565	4,987
50-59	3,128	3,540	3,654	3,730	3,902	4,116	4,346	4,613	5,279	5,612	5,966	6,473	7,046
60-64	1,825	1,958	2,158	2,506	2,760	3,010	3,143	3,512	3,842	4,174	4,488	4,957	5,147
65-69	1,933	1,903	2,157	2,852	2,934	3,170	3,663	4,199	4,757	5,548	6,101	6,847	7,447
70-79	2,183	2,329	2,808	3,999	4,222	4,767	5,520	6,239	7,120	8,551	9,492	11,151	12,487
80+	365	395	536	862	923	1,143	1,390	1,662	1,993	2,593	2,986	3,515	4,110
Missing/unknown	.	*	*	*	*	11	*	*	*	*	*	.	.
0-19	538	528	478	434	413	416	407	385	403	434	403	391	400
20-44	3,848	4,240	4,412	4,454	4,776	4,960	5,280	5,438	5,872	6,363	6,663	6,968	7,462
45-64	6,035	6,680	6,971	7,527	8,014	8,551	9,007	9,802	10,932	11,846	12,587	13,836	14,842
65-74	3,314	3,370	3,942	5,298	5,422	6,047	6,856	7,826	8,820	10,357	11,514	13,209	14,569
75+	1,167	1,257	1,559	2,415	2,657	3,033	3,717	4,274	5,050	6,335	7,065	8,304	9,475
Unknown	.	*	*	*	*	11	*	*	*	*	*	.	.
Male	8,393	9,036	9,680	11,168	11,769	12,597	13,895	15,086	16,822	19,149	20,711	22,780	24,806
Female	6,509	7,040	7,684	8,964	9,518	10,421	11,376	12,643	14,257	16,191	17,524	19,928	21,942
Unknown	.	.	.	*
White	10,320	11,073	12,012	13,595	14,395	15,433	17,057	18,518	20,550	23,362	25,256	28,044	30,598
Black	4,169	4,541	4,915	5,992	6,297	6,868	7,380	8,282	9,395	10,841	11,592	13,118	14,391
Native American	83	109	131	193	194	213	261	308	369	412	429	463	553
Asian	88	134	254	284	355	461	523	579	697	703	948	1,067	1,177
Other/unknown	242	219	52	69	46	43	50	42	68	22	*	16	29
†Hispanic													
†Non-Hispanic													
Diabetes	2,075	3,108	3,906	4,615	5,652	6,456	7,551	8,606	9,929	12,043	13,732	15,839	17,588
Hypertension	2,622	3,746	4,664	5,146	5,847	6,552	7,195	8,341	9,603	11,338	12,426	13,959	15,688
Glomerulonephritis	2,271	2,948	3,309	3,476	3,709	3,780	3,972	4,237	4,491	4,756	4,965	5,163	5,253
Cystic kidney	667	754	872	882	895	945	998	995	1,017	999	1,077	1,084	1,167
Other urologic	371	538	669	768	759	915	944	1,048	1,074	1,114	995	1,179	1,201
Other cause	1,426	1,751	2,173	2,339	2,495	2,613	2,687	2,629	2,918	3,392	3,432	3,735	4,080
Unknown cause	1,313	1,258	1,023	1,039	1,072	1,102	1,256	1,418	1,457	1,275	1,339	1,513	1,596
Missing cause	4,157	1,973	748	1,868	858	655	668	455	590	423	269	236	175
All	14,902	16,076	17,364	20,133	21,287	23,018	25,271	27,729	31,079	35,340	38,235	42,708	46,748

The incident cohorts and associated modalities are determined at the time of ESRD initiation without applying the 60-day stable modality rule.

The age of the incident patient is determined as of the date of ESRD initiation.

†CMS began collecting Hispanic ethnicity data in April 1995; cells for years prior to 1996 have therefore been left blank.

* Values for cells with 10 or fewer patients are suppressed.

." Zero values in this cell.

Table D.3

Counts of incident ESRD patients, by initial mode of therapy: hemodialysis*by demographic characteristics & primary diagnosis*

	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004
0-4	27	16	22	17	26	16	18	17	21	25	23	20
5-9	28	33	30	28	19	28	27	29	29	37	24	37
10-14	77	89	86	80	79	81	72	88	90	109	97	100
15-19	269	260	270	281	229	279	286	248	267	270	286	256
20-29	1,708	1,672	1,624	1,643	1,558	1,631	1,552	1,598	1,641	1,545	1,521	1,488
30-39	3,412	3,491	3,407	3,561	3,454	3,455	3,597	3,520	3,575	3,617	3,570	3,487
40-49	5,194	5,462	5,717	6,193	6,208	6,864	6,840	7,037	7,106	7,519	7,678	7,651
50-59	7,440	8,249	8,267	8,726	9,556	10,418	11,207	11,615	12,179	12,800	13,283	13,373
60-64	5,458	5,602	5,817	6,231	6,518	6,998	7,084	7,376	7,715	7,852	8,276	8,389
65-69	7,647	8,263	8,195	9,030	9,603	9,875	9,924	10,018	10,431	10,632	10,828	10,677
70-79	13,236	14,657	15,258	17,124	18,869	20,012	21,153	21,557	22,175	23,219	23,093	22,733
80+	4,469	5,089	5,571	6,580	7,773	8,670	9,696	10,451	11,058	12,747	13,124	13,522
Missing/unknown	.	*	*	.	*	*
0-19	401	398	408	406	353	404	403	382	407	441	430	413
20-44	7,460	7,544	7,632	7,928	7,693	8,057	8,013	8,077	8,212	8,263	8,260	8,061
45-64	15,752	16,932	17,200	18,426	19,601	21,309	22,267	23,069	24,004	25,070	26,068	26,327
65-74	15,200	16,563	16,781	18,525	19,834	20,500	20,786	21,098	21,842	22,385	22,245	21,977
75+	10,152	11,446	12,243	14,209	16,411	18,057	19,987	20,928	21,822	24,213	24,800	24,955
Unknown	.	*	*	.	*	*
Male	26,265	28,068	28,748	31,980	34,212	36,670	38,512	39,604	41,314	43,924	44,645	45,408
Female	22,700	24,817	25,518	27,514	29,681	31,657	32,942	33,950	34,973	36,448	37,151	36,313
Unknown	*	.	.	.	*	13
White	31,846	34,404	34,811	38,376	41,621	44,272	46,848	48,631	50,275	52,977	53,898	54,229
Black	15,131	16,226	16,696	18,000	19,067	20,572	20,760	21,056	21,987	23,112	23,476	22,788
Native American	567	604	641	673	725	802	861	852	870	888	868	928
Asian	1,379	1,541	1,573	1,772	1,845	2,078	2,359	2,312	2,459	2,636	2,643	2,710
Other/unknown	42	110	545	673	635	603	628	703	696	759	918	1,079
†Hispanic				6,612	6,850	7,308	8,009	9,373	8,764	9,248	9,914	9,777
†Non-Hispanic				52,882	57,043	61,019	63,447	64,181	67,523	71,124	71,889	71,957
Diabetes	18,950	21,018	23,015	25,981	28,521	30,827	32,361	33,876	35,346	36,659	37,318	37,247
Hypertension	16,144	17,451	15,654	16,798	18,246	19,391	20,406	20,751	21,718	23,134	23,924	23,350
Glomerulonephritis	5,471	5,800	5,751	6,147	6,104	6,195	6,258	5,965	5,936	6,213	5,946	5,755
Cystic kidney	1,186	1,220	1,241	1,169	1,259	1,338	1,360	1,335	1,360	1,427	1,363	1,349
Other urologic	1,019	1,044	1,524	1,782	1,808	1,862	2,154	2,017	2,115	2,260	2,166	2,151
Other cause	4,292	4,570	5,048	5,278	5,541	5,992	6,139	6,553	6,607	7,335	7,463	7,724
Unknown cause	1,616	1,678	1,919	2,233	2,326	2,609	2,653	2,922	3,093	3,139	3,232	3,072
Missing cause	287	104	114	106	88	113	125	135	112	205	391	1,086
All	48,965	52,885	54,266	59,494	63,893	68,327	71,456	73,554	76,287	80,372	81,803	81,734

The incident cohorts and associated modalities are determined at the time of ESRD initiation without applying the 60-day stable modality rule.

The age of the incident patient is determined as of the date of ESRD initiation.

†CMS began collecting Hispanic ethnicity data in April 1995; cells for years prior to 1996 have therefore been left blank.

* Values for cells with 10 or fewer patients are suppressed.

." Zero values in this cell.

Table D.4

Counts of incident ESRD patients, by initial mode of therapy: peritoneal dialysis

by demographic characteristics & primary diagnosis

	1980	1981	1982	1983	1984	1985	1986	1987	1988	1989	1990	1991	1992
0-4	*	25	47	54	63	77	74	73	75	58	54	75	75
5-9	14	17	32	38	51	41	57	48	38	42	47	36	38
10-14	12	29	70	68	88	67	67	65	74	45	69	76	69
15-19	13	27	60	67	70	78	76	92	73	96	94	72	115
20-29	70	138	211	236	286	256	313	304	348	328	390	369	387
30-39	131	231	400	488	469	534	550	588	639	714	780	865	882
40-49	146	218	411	508	510	551	618	665	720	849	965	1,162	1,141
50-59	225	417	609	714	754	765	776	831	843	996	1,118	1,184	1,325
60-64	101	232	340	431	452	517	478	530	582	591	737	783	811
65-69	80	210	322	364	417	417	445	520	503	600	750	753	734
70-79	117	218	444	520	505	554	533	647	684	761	899	976	1,065
80+	18	48	87	114	117	117	134	148	164	192	222	199	250
Missing/unknown	.	.	*	.	*	.	.	*	.	*	.	.	*
0-19	48	98	209	227	272	263	274	278	260	241	264	259	297
20-44	268	477	812	970	989	1,038	1,165	1,236	1,345	1,444	1,665	1,841	1,823
45-64	405	759	1,159	1,407	1,482	1,585	1,570	1,682	1,787	2,034	2,325	2,522	2,723
65-74	157	361	602	689	737	754	766	928	925	1,039	1,314	1,345	1,375
75+	58	115	251	309	302	334	346	387	426	514	557	583	674
Unknown	.	.	*	.	*	.	.	*	.	*	.	.	*
Male	494	902	1,588	1,921	2,079	2,111	2,278	2,422	2,562	2,854	3,314	3,621	3,727
Female	442	908	1,446	1,681	1,704	1,863	1,843	2,090	2,181	2,420	2,811	2,929	3,166
Unknown
White	757	1,407	2,274	2,678	2,827	3,081	3,152	3,423	3,617	3,969	4,660	4,892	5,141
Black	150	378	718	861	865	809	865	990	1,002	1,164	1,309	1,473	1,522
Native American	*	*	17	32	37	34	50	49	59	57	68	68	79
Asian	*	*	24	26	50	43	47	47	62	82	84	117	151
Other/unknown	15	*	*	*	*	*	*	*	*	*	*	.	.
†Hispanic													
†Non-Hispanic													
Diabetes	249	514	920	1,091	1,196	1,297	1,397	1,577	1,615	1,969	2,308	2,588	2,825
Hypertension	199	384	771	820	819	872	879	954	1,121	1,244	1,535	1,548	1,589
Glomerulonephritis	200	328	563	630	703	732	758	825	896	940	1,049	1,106	1,158
Cystic kidney	42	98	127	147	170	169	193	200	213	220	272	312	314
Other urologic	23	58	109	118	112	129	116	116	111	100	120	118	134
Other cause	127	203	336	458	477	513	477	536	479	552	612	640	670
Unknown cause	80	174	178	189	196	157	200	189	180	157	185	199	183
Missing cause	16	51	30	149	110	105	101	115	128	92	44	39	20
All	936	1,810	3,034	3,602	3,783	3,974	4,121	4,512	4,743	5,274	6,125	6,550	6,893

The incident cohorts and associated modalities are determined at the time of ESRD initiation without applying the 60-day stable modality. The age of the incident patient is determined as of the date of ESRD initiation.

†CMS began collecting Hispanic ethnicity data in April 1995; cells for years prior to 1996 have therefore been left blank.

* Values for cells with 10 or fewer patients are suppressed.

." Zero values in this cell.

Table D.4

Counts of incident ESRD patients, by initial mode of therapy: peritoneal dialysis

by demographic characteristics & primary diagnosis

	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004
0-4	72	63	78	86	71	71	61	84	69	85	72	91
5-9	43	59	46	45	39	37	49	42	45	52	36	39
10-14	76	80	80	94	79	67	70	74	60	89	80	81
15-19	97	85	103	98	84	76	71	77	79	66	71	80
20-29	400	371	353	355	295	261	233	230	219	174	183	186
30-39	860	897	861	768	647	595	583	517	476	474	432	445
40-49	1,241	1,177	1,342	1,260	1,012	950	917	914	844	734	814	774
50-59	1,362	1,479	1,533	1,489	1,414	1,342	1,291	1,203	1,349	1,245	1,191	1,192
60-64	783	831	921	842	768	681	694	685	650	648	664	601
65-69	852	806	915	839	758	763	712	706	692	678	728	676
70-79	1,116	1,191	1,197	1,314	1,225	1,161	1,160	1,102	1,122	1,080	1,051	1,038
80+	277	312	317	337	345	335	354	327	415	396	401	410
Missing/unknown	*	.	.
0-19	288	287	307	323	273	251	251	277	253	292	259	291
20-44	1,842	1,812	1,793	1,699	1,414	1,258	1,220	1,164	1,071	944	954	997
45-64	2,804	2,943	3,217	3,015	2,722	2,571	2,498	2,385	2,467	2,331	2,330	2,201
65-74	1,538	1,553	1,630	1,667	1,468	1,448	1,386	1,325	1,328	1,265	1,326	1,276
75+	707	756	799	823	860	811	840	810	901	889	854	848
Unknown	*	.	.
Male	3,825	3,906	4,137	4,036	3,561	3,321	3,272	3,199	3,162	3,053	3,075	3,053
Female	3,354	3,445	3,609	3,491	3,176	3,018	2,923	2,762	2,858	2,669	2,648	2,560
Unknown
White	5,262	5,355	5,614	5,547	5,030	4,676	4,680	4,410	4,443	4,157	4,100	4,104
Black	1,646	1,658	1,723	1,549	1,357	1,315	1,233	1,205	1,216	1,199	1,243	1,123
Native American	74	87	90	88	70	77	56	83	76	62	58	58
Asian	195	242	259	255	206	222	180	205	221	245	250	215
Other/unknown	*	*	60	88	74	49	46	58	64	59	72	113
†Hispanic				756	656	631	645	709	582	639	586	581
†Non-Hispanic				6,771	6,081	5,708	5,550	5,252	5,438	5,083	5,137	5,032
Diabetes	2,854	3,044	3,369	3,442	3,055	2,848	2,852	2,717	2,627	2,553	2,473	2,349
Hypertension	1,744	1,732	1,591	1,423	1,355	1,308	1,257	1,218	1,256	1,253	1,314	1,289
Glomerulonephritis	1,138	1,212	1,290	1,218	1,074	995	871	877	861	760	768	815
Cystic kidney	353	353	352	311	282	267	297	273	293	247	297	267
Other urologic	138	126	183	185	169	152	160	166	177	166	148	153
Other cause	714	672	731	684	582	576	564	517	583	549	528	510
Unknown cause	205	195	218	251	212	188	186	185	217	174	171	176
Missing cause	33	17	12	13	*	*	*	*	*	20	24	54
All	7,179	7,351	7,746	7,527	6,737	6,339	6,195	5,961	6,020	5,722	5,723	5,613

The incident cohorts and associated modalities are determined at the time of ESRD initiation without applying the 60-day stable modality rule.

The age of the incident patient is determined as of the date of ESRD initiation.

†CMS began collecting Hispanic ethnicity data in April 1995; cells for years prior to 1996 have therefore been left blank.

* Values for cells with 10 or fewer patients are suppressed.

." Zero values in this cell.

Table D.5

Counts of incident ESRD patients, by initial mode of therapy: transplant

by demographic characteristics & primary diagnosis

	1980	1981	1982	1983	1984	1985	1986	1987	1988	1989	1990	1991	1992
0-4	*	11	16	11	12	14	*	15	16	16	14	14	16
5-9	*	16	17	*	11	18	18	19	13	27	24	28	26
10-14	14	11	12	16	26	24	36	43	32	35	41	44	33
15-19	21	21	25	23	26	30	44	50	35	40	49	35	35
20-29	69	56	70	65	60	96	104	105	129	157	140	121	121
30-39	72	90	85	88	99	112	188	182	201	201	225	228	229
40-49	59	42	66	64	63	75	101	133	133	160	184	211	202
50-59	23	26	48	30	36	39	59	95	88	84	93	94	124
60-64	*	*	*	*	*	*	13	22	17	22	18	24	28
65-69	*	*	.	*	*	*	*	*	*	*	13	12	16
70-79	.	*	.	*	*	*	*	*	*	*	*	*	*
80+	*	*	*	.	*	*	*	*	.	.	*	.	.
Missing/unknown
0-19	49	59	70	58	75	86	106	127	96	118	128	121	110
20-44	169	169	194	187	197	256	341	365	409	451	480	480	464
45-64	55	48	77	67	63	74	124	172	159	173	180	198	240
65-74	*	*	.	*	*	11	13	12	11	13	16	14	20
75+	*	*	*	.	*	*	*	*	*	*	*	*	.
Unknown
Male	182	186	225	202	222	257	355	403	398	445	464	473	463
Female	95	93	117	113	125	175	231	277	278	311	342	341	371
Unknown	*
White	225	232	280	260	302	381	522	596	593	683	717	746	742
Black	42	42	55	46	42	37	53	64	69	54	68	48	76
Native American	.	*	*	*	*	*	*	*	*	*	*	*	*
Asian	*	*	*	*	*	*	*	12	*	16	15	11	11
Other/unknown	*	*	*	*	*	*	*	*	*	*	.	.	.
†Hispanic													
†Non-Hispanic													
Diabetes	35	55	81	62	89	104	164	169	185	233	204	228	206
Hypertension	21	20	30	24	17	20	34	58	54	46	50	49	72
Glomerulonephritis	39	43	62	67	72	106	98	142	126	158	183	161	183
Cystic kidney	*	11	*	20	16	23	38	61	55	63	74	88	90
Other urologic	20	24	13	21	21	22	41	37	28	42	38	50	36
Other cause	34	32	46	39	59	59	80	102	85	105	128	118	127
Unknown cause	15	15	31	22	18	39	43	37	45	41	52	51	50
Missing cause	111	79	71	60	56	59	88	74	98	68	77	69	70
All	277	279	342	315	348	432	586	680	676	756	806	814	834

The incident cohorts and associated modalities are determined at the time of ESRD initiation without applying the 60-day rule. The age of the incident patient is determined as of the date of ESRD initiation.

†CMS began collecting Hispanic ethnicity data in April 1995; cells for years prior to 1996 have therefore been left blank.

* Values for cells with 10 or fewer patients are suppressed.

." Zero values in this cell.

Table D.5

Counts of incident ESRD patients, by initial mode of therapy: transplant*by demographic characteristics & primary diagnosis*

	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004
0-4	17	*	13	16	12	*	*	15	16	*	11	13
5-9	12	26	21	21	20	18	14	16	15	27	21	15
10-14	34	26	33	23	26	25	30	32	24	24	24	23
15-19	44	47	50	42	34	25	33	38	37	39	51	54
20-29	105	107	95	87	87	87	79	86	95	79	90	118
30-39	206	203	235	180	144	180	135	156	171	200	171	206
40-49	188	187	207	219	186	174	208	221	270	282	304	309
50-59	123	119	128	151	152	158	162	223	266	311	348	427
60-64	33	32	36	56	54	34	55	80	93	122	130	161
65-69	18	15	26	24	44	44	54	62	78	86	124	133
70-79	11	*	11	*	*	*	27	36	33	47	49	78
80+	*	*	.	*	.	*	*
Missing/unknown	*	.	*
0-19	107	104	117	102	92	71	87	101	92	97	107	105
20-44	409	402	431	379	330	351	319	351	399	418	388	467
45-64	246	246	270	314	293	282	320	415	496	576	655	754
65-74	26	22	36	28	47	50	80	96	105	124	165	199
75+	*	*	*	*	*	*	*	*	*	*	*	14
Unknown	*	.	*
Male	464	418	480	477	444	414	466	533	614	680	751	869
Female	328	358	376	347	321	343	342	432	485	544	574	670
Unknown
White	730	698	764	744	694	692	721	859	985	1,080	1,170	1,323
Black	47	58	70	49	57	46	62	77	81	105	107	130
Native American	*	*	*	*	*	*	*	*	*	*	11	*
Asian	*	15	13	20	*	14	20	21	26	33	33	33
Other/unknown	.	*	*	*	*	*	*	*	.	*	*	47
†Hispanic				46	37	47	44	68	60	67	70	86
†Non-Hispanic				778	728	710	764	897	1,039	1,157	1,255	1,453
Diabetes	210	192	216	232	192	188	183	193	261	273	278	331
Hypertension	55	55	56	46	61	49	56	71	102	125	117	157
Glomerulonephritis	155	154	198	176	151	158	156	191	211	260	245	291
Cystic kidney	84	66	99	105	77	98	98	125	153	185	229	217
Other urologic	48	50	65	50	63	50	39	72	69	86	87	93
Other cause	119	129	132	127	124	110	121	139	188	164	190	207
Unknown cause	66	50	39	37	51	32	49	55	60	79	65	65
Missing cause	55	80	51	51	46	72	106	119	55	52	114	178
All	792	776	856	824	765	757	808	965	1,099	1,224	1,325	1,539

The incident cohorts and associated modalities are determined at the time of ESRD initiation without applying the 60-day stable modality rule.

The age of the incident patient is determined as of the date of ESRD initiation.

†CMS began collecting Hispanic ethnicity data in April 1995; cells for years prior to 1996 have therefore been left blank.

* Values for cells with 10 or fewer patients are suppressed.

." Zero values in this cell.

Table D.6

Counts of prevalent ESRD patients, by mode of therapy: all dialysis*point prevalent patients on December 31, by demographic characteristics & primary diagnosis*

	1980	1981	1982	1983	1984	1985	1986	1987	1988	1989	1990	1991	1992
0-4	38	57	80	103	113	139	157	144	145	152	145	159	171
5-9	93	84	116	142	144	120	142	146	142	153	164	149	140
10-14	233	250	286	311	317	287	280	295	313	294	326	334	332
15-19	749	774	802	749	740	782	754	786	844	910	870	886	919
20-29	3,722	3,941	4,239	4,547	4,752	4,835	4,901	5,040	5,352	5,715	5,963	6,163	6,474
30-39	6,027	6,701	7,426	8,177	8,907	9,472	9,962	10,485	11,173	12,156	12,940	13,905	14,897
40-49	7,472	8,128	8,796	10,139	10,893	11,399	11,995	12,878	14,192	15,954	17,673	19,874	21,665
50-59	11,340	12,663	13,920	15,129	16,112	16,859	17,506	18,187	19,343	20,871	22,681	24,952	27,367
60-64	5,848	6,932	7,848	9,221	10,258	11,195	11,674	12,668	13,488	14,416	15,631	16,914	18,060
65-69	5,439	6,081	6,901	8,442	9,555	10,631	12,020	13,627	14,968	16,702	18,487	20,508	22,541
70-79	5,840	6,869	8,147	10,450	12,157	13,832	15,880	17,955	20,166	23,154	26,494	31,010	35,086
80+	853	1,078	1,373	1,992	2,394	2,964	3,641	4,462	5,284	6,347	7,693	9,168	10,842
Missing/unknown	*	*	*	*	12	13	*	*	*	*	*	*	.
0-19	1,113	1,165	1,284	1,305	1,314	1,328	1,333	1,371	1,444	1,509	1,505	1,528	1,562
20-44	13,232	14,369	15,732	17,476	18,715	19,609	20,596	21,614	23,281	25,489	27,594	29,669	31,699
45-64	21,177	23,996	26,497	29,737	32,207	34,151	35,442	37,644	40,267	43,623	47,294	52,139	56,764
65-74	9,293	10,524	12,159	15,011	17,054	19,032	21,583	24,262	26,816	30,196	34,094	38,813	43,235
75+	2,839	3,504	4,262	5,873	7,052	8,395	9,958	11,782	13,602	16,007	18,580	21,873	25,234
Unknown	*	*	*	*	12	13	*	*	*	*	*	*	.
Male	25,982	28,863	32,225	37,119	40,814	43,717	46,728	50,495	54,879	60,881	67,342	75,207	82,678
Female	21,674	24,698	27,713	32,291	35,540	38,811	42,192	46,185	50,537	55,949	61,731	68,819	75,816
Unknown	.	.	.	*
White	32,060	35,574	39,463	45,196	49,084	52,588	56,094	60,295	64,953	71,331	78,146	86,540	94,334
Black	14,417	16,621	18,977	22,434	25,249	27,615	30,188	33,383	36,988	41,559	46,384	52,159	57,983
Native American	222	302	382	532	637	740	858	1,000	1,162	1,339	1,506	1,713	1,991
Asian	274	380	566	782	993	1,264	1,507	1,780	2,127	2,508	2,946	3,537	4,105
Other/unknown	683	684	550	467	391	321	273	222	186	93	91	77	81
†Hispanic													
†Non-Hispanic													
Diabetes	4,640	6,479	8,716	11,038	13,736	16,287	19,038	22,266	25,858	30,954	36,609	43,374	50,032
Hypertension	8,145	10,466	13,302	16,185	18,808	21,253	23,974	26,918	30,243	34,622	39,031	44,168	49,405
Glomerulonephritis	9,549	10,904	12,441	14,373	15,864	17,069	18,097	19,194	20,302	21,708	22,994	24,513	25,858
Cystic kidney	3,098	3,510	3,979	4,541	4,860	5,155	5,323	5,514	5,641	5,782	5,965	6,259	6,476
Other urologic	1,135	1,432	1,838	2,229	2,588	2,968	3,194	3,505	3,768	3,981	4,085	4,377	4,608
Other cause	4,649	5,449	6,603	7,821	8,760	9,573	10,162	10,683	11,490	12,641	13,515	14,471	15,524
Unknown cause	5,018	5,264	5,126	5,414	5,554	5,413	5,492	5,742	5,777	5,475	5,500	5,771	5,884
Missing cause	11,422	10,057	7,933	7,810	6,184	4,810	3,640	2,858	2,337	1,667	1,374	1,093	707
All	47,656	53,561	59,938	69,411	76,354	82,528	88,920	96,680	105,416	116,830	129,073	144,026	158,494

The point prevalent cohorts and associated modalities are determined on December 31 of each calendar year without applying the 60-day stable modality rule. The age of the point prevalent patient is determined as of December 31 of each calendar year.

†CMS began collecting Hispanic ethnicity data in April 1995; cells for years prior to 1996 have therefore been left blank.

* Values for cells with 10 or fewer patients are suppressed.

." Zero values in this cell.

Table D.6

Counts of prevalent ESRD patients, by mode of therapy: all dialysis*point prevalent patients on December 31, by demographic characteristics & primary diagnosis*

	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004
0-4	169	164	170	167	144	139	133	153	153	168	159	177
5-9	147	171	160	148	159	138	139	164	150	170	144	152
10-14	345	367	354	343	335	340	303	336	336	395	354	360
15-19	948	889	911	940	930	958	907	860	904	921	938	916
20-29	6,638	6,798	6,860	7,009	7,050	7,233	7,246	7,259	7,262	7,148	7,092	7,022
30-39	15,664	16,422	16,738	17,031	17,177	17,306	17,813	18,134	18,381	18,544	18,794	18,616
40-49	23,489	25,299	27,331	29,197	30,331	31,832	33,130	34,408	35,469	36,445	37,288	37,849
50-59	29,717	32,399	34,398	36,971	39,672	42,823	46,138	49,310	52,598	55,350	57,962	60,378
60-64	19,156	20,380	21,577	22,773	23,972	25,331	26,491	27,735	29,038	30,798	32,385	33,433
65-69	24,267	25,981	27,186	28,821	30,003	31,166	32,436	33,447	34,574	35,523	36,502	37,263
70-79	38,551	43,130	46,719	51,025	55,342	58,814	61,867	64,360	65,939	68,124	69,293	70,003
80+	12,471	14,201	15,889	17,870	19,992	22,172	24,394	27,163	29,475	32,141	34,473	36,321
Missing/unknown	.	*	*	*	*
0-19	1,609	1,591	1,595	1,598	1,568	1,575	1,482	1,513	1,543	1,654	1,595	1,605
20-44	33,297	34,703	35,825	37,010	37,799	38,641	39,574	40,228	40,933	41,102	41,536	41,553
45-64	61,367	66,595	71,079	75,971	80,403	85,884	91,244	96,618	101,815	107,183	111,985	115,745
65-74	47,096	51,292	54,154	57,675	60,806	63,383	65,742	67,727	69,565	71,340	72,774	73,844
75+	28,193	32,020	35,640	40,041	44,531	48,769	52,955	57,243	60,423	64,448	67,494	69,743
Unknown	.	*	*	*	*
Male	89,751	97,398	103,559	111,360	118,608	126,096	133,636	140,847	147,357	154,471	160,184	164,934
Female	81,811	88,805	94,734	100,935	106,499	112,156	117,359	122,481	126,921	131,256	135,197	137,545
Unknown	*	*	*	*	*	13
White	100,880	108,992	114,820	122,334	129,158	135,769	143,124	150,379	156,631	162,500	167,526	171,565
Black	63,486	68,997	73,820	78,948	83,839	89,229	93,529	97,526	101,095	105,417	109,032	111,123
Native American	2,214	2,428	2,675	2,858	3,059	3,341	3,525	3,686	3,908	4,065	4,170	4,316
Asian	4,887	5,612	6,291	6,931	7,506	8,176	8,947	9,693	10,398	11,227	11,771	12,157
Other/unknown	95	174	687	1,224	1,545	1,737	1,872	2,045	2,247	2,519	2,886	3,331
†Hispanic				21,467	23,910	26,094	28,632	31,989	33,971	36,096	38,316	40,069
†Non-Hispanic				190,828	201,197	212,158	222,365	231,340	240,308	249,632	257,069	262,423
Diabetes	55,960	63,028	70,252	78,390	86,219	93,628	100,465	107,680	113,947	120,132	125,509	129,620
Hypertension	53,838	58,515	60,235	62,879	65,518	68,567	71,774	74,665	77,393	80,655	83,915	85,732
Glomerulonephritis	27,306	28,923	30,121	31,486	32,365	33,211	33,977	34,373	34,804	35,104	35,123	35,036
Cystic kidney	6,718	7,034	7,194	7,239	7,252	7,423	7,630	7,821	7,923	8,031	8,068	8,089
Other urologic	4,593	4,770	5,275	5,813	6,239	6,583	7,113	7,430	7,686	7,954	8,017	8,020
Other cause	16,384	17,186	18,188	18,936	19,606	20,527	21,263	22,109	22,784	23,617	24,120	24,626
Unknown cause	6,117	6,396	6,716	7,273	7,673	8,112	8,571	9,074	9,565	9,946	10,187	10,194
Missing cause	646	351	312	279	235	201	204	177	177	289	446	1,175
All	171,562	186,203	198,293	212,295	225,107	238,252	250,997	263,329	274,279	285,728	295,385	302,492

The point prevalent cohorts and associated modalities are determined on December 31 of each calendar year without applying the 60-day stable modality rule.

The age of the point prevalent patient is determined as of December 31 of each calendar year.

†CMS began collecting Hispanic ethnicity data in April 1995; cells for years prior to 1996 have therefore been left blank.

* Values for cells with 10 or fewer patients are suppressed.

." Zero values in this cell.

Table D.7

Counts of prevalent ESRD patients, by mode of therapy: hemodialysis*point prevalent patients on December 31, by demographic characteristics & primary diagnosis*

	1980	1981	1982	1983	1984	1985	1986	1987	1988	1989	1990	1991	1992
0-4	21	18	20	28	26	35	39	37	36	41	36	37	26
5-9	67	54	63	59	48	32	42	48	47	43	47	50	46
10-14	194	200	199	179	158	142	118	132	135	126	143	131	143
15-19	706	717	679	606	572	542	527	541	578	605	560	554	567
20-29	3,522	3,669	3,843	3,953	4,005	4,015	3,993	4,048	4,241	4,545	4,682	4,773	4,978
30-39	5,619	6,136	6,583	7,009	7,398	7,684	8,092	8,473	8,965	9,702	10,124	10,772	11,507
40-49	6,983	7,503	7,914	8,830	9,292	9,499	9,919	10,626	11,642	13,034	14,286	15,839	17,231
50-59	10,575	11,631	12,635	13,413	13,961	14,386	14,885	15,428	16,457	17,664	18,974	20,783	22,679
60-64	5,545	6,405	7,154	8,235	9,012	9,649	10,036	10,866	11,556	12,364	13,283	14,448	15,257
65-69	5,098	5,656	6,286	7,694	8,500	9,264	10,427	11,867	13,064	14,584	16,087	17,680	19,460
70-79	5,429	6,354	7,380	9,573	11,005	12,319	14,130	15,977	17,991	20,759	23,641	27,529	31,088
80+	774	969	1,225	1,822	2,175	2,688	3,313	4,072	4,837	5,821	7,072	8,412	9,999
Missing/unknown	*	*	*	*	12	12	*	*	*	*	*	*	.
0-19	988	989	961	872	804	751	726	758	796	815	786	772	782
20-44	12,395	13,224	14,075	15,092	15,679	16,094	16,772	17,464	18,692	20,442	21,787	23,125	24,650
45-64	19,849	22,120	24,054	26,348	27,989	29,139	30,153	31,977	34,169	36,867	39,562	43,490	47,002
65-74	8,691	9,773	11,043	13,696	15,246	16,700	18,880	21,272	23,527	26,594	29,926	33,807	37,648
75+	2,610	3,206	3,848	5,393	6,434	7,571	8,990	10,644	12,365	14,570	16,874	19,814	22,899
Unknown	*	*	*	*	12	12	*	*	*	*	*	*	.
Male	24,353	26,651	29,073	32,750	35,304	37,130	39,631	42,750	46,493	51,584	56,720	62,918	69,166
Female	20,182	22,664	24,912	28,659	30,860	33,137	35,897	39,371	43,062	47,710	52,221	58,094	63,815
Unknown	.	.	.	*
White	29,670	32,288	34,933	39,042	41,337	43,228	46,004	49,415	53,177	58,478	63,466	69,860	75,972
Black	13,751	15,726	17,651	20,758	23,047	24,989	27,209	30,107	33,333	37,391	41,525	46,543	51,734
Native American	209	283	348	472	533	631	722	825	988	1,145	1,277	1,450	1,680
Asian	266	359	521	707	883	1,129	1,351	1,579	1,898	2,205	2,598	3,087	3,519
Other/unknown	639	659	532	431	364	290	242	195	159	75	75	72	76
†Hispanic													
†Non-Hispanic													
Diabetes	4,101	5,651	7,424	9,359	11,485	13,515	15,901	18,687	21,819	26,211	30,864	36,457	42,038
Hypertension	7,665	9,719	12,102	14,671	16,818	18,749	21,077	23,740	26,676	30,553	34,145	38,421	43,046
Glomerulonephritis	8,890	9,967	11,101	12,469	13,382	14,110	14,911	15,731	16,526	17,649	18,562	19,636	20,609
Cystic kidney	2,933	3,244	3,597	4,024	4,221	4,384	4,464	4,623	4,738	4,832	4,877	5,072	5,158
Other urologic	1,064	1,313	1,640	1,988	2,273	2,582	2,773	3,076	3,302	3,484	3,524	3,784	3,946
Other cause	4,345	4,997	5,900	6,798	7,453	7,966	8,468	8,848	9,530	10,424	11,102	11,771	12,545
Unknown cause	4,676	4,830	4,630	4,824	4,873	4,688	4,733	4,962	4,985	4,720	4,707	4,956	5,057
Missing cause	10,861	9,594	7,591	7,277	5,659	4,273	3,201	2,454	1,979	1,421	1,160	915	582
All	44,535	49,315	53,985	61,410	66,164	70,267	75,528	82,121	89,555	99,294	108,941	121,012	132,981

The point prevalent cohorts and associated modalities are determined on December 31 of each calendar year without applying the 60-day stable modality. The age of the point prevalent patient is determined as of December 31 of each calendar year.

†CMS began collecting Hispanic ethnicity data in April 1995; cells for years prior to 1996 have therefore been left blank.

* Values for cells with 10 or fewer patients are suppressed.

." Zero values in this cell.

Table D.7

Counts of prevalent ESRD patients, by mode of therapy: hemodialysis*point prevalent patients on December 31, by demographic characteristics & primary diagnosis*

	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004
0-4	35	37	34	34	25	26	31	29	38	32	28	38
5-9	51	52	52	59	63	62	55	71	54	58	43	62
10-14	145	168	163	151	143	158	148	165	170	186	168	165
15-19	587	549	594	596	611	653	628	587	614	658	669	646
20-29	5,043	5,185	5,280	5,500	5,612	5,898	5,990	6,017	6,034	6,003	5,959	5,924
30-39	12,018	12,576	13,042	13,553	14,022	14,404	14,924	15,394	15,690	15,906	16,216	16,161
40-49	18,661	20,082	21,875	23,933	25,367	27,271	28,767	30,076	31,272	32,334	33,160	33,760
50-59	24,672	26,773	28,582	31,126	34,131	37,555	40,798	44,082	47,345	49,989	52,494	54,963
60-64	16,341	17,338	18,391	19,755	21,133	22,784	24,034	25,332	26,683	28,365	29,774	30,871
65-69	20,864	22,401	23,551	25,261	26,841	28,326	29,648	30,747	31,978	32,932	33,928	34,700
70-79	34,142	38,282	41,774	45,998	50,630	54,453	57,588	60,280	62,142	64,215	65,344	66,091
80+	11,514	13,100	14,730	16,635	18,800	21,119	23,229	26,037	28,369	30,962	33,237	35,078
Missing/unknown	.	*	*
0-19	818	806	843	840	842	899	862	852	876	934	908	911
20-44	25,729	26,784	27,996	29,583	30,870	32,338	33,415	34,295	35,114	35,539	35,983	36,160
45-64	51,006	55,170	59,174	64,284	69,395	75,574	81,098	86,606	91,910	97,058	101,620	105,519
65-74	40,856	44,627	47,445	51,041	54,833	57,913	60,402	62,653	64,778	66,516	67,876	69,027
75+	25,664	29,156	32,610	36,853	41,438	45,985	50,063	54,411	57,711	61,593	64,633	66,842
Unknown	.	*	*
Male	75,187	81,767	87,740	95,952	104,364	112,998	120,698	128,184	135,097	141,982	147,572	152,430
Female	68,886	74,777	80,328	86,649	93,014	99,711	105,140	110,632	115,291	119,657	123,444	126,017
Unknown	*	*	*	*	*	13
White	81,450	88,289	93,879	101,828	110,136	118,443	126,065	133,883	140,651	146,446	151,358	155,634
Black	56,516	61,343	66,092	71,431	76,777	82,596	87,025	91,159	94,885	99,175	102,693	104,888
Native American	1,895	2,072	2,306	2,509	2,708	3,017	3,245	3,380	3,586	3,734	3,843	4,015
Asian	4,126	4,685	5,207	5,813	6,440	7,119	7,840	8,552	9,247	10,014	10,500	10,899
Other/unknown	86	155	584	1,020	1,317	1,534	1,665	1,843	2,020	2,271	2,626	3,024
†Hispanic				18,718	21,164	23,467	25,901	29,189	31,240	33,208	35,427	37,247
†Non-Hispanic				163,883	176,214	189,242	199,939	209,628	219,149	228,432	235,593	241,213
Diabetes	47,396	53,486	60,215	68,222	76,657	84,827	91,715	99,082	105,598	111,667	116,959	121,316
Hypertension	46,837	50,899	52,775	55,831	59,071	62,687	65,882	68,911	71,963	75,048	78,147	79,975
Glomerulonephritis	21,665	22,891	23,931	25,376	26,593	27,872	28,944	29,491	30,082	30,449	30,513	30,454
Cystic kidney	5,318	5,581	5,720	5,824	5,990	6,288	6,502	6,721	6,829	7,024	6,973	7,012
Other urologic	3,903	4,050	4,516	5,025	5,480	5,859	6,368	6,691	6,942	7,185	7,247	7,267
Other cause	13,192	13,911	14,900	15,807	16,679	17,807	18,561	19,551	20,125	20,916	21,452	22,015
Unknown cause	5,220	5,430	5,747	6,279	6,706	7,192	7,692	8,222	8,691	9,100	9,341	9,358
Missing cause	542	296	264	237	202	177	176	148	159	251	388	1,063
All	144,073	156,544	168,068	182,601	197,378	212,709	225,840	238,817	250,389	261,640	271,020	278,460

The point prevalent cohorts and associated modalities are determined on December 31 of each calendar year without applying the 60-day stable modality rule.

The age of the point prevalent patient is determined as of December 31 of each calendar year.

†CMS began collecting Hispanic ethnicity data in April 1995; cells for years prior to 1996 have therefore been left blank.

* Values for cells with 10 or fewer patients are suppressed.

." Zero values in this cell.

Table D.8

Counts of prevalent ESRD patients, by mode of therapy: peritoneal dialysis*point prevalent patients on December 31, by demographic characteristics & primary diagnosis*

	1980	1981	1982	1983	1984	1985	1986	1987	1988	1989	1990	1991	1992
0-4	11	29	48	63	72	95	110	90	92	92	89	111	129
5-9	*	15	42	67	79	69	86	85	78	96	104	89	83
10-14	17	30	59	113	138	129	144	147	154	142	162	181	175
15-19	*	36	86	116	146	216	199	220	228	271	275	304	326
20-29	60	184	285	505	649	727	809	886	992	1,033	1,175	1,295	1,399
30-39	127	357	588	974	1,292	1,596	1,653	1,814	2,004	2,212	2,595	2,908	3,189
40-49	129	370	638	1,122	1,377	1,689	1,867	2,014	2,286	2,661	3,122	3,745	4,138
50-59	203	616	923	1,442	1,878	2,200	2,385	2,516	2,614	2,943	3,408	3,842	4,371
60-64	65	318	509	823	1,078	1,382	1,503	1,662	1,758	1,906	2,167	2,274	2,611
65-69	84	246	424	593	870	1,175	1,414	1,549	1,687	1,895	2,159	2,585	2,861
70-79	89	241	510	625	906	1,249	1,462	1,693	1,850	2,104	2,485	3,108	3,596
80+	12	47	91	108	144	199	245	304	347	427	507	620	720
Missing/unknown	*	.	*
0-19	45	110	235	359	435	509	539	542	552	601	630	685	713
20-44	247	722	1,167	2,008	2,602	3,115	3,402	3,721	4,138	4,541	5,334	6,081	6,614
45-64	337	1,123	1,776	2,858	3,672	4,479	4,815	5,171	5,516	6,214	7,133	7,983	9,094
65-74	141	407	773	999	1,472	2,004	2,352	2,607	2,884	3,222	3,727	4,546	5,125
75+	44	127	252	327	448	619	769	939	1,000	1,204	1,424	1,767	2,052
Unknown	*	.	*
Male	435	1,294	2,210	3,580	4,636	5,739	6,255	6,846	7,411	8,328	9,595	11,205	12,456
Female	379	1,195	1,993	2,971	3,993	4,988	5,622	6,135	6,679	7,454	8,653	9,857	11,142
Unknown
White	676	1,976	3,254	5,087	6,608	8,248	9,016	9,767	10,539	11,609	13,325	15,194	16,918
Black	121	469	871	1,313	1,796	2,234	2,564	2,847	3,168	3,701	4,382	5,204	5,824
Native American	*	15	32	56	100	101	133	167	163	183	217	251	302
Asian	*	17	37	65	104	122	143	184	208	280	318	411	552
Other/unknown	*	12	*	30	21	22	21	16	12	*	*	*	*
†Hispanic													
†Non-Hispanic													
Diabetes	187	526	983	1,366	1,929	2,446	2,828	3,222	3,618	4,325	5,204	6,269	7,401
Hypertension	151	467	894	1,262	1,671	2,162	2,554	2,822	3,155	3,647	4,458	5,278	5,917
Glomerulonephritis	205	576	915	1,572	2,131	2,630	2,848	3,121	3,402	3,684	4,065	4,533	4,931
Cystic kidney	58	170	292	449	571	705	789	810	816	859	989	1,113	1,223
Other urologic	23	82	137	192	271	342	365	377	406	463	507	540	594
Other cause	98	289	522	859	1,126	1,429	1,520	1,652	1,775	1,983	2,214	2,497	2,715
Unknown cause	67	206	294	428	513	562	600	635	626	617	658	711	738
Missing cause	25	173	166	423	417	451	373	342	292	204	153	121	79
All	814	2,489	4,203	6,551	8,629	10,727	11,877	12,981	14,090	15,782	18,248	21,062	23,598

The point prevalent cohorts and associated modalities are determined on December 31 of each calendar year without applying the 60-day stay rule. The age of the point prevalent patient is determined as of December 31 of each calendar year.

†CMS began collecting Hispanic ethnicity data in April 1995; cells for years prior to 1996 have therefore been left blank.

* Values for cells with 10 or fewer patients are suppressed.

." Zero values in this cell.

Table D.8

Counts of prevalent ESRD patients, by mode of therapy: peritoneal dialysis*point prevalent patients on December 31, by demographic characteristics & primary diagnosis*

	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004
0-4	115	122	132	128	115	111	101	122	113	135	129	135
5-9	83	108	100	88	92	69	80	92	95	110	100	89
10-14	180	185	178	181	183	174	151	164	159	202	180	189
15-19	330	313	303	322	300	274	258	252	260	242	252	257
20-29	1,510	1,538	1,508	1,464	1,387	1,274	1,180	1,166	1,172	1,104	1,095	1,062
30-39	3,453	3,632	3,565	3,349	3,041	2,784	2,748	2,616	2,577	2,536	2,463	2,368
40-49	4,570	4,929	5,223	5,090	4,798	4,401	4,192	4,146	4,036	3,966	3,999	3,967
50-59	4,800	5,359	5,631	5,672	5,420	5,110	5,130	5,038	5,123	5,214	5,322	5,267
60-64	2,677	2,870	3,054	2,928	2,777	2,487	2,373	2,333	2,299	2,366	2,544	2,497
65-69	3,187	3,331	3,422	3,291	3,000	2,715	2,629	2,544	2,550	2,543	2,513	2,504
70-79	4,077	4,469	4,600	4,495	4,340	4,097	3,878	3,730	3,730	3,810	3,846	3,851
80+	835	969	1,030	1,025	1,039	959	973	970	1,066	1,123	1,187	1,217
Missing/unknown	*	*	*
0-19	708	728	713	719	690	628	590	630	627	689	661	670
20-44	7,166	7,492	7,512	7,173	6,674	6,045	5,852	5,649	5,562	5,354	5,344	5,208
45-64	9,844	10,836	11,469	11,330	10,749	10,011	9,771	9,650	9,645	9,832	10,079	9,953
65-74	5,815	6,200	6,294	6,069	5,608	5,209	4,970	4,733	4,708	4,727	4,778	4,720
75+	2,284	2,569	2,758	2,742	2,771	2,562	2,510	2,511	2,638	2,749	2,768	2,852
Unknown	*	*	*
Male	13,647	14,616	15,016	14,530	13,630	12,534	12,134	11,958	11,869	12,073	12,191	12,134
Female	12,170	13,209	13,730	13,503	12,862	11,921	11,559	11,215	11,311	11,279	11,440	11,270
Unknown
White	18,174	19,345	19,839	19,203	18,061	16,514	15,956	15,494	15,497	15,544	15,636	15,492
Black	6,599	7,221	7,395	7,208	6,821	6,399	6,208	6,085	6,014	6,055	6,168	6,077
Native American	309	346	362	341	343	320	276	294	316	329	322	298
Asian	729	898	1,052	1,088	1,044	1,022	1,053	1,105	1,128	1,181	1,252	1,240
Other/unknown	*	15	98	193	223	200	200	195	225	243	253	297
†Hispanic				2,653	2,655	2,534	2,615	2,710	2,658	2,807	2,819	2,741
†Non-Hispanic				25,380	23,837	21,921	21,078	20,463	20,522	20,545	20,812	20,663
Diabetes	8,038	8,940	9,540	9,592	9,163	8,461	8,246	8,147	8,181	8,275	8,330	8,129
Hypertension	6,602	7,168	7,112	6,623	6,147	5,635	5,542	5,410	5,288	5,442	5,616	5,655
Glomerulonephritis	5,351	5,710	5,937	5,860	5,549	5,097	4,752	4,638	4,520	4,462	4,456	4,418
Cystic kidney	1,316	1,365	1,411	1,354	1,221	1,104	1,086	1,041	1,061	976	1,055	1,035
Other urologic	647	666	704	717	701	683	685	690	712	736	737	725
Other cause	2,985	3,062	3,118	2,950	2,787	2,590	2,555	2,435	2,557	2,619	2,581	2,537
Unknown cause	811	875	895	915	907	872	811	796	850	824	817	823
Missing cause	67	39	29	22	17	13	16	16	11	18	39	82
All	25,817	27,825	28,746	28,033	26,492	24,455	23,693	23,173	23,180	23,352	23,631	23,404

The point prevalent cohorts and associated modalities are determined on December 31 of each calendar year without applying the 60-day stable modality rule.

The age of the point prevalent patient is determined as of December 31 of each calendar year.

†CMS began collecting Hispanic ethnicity data in April 1995; cells for years prior to 1996 have therefore been left blank.

* Values for cells with 10 or fewer patients are suppressed.

." Zero values in this cell.

Table D.9

Counts of prevalent ESRD patients, by mode of therapy: transplant*point prevalent patients on December 31, by demographic characteristics & primary diagnosis*

	1980	1981	1982	1983	1984	1985	1986	1987	1988	1989	1990	1991	1992
0-4	23	38	66	84	94	112	121	148	148	140	127	133	137
5-9	99	113	129	137	173	222	245	259	265	291	311	341	342
10-14	248	255	283	308	352	399	427	447	452	465	472	480	511
15-19	548	582	604	659	714	779	863	918	868	859	879	882	867
20-29	2,098	2,437	2,755	3,096	3,416	3,749	4,134	4,337	4,312	4,368	4,606	4,848	4,943
30-39	2,266	2,924	3,517	4,282	5,149	6,104	7,160	7,882	8,309	8,716	9,454	10,151	10,774
40-49	1,636	1,982	2,457	2,974	3,603	4,510	5,563	6,619	7,496	8,323	9,215	10,408	11,552
50-59	982	1,257	1,573	1,934	2,475	3,008	3,864	4,702	5,330	5,945	6,734	7,468	8,431
60-64	137	183	242	330	475	668	933	1,262	1,653	2,013	2,354	2,733	3,052
65-69	56	70	91	118	164	252	370	572	802	1,045	1,343	1,687	2,115
70-79	15	21	26	35	50	75	112	174	239	317	454	657	869
80+	*	*	*	*	*	11	14	13	14	14	22	27	26
Missing/unknown
0-19	918	988	1,082	1,188	1,333	1,512	1,656	1,772	1,733	1,755	1,789	1,836	1,857
20-44	5,239	6,441	7,613	8,986	10,515	12,260	14,318	15,812	16,643	17,600	19,128	20,656	21,834
45-64	1,880	2,342	2,931	3,630	4,603	5,779	7,336	8,990	10,457	11,765	13,235	14,952	16,918
65-74	68	84	110	146	206	313	461	708	992	1,296	1,711	2,233	2,854
75+	*	*	*	*	12	25	35	51	63	80	108	138	156
Unknown
Male	4,939	6,109	7,375	8,772	10,458	12,524	14,976	17,137	18,600	20,082	22,114	24,280	26,494
Female	3,171	3,754	4,371	5,186	6,209	7,363	8,828	10,194	11,286	12,413	13,856	15,534	17,124
Unknown	*	*	*	*	*	*	*	*	*
White	6,570	7,984	9,433	11,125	13,251	15,722	18,777	21,542	23,445	25,518	28,235	31,176	34,046
Black	1,301	1,607	2,005	2,466	2,985	3,649	4,367	4,977	5,473	5,867	6,435	7,148	7,864
Native American	50	67	83	96	119	149	180	202	242	292	326	377	408
Asian	67	93	135	187	253	313	429	550	675	773	933	1,071	1,261
Other/unknown	122	112	90	84	61	56	53	62	53	46	42	43	40
†Hispanic													
†Non-Hispanic													
Diabetes	754	1,093	1,512	2,032	2,625	3,408	4,426	5,284	6,018	6,845	7,832	8,891	9,812
Hypertension	849	1,099	1,409	1,819	2,206	2,742	3,345	3,923	4,424	4,891	5,513	6,125	6,858
Glomerulonephritis	2,687	3,247	3,919	4,584	5,516	6,512	7,581	8,607	9,286	9,945	10,936	12,062	13,134
Cystic kidney	367	485	622	764	1,008	1,237	1,587	1,967	2,299	2,562	2,898	3,227	3,646
Other urologic	302	399	465	582	671	803	939	1,053	1,112	1,188	1,299	1,468	1,626
Other cause	1,278	1,545	1,799	2,168	2,605	3,104	3,689	4,120	4,312	4,572	4,936	5,413	5,859
Unknown cause	681	812	887	949	1,017	1,101	1,241	1,377	1,447	1,553	1,624	1,697	1,760
Missing cause	1,192	1,183	1,133	1,060	1,021	982	998	1,002	990	940	933	932	924
All	8,110	9,863	11,746	13,958	16,669	19,889	23,806	27,333	29,888	32,496	35,971	39,815	43,619

The point prevalent cohorts and associated modalities are determined on December 31 of each calendar year without applying the 60-day stable mc. The age of the point prevalent patient is determined as of December 31 of each calendar year.

†CMS began collecting Hispanic ethnicity data in April 1995; cells for years prior to 1996 have therefore been left blank.

* Values for cells with 10 or fewer patients are suppressed.

." Zero values in this cell.

Table D.9

Counts of prevalent ESRD patients, by mode of therapy: transplant*point prevalent patients on December 31, by demographic characteristics & primary diagnosis*

	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004
0-4	151	144	144	151	149	143	150	153	160	159	158	166
5-9	344	327	351	365	392	405	409	356	364	318	321	296
10-14	542	524	563	619	613	654	684	678	636	652	573	554
15-19	909	968	1,074	1,077	1,087	1,076	1,151	1,133	1,129	1,081	1,025	1,084
20-29	5,033	4,874	5,029	5,121	5,148	5,159	5,194	5,114	5,033	4,907	4,590	4,678
30-39	11,440	11,645	12,198	12,664	12,777	13,157	13,324	13,472	13,470	13,521	13,107	13,149
40-49	12,788	13,598	14,851	15,971	16,781	17,670	18,632	19,544	20,260	20,947	21,008	21,783
50-59	9,464	10,167	11,253	12,551	14,205	15,732	17,396	18,716	20,254	21,805	22,605	24,559
60-64	3,402	3,616	4,089	4,683	5,194	5,751	6,423	6,925	7,570	8,427	8,973	9,781
65-69	2,597	3,035	3,503	4,036	4,528	5,024	5,627	6,249	6,949	7,731	8,632	9,498
70-79	1,100	1,400	1,742	2,115	2,600	3,130	3,640	4,344	5,023	5,946	6,811	7,793
80+	29	41	48	73	84	116	160	210	266	336	417	534
Missing/unknown	*	.	*	*	*	*	*	*	*	*	.	.
0-19	1,946	1,963	2,132	2,212	2,241	2,278	2,394	2,320	2,289	2,210	2,077	2,100
20-44	23,153	23,448	24,687	25,660	26,227	27,024	27,723	28,053	28,193	28,364	27,559	27,945
45-64	18,974	20,452	22,733	25,330	27,878	30,445	33,246	35,718	38,394	41,243	42,724	46,005
65-74	3,522	4,210	4,920	5,734	6,594	7,474	8,430	9,579	10,774	12,184	13,656	15,209
75+	204	266	373	490	618	796	997	1,224	1,464	1,829	2,204	2,616
Unknown	*	.	*	*	*	*	*	*	*	*	.	.
Male	28,962	30,592	33,361	35,989	38,433	40,935	43,660	46,048	48,522	51,263	52,917	56,203
Female	18,837	19,747	21,485	23,438	25,126	27,083	29,131	30,846	32,592	34,567	35,303	37,672
Unknown	*	*	*	*	.	.
White	37,196	38,838	42,133	45,420	48,357	51,552	54,933	57,881	60,805	64,287	65,794	69,566
Black	8,686	9,384	10,362	11,374	12,345	13,313	14,335	15,195	16,189	17,141	17,794	19,055
Native American	452	506	550	610	648	692	780	820	856	893	897	969
Asian	1,426	1,570	1,758	1,974	2,156	2,398	2,671	2,912	3,173	3,403	3,565	3,916
Other/unknown	40	41	43	49	53	63	72	87	92	107	170	369
†Hispanic				5,352	5,904	6,651	7,392	8,061	8,758	9,551	9,954	10,821
†Non-Hispanic				54,075	57,655	61,367	65,399	68,834	72,357	76,280	78,266	83,054
Diabetes	11,127	12,146	13,423	14,747	15,933	17,158	18,416	19,598	20,759	22,197	23,140	24,666
Hypertension	7,603	8,237	9,090	9,946	10,637	11,325	12,057	12,653	13,398	14,189	14,479	15,444
Glomerulonephritis	14,094	14,545	15,703	16,855	17,855	18,981	20,183	21,217	22,111	23,154	23,532	24,801
Cystic kidney	3,982	4,163	4,566	4,979	5,403	5,823	6,254	6,626	7,035	7,535	7,922	8,452
Other urologic	1,815	1,847	1,989	2,135	2,286	2,449	2,621	2,733	2,910	3,073	3,170	3,361
Other cause	6,419	6,647	7,217	7,798	8,291	8,888	9,575	10,072	10,621	11,099	11,151	11,934
Unknown cause	1,831	1,845	1,934	2,043	2,209	2,398	2,616	2,834	3,082	3,352	3,505	3,703
Missing cause	929	909	924	924	945	996	1,069	1,162	1,199	1,232	1,321	1,514
All	47,800	50,339	54,846	59,427	63,559	68,018	72,791	76,895	81,115	85,831	88,220	93,875

The point prevalent cohorts and associated modalities are determined on December 31 of each calendar year without applying the 60-day stable modality rule.

The age of the point prevalent patient is determined as of December 31 of each calendar year.

†CMS began collecting Hispanic ethnicity data in April 1995; cells for years prior to 1996 have therefore been left blank.

* Values for cells with 10 or fewer patients are suppressed.

." Zero values in this cell.

Table D.15

Incident ESRD patients, 2004: by modality & state/territory*incident ESRD patients*

	Center hemo	Center self hemo	Home hemo	CAPD	CCPD	Other PD	Uncertain dialysis	Transplant	All
Alabama	1,567	.	.	92	15	*	.	18	1,694
Alaska	82	.	.	22	*	.	.	*	109
Arizona	1,569	.	*	80	11	.	*	26	1,693
Arkansas	795	*	.	62	24	*	*	*	896
California	8,740	*	*	438	100	15	52	116	9,467
Colorado	723	.	.	45	*	.	*	17	792
Connecticut	735	.	*	72	34	*	.	15	859
Delaware	264	.	.	14	*	.	.	*	285
District of Columbia	318	.	*	*	*	.	.	*	332
Florida	5,271	.	65	183	114	16	*	68	5,727
Georgia	2,618	.	.	180	42	*	*	54	2,899
Hawaii	494	.	.	15	.	.	*	*	520
Idaho	193	*	.	50	*	*	.	13	261
Illinois	3,332	.	44	168	72	20	*	85	3,728
Indiana	1,607	.	*	125	19	*	*	54	1,814
Iowa	607	*	*	22	17	.	*	42	695
Kansas	545	.	*	65	*	.	*	11	633
Kentucky	1,138	.	*	77	17	21	*	16	1,277
Louisiana	1,907	.	.	63	13	*	*	20	2,006
Maine	278	.	.	*	*	.	.	15	312
Maryland	1,770	.	*	73	52	*	*	29	1,938
Massachusetts	1,348	.	.	56	29	.	*	27	1,462
Michigan	2,901	.	*	187	26	*	*	71	3,198
Minnesota	952	.	*	36	18	*	*	84	1,095
Mississippi	1,066	.	*	91	36	*	.	*	1,207
Missouri	1,578	.	*	98	33	*	*	23	1,745
Montana	146	.	*	17	*	*	.	*	175
Nebraska	432	.	*	13	16	.	.	12	474
Nevada	597	.	.	29	*	.	*	*	640
New Hampshire	226	.	.	16	*	.	*	*	260
New Jersey	2,713	.	*	62	59	*	*	50	2,901
New Mexico	505	.	.	29	15	*	.	*	556
New York	5,380	*	*	137	102	55	*	71	5,764
North Carolina	2,722	.	*	111	110	*	*	33	2,987
North Dakota	131	.	.	18	*	*	*	15	170
Ohio	3,519	.	12	219	47	*	*	71	3,876
Oklahoma	999	.	*	55	34	*	*	*	1,110
Oregon	673	.	*	56	12	.	*	16	764
Pennsylvania	3,786	*	*	90	83	*	*	76	4,049
Rhode Island	253	.	.	*	.	.	.	*	268
South Carolina	1,635	.	.	77	58	.	*	17	1,793
South Dakota	175	.	.	*	12	.	*	14	209
Tennessee	1,709	.	*	44	86	.	*	25	1,867
Texas	6,576	.	15	209	168	40	20	70	7,098
Utah	269	*	.	18	*	.	*	24	327
Vermont	96	.	.	*	*	.	.	*	113
Virginia	2,213	.	.	110	33	15	*	34	2,415
Washington	1,103	14	*	61	14	*	*	32	1,229
West Virginia	623	.	.	29	47	*	.	14	723
Wisconsin	1,401	.	.	70	37	*	*	83	1,599
Wyoming	76	.	.	*	*	*	.	*	84
American Samoa	20	20
Guam	46	.	.	*	52
Puerto Rico	1,016	.	*	15	91	*	.	.	1,124
Saipan	19	.	.	*	20
Virgin Islands	44	.	.	*	*	.	.	.	46
Foreign	*	.	.	*	11
Unknown	*	.	.	*	*
All	81,511	25	198	3,847	1,766	268	216	1,539	89,370

The incident cohorts and associated modalities are determined at the time of ESRD initiation without applying the 60-day stable modality rule.

* Values for cells with 10 or fewer patients are suppressed.

." Zero values in this cell.

Table D.16

Prevalent ESRD patients, 2004: by modality & state/territory*point prevalent ESRD patients on December 31*

	Center hemo	Center self hemo	Home hemo	CAPD	CCPD	Other PD	Uncertain dialysis	Transplant	All
Alabama	5,692	*	24	219	298	.	*	1,619	7,857
Alaska	225	.	*	34	47	.	*	145	455
Arizona	4,932	*	16	154	201	.	*	1,665	6,979
Arkansas	2,264	13	20	171	184	.	*	903	3,562
California	31,321	28	84	1,235	1,530	*	95	10,015	44,310
Colorado	2,553	*	*	84	134	.	11	1,170	3,957
Connecticut	2,591	.	*	132	319	.	*	907	3,962
Delaware	941	.	*	16	41	.	*	369	1,369
District of Columbia	1,433	.	*	24	26	.	*	197	1,685
Florida	16,544	*	169	353	858	*	37	5,267	23,230
Georgia	10,641	*	36	427	537	*	13	2,388	14,044
Hawaii	1,848	.	*	75	42	.	*	360	2,329
Idaho	585	*	*	93	64	.	*	402	1,157
Illinois	11,247	*	278	398	638	*	23	4,400	16,996
Indiana	5,348	.	37	320	265	.	11	2,021	8,002
Iowa	1,970	.	17	50	93	.	*	1,130	3,265
Kansas	1,881	*	17	134	105	.	*	706	2,852
Kentucky	3,409	*	26	128	191	*	17	1,254	5,027
Louisiana	6,389	*	22	116	207	.	12	1,371	8,119
Maine	869	.	*	26	52	.	*	415	1,372
Maryland	6,082	.	21	232	317	*	17	2,306	8,976
Massachusetts	4,139	.	31	119	188	.	14	1,952	6,443
Michigan	9,606	*	45	494	369	.	26	3,634	14,175
Minnesota	2,806	*	13	70	121	.	*	2,093	5,109
Mississippi	4,338	*	45	166	275	*	*	927	5,758
Missouri	5,179	.	60	246	323	.	*	1,800	7,614
Montana	518	.	*	43	37	.	.	336	937
Nebraska	1,240	.	16	43	89	.	*	603	1,992
Nevada	1,602	.	*	72	93	.	*	588	2,365
New Hampshire	638	.	*	25	32	.	*	349	1,050
New Jersey	8,649	*	25	176	335	*	17	2,819	12,028
New Mexico	1,973	.	*	93	123	*	*	580	2,781
New York	18,671	79	112	478	575	*	35	5,438	25,392
North Carolina	10,413	*	36	274	596	*	11	2,602	13,935
North Dakota	365	.	*	39	14	.	*	300	722
Ohio	11,199	*	116	488	528	*	27	3,892	16,254
Oklahoma	3,077	*	18	136	230	.	*	1,002	4,470
Oregon	2,078	.	14	109	154	.	*	1,080	3,441
Pennsylvania	11,511	23	47	273	393	*	46	4,829	17,124
Rhode Island	766	.	*	13	14	*	*	386	1,186
South Carolina	5,956	*	21	161	333	*	*	1,423	7,901
South Dakota	566	.	.	24	45	.	*	320	958
Tennessee	5,715	*	37	157	367	.	*	2,001	8,287
Texas	24,614	.	134	577	1,236	*	36	6,510	33,113
Utah	873	*	11	76	24	.	*	821	1,813
Vermont	355	.	*	*	15	.	*	170	552
Virginia	7,781	*	68	260	386	*	11	2,461	10,969
Washington	3,664	*	101	145	171	*	*	1,779	5,877
West Virginia	1,568	.	*	66	170	*	*	697	2,510
Wisconsin	3,929	*	21	146	188	.	*	2,482	6,777
Wyoming	233	.	*	15	12	.	*	160	426
American Samoa	83	83
Guam	245	.	.	18	*	.	.	*	271
Puerto Rico	3,036	.	*	67	299	*	*	760	4,175
Saipan	88	.	.	*	.	.	.	*	104
Virgin Islands	148	.	.	*	*	.	.	16	172
Foreign	51	.	*	*	*	.	.	37	92
Unknown	*	.	.	*	*
All	276,438	193	1,829	9,511	13,893	39	589	93,875	396,367

The point prevalent cohorts and associated modalities are determined on December 31 of each calendar year without applying the 60-day stable modality rule.

* Values for cells with 10 or fewer patients are suppressed.

." Zero values in this cell.