

# Chapter XV

## Analytical Methods

This chapter discusses the analytical methods used in this 1994 Annual Data Report (ADR), and the differences between this ADR and the 1993 ADR. This chapter should be read in conjunction with Chapter XIV, which describes the USRDS database. The *Researcher's Guide to the USRDS Database* provides additional detail about the database and the Standard Analysis Files.

---

### Differences Between 1993 and 1994 Reports

**1) New Data.** The largest change is the addition of a new year of data. Data are now reported through 1991 for patient data and through 1992 for provider-based data. There are also changes in data which apply to years reported in past ADRs. This includes data which are delayed in reaching the USRDS database, and new sources of data, such as increased reporting by Department of Veterans Affairs facilities. See "New Data in the Database" below.

**2) Revised death and hospitalization rate methodology.** The methodology for computing annual death rates in Section D of the Reference Tables has been revised in response to feedback from users of these tables. The

tables now include incident patients and dialysis patients who have had a prior transplant. Death rates for patients with a functioning transplant are computed separately using a somewhat different methodology. See "Annual Death Rate Methodology" below.

**3) Standard Populations.** The 1990 Census is now used as the standard population for adjustment of incidence and prevalence rates. Prior ADRs used the 1980 Census. For adjustment of survival rates, the standard population is drawn from the ESRD population rather than from the overall population. We have adopted the standard of using the most recent year. See "Adjustment or Standardization of Rates" below.

**4) Patient Residence.** In the 1993 ADR, for prevalent counts, patients were reported by state and ESRD Network based on the place of treatment. The USRDS now contains a residence history for each patient, so patients are now always reported based on the place of residence. The residence history goes back to the address reported at the start of ESRD for each patient, but the date at which the patient moved is not precisely known. See "Geography" below.

**5) Death Notification Form** A new ESRD Death Notification Form (HCFA-2746) came into use during 1990. More

detailed cause of death categories are used on the new form, and stoppage of ESRD treatment prior to death is asked separately from the cause of death question. For most cause of death tables in Section D of the Reference Tables, the new codes are recoded into the old coding system. However, we now have one full year (1991) of data which uses primarily the new codes, so a table of death rates by the new codes is also provided. Chapter VII discusses the impact of this change. See also "Death Notification Form" below.

**6) Minimum Aggregation Size.** In the Reference Tables and in the graphics and tables in the text, aggregations of less than 10 persons generally are not reported. This brings the ADR into compliance with Section F of the USRDS Agreement for Release of Data, which is included in Chapter II.

This restriction is imposed to insure the confidentiality of patient-specific data. In a cross-tabulation of patients by patient characteristics, if an individual cell in the table represents only one person, then it is conceivable that one could identify that person and learn something about that person. By restricting cell sizes to 10 or more persons, such identification becomes virtually impossible. The general convention in the Reference Tables is to replace the number with an asterisk (\*) in cells which are based on less than 10 patients.

**7) Standard Analysis Files.** The Standard Analysis Files (SAFs) described in Chapters II and XIV are the starting point for almost all of the tables and analyses presented in this ADR. This clearly illustrates the ability of the

SAFs to support a broad range of research topics and methodologies.

**8) Hospital Length of Stay** The algorithm used for calculating hospital length of stay has been changed to conform to the method generally used in hospital utilization research. This affects Section H of the Reference Tables. See "Hospitalization" below

These changes in data, definitions, and methods mean that care must be used when comparing data between ADRs for different years. However, this ADR follows the USRDS policy that each ADR present data for a series of years using a consistent set of definitions so that the reader can make valid comparisons among years within that report.

---

## New Data in the Database

Table XV-1 shows the incidence counts by year as reported in the 1989, 1990, 1991, and 1993 ADRs. Each ADR shows higher incident counts for all years than did the previous ADR. The greatest increases occur in the most recent two years.

Table XV-2 presents these changes from a different perspective. The years are shown relative to the year of the update rather than as calendar years. Table XV-2 illustrates the reason for this: the data for the year prior to an update show changes on the order of 8 percent to 14 percent on the next update. The incidence counts for earlier years are minimal.

The increases in incidence counts for past years from update to update may be due to at least three factors:

**Incidence Counts by Year by Database Update Year and Change in Counts  
Between Updates For Updates from 1989 through 1993**

Incident Year	Year of Update										
	1989		1990		1991		1992		1993		Cumulative Change
	(1989 ADR)		(1990 ADR)		(1991 ADR)		(1992 ADR)		(1993 ADR)	(1994 ADR)	
1983	24929	0.25%	24991	0.12%	25020	0.57%	25,163	0.02%	25169		0.96%
1984	26336	0.24%	26399	0.29%	26475	0.59%	26,631	0.19%	26681		1.31%
1985	29156	0.25%	29228	0.46%	29361	0.79%	29593	0.18%	29647		1.68%
1986	31195	0.64%	31394	0.64%	31594	0.68%	31,810	0.21%	31876		2.18%
1987	33578	1.43%	34058	1.10%	34433	0.88%	34,737	0.49%	34908		3.96%
1988			36160	2.69%	37133	1.59%	37,723	0.72%	37994		5.07%
1989					41269	2.44%	42,278	1.00%	42702		3.47%
1990					39863	14.76%	45,745	1.56%	46458		1.56%
1991					1938		46803	8.30%	50689		
1992							5480		50523		
1993									5746		

Note: The horizontal line indicates the last year reported from this update. The cumulative change is measured from the year in which the incident year was first reported.

**Table XV-1**

**Delays in receiving data.** This is probably the main factor in the most recent years. The USRDS generally waits 15 months before reporting data for a given time period because of this problem. Table XV-1 suggests that we are missing more than 8 percent of new patients 3 months after the end of the

year, about 3 to 7 percent after 15 months, and about 1.5 to 3 percent after 27 months. We can expect the incidence count for 1991 to be 2 percent to 3 percent higher when reported in the 1995 ADR and another 1 percent higher when reported in the 1996 ADR.

**Change in Yearly Incident Counts From Prior Update,  
by Number of Years Before Update Year**

Prior Year	Year of USRDS Database Update				Average
	1990 (1990 ADR)	1991 (1991 ADR)	1992 (1993 ADR)	1993 (1994 ADR)	
-9			0.00%	0.02%	0.01%
-8			0.57%	0.19%	0.38%
-7		0.12%	0.59%	0.18%	0.30%
-6	0.25%	0.29%	0.79%	0.21%	0.38%
-5	0.24%	0.46%	0.68%	0.49%	0.47%
-4	0.25%	0.64%	0.88%	0.72%	0.62%
-3	0.64%	1.10%	1.59%	1.00%	1.08%
-2	1.43%	2.69%	2.44%	1.56%	2.03%
-1			14.76%	8.30%	11.53%

Example: The outlined cell refers to the second year before 1993 (1991). The incidence count reported for 1991 increased by 1.56% from the 1992 update (reported in 1993 ADR) to the 1993 update (reported 1994 ADR). Data below the line are not reported.

**Table XV-2**

An example of a delay in receiving data is a patient who is covered under private health insurance and for whom Medicare would be the secondary payer for the first 21 months. If the private health insurance covers all of the dialysis costs, then the Medical Evidence Form might not be filed until a year or more after the start of ESRD. Since there are no Medicare claims, the patient will not appear in the USRDS database until the Medical Evidence Form is filed. When the form is filed, the date of start of chronic renal dialysis entered on the form should be the date dialysis started, even though it was not covered by Medicare at that point. Thus this patient appears as an incident case in the earlier year rather than the year in which the form was filed.

**New data sources.** The data starting with the 1993 ADR include Department of Veterans Affairs (DVA) patients being reported to HCFA for the first time. This is one reason the percentage increases shown in Table XV-1 are higher for the 1993 ADR than for prior ADRs.

**Delays in Medicare eligibility.** If a patient is covered by some other insurance (such as DVA) for four years and then becomes eligible for Medicare and has a Medical Evidence Form filed, his/her first service date would be four years prior to appearing in the file.

**Errors in first service date.** The physician completing the Medical Evidence Form may report the first dialysis ever rather than the start of maintenance dialysis. There may be clerical or keypunch errors in the year of the first dialysis date. The software for determining first service date may be wrong. Thus, patients added in a given

update can appear as incident in a much earlier year. If a first service date is corrected, then a patient may appear as incident in a different year than in a previous database update.

### **The “Fifteen Month Rule”**

The USRDS generally does not report data for periods later than 15 months prior to the update date, which is generally in April or May. Because of delays in the submission and processing of Medicare bills, these data generally are not considered to be complete until 15 months have passed. Thus, although we have data through May 1993, the data are only considered complete through February 1992, and this ADR reports results only through December 31, 1991.

The USRDS CC is currently evaluating this rule. Some analyses in this ADR use data from 1992. These analyses are labeled as preliminary and should be treated with caution.

---

### **Death Notification Form**

In March of 1990, a new ESRD Death Notification Form (HCFA-2746) came into use. As Table XV-3 shows, a few deaths from 1989 were reported (in 1990) using the new form. Both forms were in use in 1990, after which only a handful of old forms were filed.

The new form uses a new set of codes and categories for cause of death. As Table XV-4 shows, these codes were designed so that they could be recoded as pre-1990 categories. Another substantial change is that withdrawal from dialysis is no longer a cause of death, but a separate question asks whether the ESRD treatment was

<b>Death Notification Forms By Year of Death, By Form Type and By Cause of Death Using Pre-1990 Categories</b>				
<b>Form Type</b>	<b>Year of Death</b>			
	<b>1989</b>	<b>1990</b>	<b>1991</b>	<b>1992</b>
New Form	556	12134	27572	24576
Pre 1990 Form	23381	13567	16	14
Missing form	4190	4830	5645	5397
Total	28127	30531	33233	29987

  

<b>Primary Cause of Death, Pre-1990 Coding</b>				
	<b>1989</b>	<b>1990</b>	<b>1991</b>	<b>1992</b>
1 Pericarditis(incl. Cardiac Tamponade)	0.73%	0.56%	0.25%	0.20%
2 Myocardial Infarction, Acute	14.00%	12.82%	11.44%	10.83%
3 Cardiac (Other Than 01 Or 02)	29.26%	30.05%	31.22%	32.29%
4 Cerebrovascular	6.14%	5.58%	4.86%	4.79%
5 Embolism, Air	0.02%	0.02%	0.02%	0.03%
6 Embolism, Pulmonary	0.51%	0.50%	0.41%	0.40%
7 Gi Hemorrhage	1.52%	1.58%	2.00%	2.00%
8 Vascular Access	0.13%	0.16%	0.13%	0.13%
9 Hemorrhage (Other Than 04, 07, Or 08)	0.66%	0.74%	0.74%	0.74%
10 Pulmonary Infection	3.28%	3.17%	2.55%	2.31%
11 Septicemia	9.93%	9.57%	9.65%	9.59%
12 Viral Hepatitis	0.12%	0.07%	0.03%	0.04%
13 Infection (Other Than 10, 11, Or 12)	1.62%	1.33%	1.28%	1.20%
14 Hyperkalemia	1.19%	1.14%	1.10%	1.17%
15 Pancreatitis	0.31%	0.26%	0.21%	0.26%
16 Malignancy	3.28%	3.23%	2.75%	2.78%
17 Withdrawl From Dialysis	9.70%	12.61%	16.36%	16.32%
18 Suicide	0.28%	0.25%	0.20%	0.18%
19 Accidental, Treatment Related (Not 05)	0.04%	0.04%	0.01%	0.02%
20 Accidental, Not Treatment Related	0.43%	0.30%	0.30%	0.31%
21 Unknown Cause	8.40%	7.40%	6.26%	6.09%
22 Other (Specify In Remarks)	8.45%	8.61%	8.20%	8.32%

**Table XV-3**

stopped before the patient died and the reason for such stoppage.

Table XV-3 shows the distribution of causes of death from 1989 through 1992 using the pre-1990 coding. The most significant change caused by the new coding is an increase in the withdrawal from dialysis category, which rose from 9.7 percent to 16.3 percent from 1989 to 1991. If the new form indicates that that ESRD treatment was stopped before

death, then the cause of death is recoded to the old withdrawal from dialysis category. It is plausible that most of this change is due to the change in the form rather than to a change in medical practice.

### Recoding of New Cause of Death Codes Into Pre-1990 Categories For Death Notification Forms for Deaths in 1989-1992

Pre-1990 Coding	Coding on New Form	COUNT
Pericarditis (incl. Cardiac Tamponade)	1 25 Pericarditis, Incl. Cardiac Tamponade	160
Myocardial Infarction, Acute	2 23 Myocardial Infarction, Acute	7330
Cardiac (Other Than 01 Or 02)	3 26 Atherosclerotic Heart Disease	2472
	27 Cardiomyopathy	2349
	28 Cardiac Arrhythmia	3959
	29 Cardiac Arrest, Cause Unknown	10861
	30 Valvular Heart Disease	359
	31 Pulmonary Edema Due To Exogenous Fluid	425
Cerebrovascular	4 36 Cerebro-Vascular Accident Including Intracranial Hemorrhage	2682
	37 Ischemic Brain Damage/Anoxic Encephalopathy	441
Embolism, Air	5 89 Air Embolism	16
Embolism, Pulmonary	6 35 Pulmonary Embolus	277
Gi Hemorrhage	7 44 Mesenteric Infarction/Ischemic Bowel	536
	72 Gastro-Intestinal Hemorrhage	718
Vascular Access	8 39 Hemorrhage From Vascular Access	96
Hemorrhage (Other Than 04, 07, Or 08)	9 38 Hemorrhage From Transplant Site	9
	41 Hemorrhage From Ruptured Vascular Aneurysm	193
	42 Hemorrhage From Surgery (Not 38,39 Or 40)	47
	43 Other Hemorrhage (Not Codes 38-42,72)	238
Pulmonary Infection	10 53 Pulmonary Infection (Bacterial)	1258
	54 Pulmonary Infection (Fungal)	72
	55 Pulmonary Infection (Other)	301
Septicemia	11 49 Septicemia, Due To Vascular Access	636
	50 Septicemia, Due To Peritonitis	657
	51 Septicemia, Due To Peripheral Vascular Disease, Gangrene	1201
	52 Septicemia, Other	3728
Viral Hepatitis	12 64 Hepatitis B	21
Infection (Other Than 10, 11, Or 12)	13 56 Viral Infection, Cmv	38
	57 Viral Infection, Other (Not 64 Or 65)	49
	58 Tuberculosis	25
	59 Aids	362
	60 Infection, Other	359
Hyperkalemia	14 24 Hyperkalemia	740
Pancreatitis	15 73 Pancreatitis	152
Malignancy	16 82 Malignant Disease, Patient Ever On Immunosuppressive Therapy	596
	83 Malignant Disease (Not 82)	1249
Withdrawl From Dialysis	17 All codes if treatment stopped before death	10402
Suicide	18 92 Suicide	135
Accidental, Treatment Related (Not 05)	19 40 Hemorrhage From Dialysis Circuit	7
	90 Accident Related To Treatment	7
Accidental, Not Treatment Related	20 91 Accident Unrelated To Treatment	194
Unknown Cause	21 99 Unknown	4116
Other (Specify In Remarks)	22 65 Other Viral Hepatitis	43
	66 Liver-Drug Toxicity	7
	67 Cirrhosis	159
	68 Polycystic Liver Disease	6
	69 Liver Failure, Cause Unknown Other	190
	74 Fungal Peritonitis	27
	75 Perforation Of Peptic Ulcer	54
	76 Perforation Of Bowel (Not 75)	215
	80 Bone Marrow Depression	18
	81 Cachexia	413
	84 Dementia, Incl. Dialysis Dementia, Alzheimers	171
	85 Seizures	128
	86 Diabetic Coma, Hyperglycemia, Hypoglycemia	150
	87 Chronic Obstructive Lung Disease (COPD)	252
	88 Complications Of Surgery	152
	93 Drug Overdose (Street Drugs)	45
	94 Drug Overdose (Not 92 Or 93)	22
	98 Other Identified Cause of Death, Please Specify	3313

Table XV-4

---

## Geography

Starting with the 1993 ADR, Puerto Rico and United States Territories have been reported separately from the 50 states and the District of Columbia in the incidence and prevalence Reference Tables (Tables A.1, A.3, B.1, B.4).

The population base used in computing incidence and prevalence rates is United States resident population, which includes only the 50 states and D.C. Thus, incidence rates in prior ADRs were overstated by about one percent because of the inclusion of Puerto Rico and United States Territories in the numerator.

The 1990 Census provides population counts for Puerto Rico for calculating 1990 incidence and prevalence rates. Crude rates are computed by age and by sex. Rates are not computed by race since the Census Bureau does not report the population of Puerto Rico by race. The population of Puerto Rico is more racially mixed than is the overall population of the United States Rates, so that classification by race is not meaningful. Adjusted rates are not computed for Puerto Rico because of the lack of the race variable. Rates are not computed for earlier years because we do not have a good longitudinal series of population data for Puerto Rico by age and sex.

Puerto Rico, the Virgin Islands, and the Pacific islands are not included in computing the incidence and prevalence rates in Sections A and B for the New Jersey and Northern California ESRD Networks, respectively. They also were not included in the 1991 ADR incidence and prevalence rates by Network. In

Reference Tables sections other than A and B, all patients are counted, and tables by Network include Puerto Rico, the Virgin Islands, and the Pacific Islands.

In the 1993 ADR prevalence tables, patients were classified based on the location of the facility at which they were being treated. Since that time, the USRDS has constructed a file of the place of residence of each patient since the start of ESRD. Using this file, we are able to do all tabulations in the 1994 ADR based on the place of residence of the patient at the point in time relevant to the particular analysis.

Table XV-5, repeated from the 1993 ADR Reference Tables, shows the distribution of patients by state of residence and state of treatment. This table indicates the magnitude of the effect this change is likely to have on prevalence counts by state.

A patient may be treated in a state different from the state of residence for a number of reasons. The patient may live near the state border, and the closest facility may be in the neighboring state. The patient may be traveling and be temporarily treated in another state. The patient may change residence seasonally.

This difference may also be an artifact of imprecision in the date of change of residence. One limitation of the new longitudinal residence record is that it is based on residence at no more than one or two points in time each year back to 1987 and one point during 1982. Thus the exact dates at which the patient moved are not known. The imprecision of this record must be taken into account when using the Residence SAF.

---

**Measurement Concepts: Incidence, Acceptance, and**


---

**Percentage of Patients Treated in Home State and Other States  
By State of Residence  
For Patients Alive on December 31, 1990**

Residence	Treatment	%	Residence	Treatment	%	Residence	Treatment	%
Alabama	AL	95	Louisiana	LA	97	Oklahoma	OK	91
Alaska	AK	79	Maine	ME	90		TX	3
	WA	10		FL	3		AR	2
	CA	3		NH	3	Oregon	OR	94
	OR	3	Maryland	MD	90		CA	2
	FL	1		DC	7	Pennsylvania	PA	96
	IL	1	Massachusetts	MA	94	Puerto Rico	PR	96
	IN	1		RI	2	Rhode Island	RI	96
Arizona	AZ	95	Michigan	MI	95	South Carolina	SC	93
	CA	1	Minnesota	MN	95		GA	3
Arkansas	AR	88		ND	2	South Dakota	SD	72
	TX	4	Mississippi	MS	92		MN	23
	TN	3		TN	4		IA	2
California	CA	97	Missouri	MO	95	Tennessee	TN	96
Colorado	CO	91		KS	2	Texas	TX	98
	NM	2	Montana	MT	83	Utah	UT	95
	CA	1		WA	5		CA	1
	TX	1		UT	2	Vermont	VT	63
Connecticut	CT	94		CA	2		NH	20
	RI	2		MN	2		MA	11
Delaware	DE	90		ND	1		FL	3
	PA	4		OR	1	Virgin Islands	VI	74
	MD	4		CO	1		PR	14
District of Col	DC	89		PA	1		FL	2
	MD	8	Nebraska	NE	92		NJ	2
Florida	FL	95		IA	3		NY	2
	NY	1		CO	2		SC	2
Georgia	GA	95	Nevada	NV	92	Virginia	VA	93
Hawaii	HI	97		CA	3		DC	2
Idaho	ID	69		UT	2		TN	2
	WA	17	New Hampshire	NH	77	Washington	WA	95
	UT	6		MA	16		OR	2
	CA	3		FL	2	West Virginia	WV	85
Illinois	IL	92	New Jersey	NJ	91		VA	6
	MO	2		PA	4		OH	3
	IN	2	New Mexico	NM	90		MD	2
Indiana	IN	94		TX	7	Wisconsin	WI	93
	KY	2	New York	NY	96		MN	4
Iowa	IA	85	North Carolina	NC	96	Wyoming	WY	45
	MN	6	North Dakota	ND	79		CO	23
	NE	4		MN	16		UT	20
Kansas	KS	84		SD	1		MT	5
	MO	11	Ohio	OH	93		SD	2
Kentucky	KY	90		WV	2		ID	1
	OH	4						
	TN	2						

Note: For each state of residence, states of treatment are listed in order of descending percentage until the sum reaches 95%. The remaining percentage is accounted for by one or more other states of treatment with smaller percentages than the last state listed. Some of the differences are due to imprecision in the date of change of residence.

---

**Table XV-5**

## Prevalence

Incidence and prevalence are two concepts used in measuring the extent of a disease. The incidence of a disease is the number of persons diagnosed with that disease in a given time period, typically a year. Prevalence is the number of persons who have that disease at a given point in time (point prevalence) or who have the disease during a given time period (period prevalence). Thus, annual period prevalence consists of persons who have the disease at the end of the year plus those who had the disease at some time during the year and died before the end of the year. The prevalence measures combine the effects of those who get the disease (incidence) and those who die (mortality).

Note that the USRDS considers successful transplantation as a therapy rather than a “recovery” from ESRD.

The terms “incidence” and “prevalence” of ESRD imply that information is available about all patients with ESRD, but, as discussed earlier, this is not the case. Actually, the data are for persons receiving ESRD therapy as reported through HCFA and do not include patients who die of ESRD before receiving treatment and those who are not reported through HCFA. For these reasons, the terms incidence and prevalence are qualified as incidence and prevalence of reported ESRD therapy.

As discussed earlier and in Chapter XIV, patients are classified as lost to follow-up if one year has passed with no dialysis, death, or transplant data. Starting with the 1992 ADR, these patients are not included in the point prevalence counts. The Reference Tables

show the number of patients not counted for this reason.

The term “acceptance into ESRD therapy” is used by some other ESRD registries, such as the European Dialysis and Transplantation Association. The USRDS chose not to use this term because “incidence of reported ESRD therapy” is more precise, and because “acceptance” implies that the remaining patients are rejected, when in fact they may be unidentified as ESRD cases or unreported through HCFA.

Incidence tends to be more a useful measure for medical and epidemiological research which examines the causes of disease and the differences in how sub-populations are affected by these causes. Point prevalence may be more useful for public health research, since it measures the overall burden of the disease on the health care delivery system. Period prevalence may be more useful for cost analysis, since it indicates the total burden of the disease over the course of the year.

Sections A and B of the Reference Tables present parallel sets of counts and rates for incidence and December 31 point prevalence. Section B also presents annual period prevalence counts (but not rates). This de-emphasis of period prevalence reflects our experience over the past year with the general lack of demand for more than overall counts of period prevalence.

## Census Population Base

Incidence and prevalence counts indicate the absolute level of occurrence of ESRD as defined by its treatment. Incidence and prevalence rates per million population indicate the level of

the disease relative to the population and provide a convenient method of comparing the occurrence of the disease over time, among population subgroups, and with other diseases. This report generally uses rates per million population, although for some tables rates per ten million population are used. The size of the population base is chosen largely to produce rates with three significant digits to the left of the decimal point. Rates for diseases which occur more frequently than ESRD might use a smaller population base size.

Note that rates for a specific age/sex/race population group use that group for both the numerator and denominator of the rate. Thus the incidence rate for 40-45 year old white males uses the Census population counts for 40-45 year old white males as the base. Rates by primary disease causing ESRD for 40-45 year old white males also use the Census population counts for 40-45 year old white males as the base.

The population base for computing incidence rates is the total United States resident population on July 1 of each year by race, sex, and age. For point prevalence rates, the population on December 31 is used. Race is divided among four race groups: black, white, Asian, and Native American. Age is divided among eighteen five-year age groups: the first group includes persons who are 0-4 years of age, while the 18th group includes persons who are 85 years of age and older. For this ADR, rates are reported for 1982 through 1991. Section J of the Reference Tables shows the July 1 population counts. The December 31 counts were computed as the midpoint

between the July 1 counts for adjacent years.

The population data used for this 1994 ADR are based on data from the 1990 Census and from the Census Bureau's Current Population Reports. When the 1990 Census data became available, the Census Bureau revised the estimates of the overall population for the 1980's. The estimate of overall population was only 0.6 percent under the population counted by the Census. However, the differences were much higher for some subgroups. For instance, the Native American population was almost 15 percent higher in the 1990 Census than in the estimates reported in the Current Population Reports before the 1990 Census was tabulated. The Asian/Pacific Islander population was 3 percent higher.

For the 1993 ADR, the USRDS recomputed all population counts from 1980 to 1990. The differences between the new counts and those used for the 1980s in earlier ADRs are very small for 1980 and increase steadily over the decade. The process involved the following basic steps:

1. Establish July 1, 1980, counts by age/race/sex group to be used as the starting point.
2. Establish July 1, 1990, counts by age/race/sex group to be used as the endpoint.
3. Linearly interpolate 1990 counts by age/race/sex group for intervening years.
4. Adjust each year so that the totals match published totals for that year.

The starting counts are, with a slight adjustment to be described below, those

reported for July 1, 1980, in the 1991 ADR. These counts were derived by combining data from United States Bureau of the Census Current Population Reports (1990, Nos. 1045 and 1057) and the 1980 Census (1983).

The Current Population Reports give population estimates by year, by one and five-year age groups, sex, and black, white, and "other" race groups. In 1990, population estimates by year and sex were published for Native American/Alaskan Native and Asian/Pacific Islander race groups. The "other" race category is divided entirely between these two race groups. In making these estimates, the Census Bureau allocated persons with race "unknown" or other than black, white, Native American, or Asian into one of these four groups. The treatment of race is discussed in more detail below.

In order to use all four race groups for computing and adjusting rates, it was necessary to estimate population by five-year age groups for the Native American and Asian groups. The Native American population estimates for each year were allocated to age groups using the 1980 Census age distribution for Native Americans. (Bureau of the Census, 1983). The Asian population estimate for each year was then allocated to age groups by subtracting the estimate of the Native American population for each age group from the entire Census Bureau estimate of the "other" race population for that age group.

In May 1992, the Current Population Reports published estimates of the total United States resident population for every January 1 and July 1 between 1980 and 1992 (Bureau of the Census, 1992). The estimate for July 1, 1980, differed

slightly from the total derived via the process described above and published in the 1991 ADR. Accordingly, the counts for all subgroups were adjusted proportionately to bring the total for 1980 into agreement with the Current Population Reports estimate for July 1, 1980.

The counts for July 1, 1990 were derived from the Bureau of the Census Modified Age/Race, Sex and Hispanic Origin (MARS) file (Bureau of the Census, 1992). Based on the 1990 Census, this file contains 1616 counts (101 ages by 2 sexes by 4 races by Hispanic/non-Hispanic) for every county and state on April 1, 1990. These data were aggregated into the USRDS age and race groups, and the total for all subgroups were multiplied by a constant to bring the total to the Current Population Reports estimate for July 1, 1990.

Counts for points between July 1, 1981 and July 1, 1990, were derived by first applying straight-line interpolation. All subgroups in each year were multiplied by a constant factor to bring the total for the year to the Current Population Reports estimate of the total population for that date. The January 1 Current Population Reports estimates were used for December 31 of the prior year.

For 1990 only, incidence and prevalence rates are presented by state and Network. The July 1 and December 31 population counts used as a base were derived from the 1990 MARS file and the Current Population Reports estimates in the manner described above.

## Race Classifications

The race codes in the USRDS database classify patients into the categories of Native American/Alaskan Native, Asian/Pacific Islander, Black, White, Other, and Unknown. As mentioned earlier, the Native American and Asian categories became available only with patients starting ESRD therapy in 1982. Since 1982, the proportion of new patients in the Other and Unknown categories has remained under two percent.

Incidence and prevalence rates are computed and adjusted by race using only the following categories: black, white, Asian, and Native American. Only counts are reported for the “other” and “unknown” categories, and these categories are dropped from all tables which report rates by race. In addition, rates by race are only reported for years starting with 1982 because of the change in the reporting of race that year.

Only four categories are used in computing incidence and prevalence rates because the Census population estimates used in computing the rates place the entire population into four categories and do not include a residual “other” or an “unknown” category. Since almost two percent of the patients in the USRDS database fall into the “other” and “unknown” categories, the rates reported in this report will be biased slightly low. If bias is uniform across races, then comparisons of rates by race are not significantly biased.

The USRDS is not able to produce tabulations for the Hispanic population because it does not have this data item for all patients. An item on Hispanic ethnicity will appear on the new HCFA

Medical Evidence Form, which will go into use in 1995, so that this item will be available for most patients starting ESRD in 1995 or later.

---

## Adjustment and Standardization of Rates

Adjustment or standardization of rates is a method for comparing rates between different sub populations or time periods to remove the effect of differences in the composition of the populations being compared. For instance, blacks with ESRD have a younger age distribution than do whites. One way to remove the effect of this age difference on observed rates is to compare the rates for specific age groups between blacks and whites. Age adjustment provides a method of obtaining summary rates (across age groups) for blacks and whites with the effect of age differences removed.

Age adjusted rates indicate what the rates for the two groups, for example blacks and whites, would have been if the age specific rates for each age group had occurred in the same standard population. Age adjustment operates by taking the age specific rates for blacks, applying them to a standard population distribution stratified by age, summing the resulting ESRD population, and computing a new overall rate per million population using the total of the standard population. The same procedure is followed for whites. After age adjustment, the overall 1986 incidence rate for blacks changes from 300 to 373, and that for whites changes from 105 to 99. Blacks have higher rates of ESRD than whites at almost every age group, but the fact that the black population has a younger age distribution masks some

of this difference in the across age unadjusted rates, since there are relatively fewer patients in the younger age groups.

The standard population used in this report for adjustment of incidence and prevalence rates is the July 1, 1990, United States resident population by five-year age groups, four race categories, and sex. Prior ADRs used 1980 population as the standard.

Starting with this ADR, the standard population used for adjustment of patient survival probabilities will be the ESRD incident population for the most recent year. This year it is the 1991 ESRD incident population by five-year age groups, three race groups, sex, and four primary disease groups. This ESRD population allows adjustment to be made for primary disease as well as for race, sex, and age. Prior ADRs used the 1985 incident ESRD population as the standard population.

The adjustment method described here is the “direct standardization” method and is the method used through most of this report. For the rates by state and ESRD Network in Sections A and B of the Reference Tables, the “indirect standardization” method is used because of problems with small population counts (Fleiss).

In addition to adjustment by age, the tables in this report frequently adjust by race and sex or appropriate combinations of the three factors. For instance, in one of the standard table formats, rates by age are adjusted by race and sex, rates by race are adjusted by age and sex, rates by sex are adjusted by age and race, and rates by primary diagnosis and overall rates by year are adjusted by age, race,

and sex. Adjustment for multiple factors requires stratifying the populations by the multiple factors rather than just by age. Such adjustments are particularly useful when comparing rates over time to control for the changing characteristics of the ESRD population. Most tables showing adjusted rates are followed by a parallel table showing standard errors of the adjusted rates.

It should be noted that the process of standardization across age, race, and sex produces rates that are different than those encountered if considering one rate separately. For example, someone familiar with the total ESRD incidence rate for 1987 would find the adjusted rate (135/million population, see Reference Table A.6) to be lower than the real or “crude” rate (139/million). This difference does not indicate that the standardized rate is incorrect; rather, the standardized rate is “adjusted” so that rates for 1987 can be compared to rates for all of the other years (1982-90), regardless of changing age, race, and sex compositions of the ESRD population during these years. Thus, while standardized rates are not “real” in one sense, they are the most useful measures for comparing rates across strata or years.

---

## Annual Death Rate Methodology

Section D of the Reference Tables presents death counts and death rates expressed as the number of ESRD patient deaths per 1,000 patient years at risk. The death rates are computed based on calendar year periods. In order to increase the stability of the estimated death rates, three years (1988-1990) are pooled for the tables. Death rates are

presented by age, race, sex, and primary disease.

This death rate methodology has been used by the USRDS since 1991. It was designed to be used by ESRD Networks and dialysis and transplant facilities based on data about a group of patients. These death rate tables can be used to compare local ESRD mortality rates to national rates. (Wolfe) The time period for the local data does not need to be a calendar year and can be more or less than one year in length as long as the period at risk is measured in units of years and all other aspects of the methodology are followed.

Tables of death rates by cause of death age, race, sex, and primary disease are also included.

In response to suggestions from users of this methodology, some major changes have been made starting with this year: patients starting ESRD therapy during the year are included, previously transplanted dialysis patients are included, different categories of patients are used, and a different methodology is used for patients with functioning transplants.

**Incident Patients.** The original methodology was based only on patients alive on January 1 who had reached day 91 of ESRD and followed those patients for the rest of the calendar year. Patients who entered ESRD therapy during the year did not enter the analysis until January 1 of the following year. This means that one third of the patients in a given dialysis unit are likely to be excluded from the analysis. In the new methodology, patients who reach day 91 of ESRD during the year are included in the analysis and are followed from that

day until the end of the calendar year (or until the censoring date for other time periods).

**Previously Transplanted Patients.** In the original methodology, previously transplanted dialysis patients were excluded. The 1992 ADR reported that previously transplanted dialysis patients had a nine percent lower mortality rate than did never-transplanted dialysis patients and accounted for 13 percent of all dialysis patients. In order not to lose those 13 percent from the analysis, the new methodology has patients re-enter the analysis on the January 1 after a transplant failure. In the future, this may be revised to have the patient re-enter the analysis 60 days after the transplant failure.

**New categories.** Below is a list of the categories for which mortality rates are reported in this ADR, along with an explanation of the changes in the categories:

- All dialysis patients. *No longer restricted to those who have not received a transplant.* The period at risk is censored at transplant date if a transplant occurs during the year.
- Dialysis patients who are on hemodialysis (including home hemodialysis) at the start of the period and who have been on this modality for at least 60 days. *No longer restricted to those who have not received a transplant.* The period at risk is censored at the time of a new transplant.
- Dialysis patients not yet transplanted who are on CAPD or CCPD at the start of the period

and who have been on this modality for at least 60 days. *No longer restricted to those who have not received a transplant.* The period at risk is censored at the time of a new transplant.

- All ESRD patients.

The “All ESRD” category will be less useful than the others but is included for completeness and as an overall reference. The following categories have been replaced by the separate methodology for functioning graft patients, which is described later.

- All patients with a functioning transplant.
- Patients with a functioning cadaveric first transplant.

The cohort of patients used in this method is defined as those who are alive and not lost to follow-up at the beginning of a year, and who have had ESRD for at least 90 days, plus those patients who reach day 91 during the year. A patient who dies on December 31 of the prior year is excluded, but a patient who dies on January 1 is included with one day at risk. In effect, we are assuming that all deaths occur at 11:59 PM on the death date. Eliminating lost to follow-up patients is a change introduced with the 1992 Annual Data Report.

### **Annual Death Rates for Dialysis and All ESRD**

All patients who have reached day 91 of ESRD on January 1 or who reach day 91 later in the year are included and are followed until the end of the calendar year. The starting date is either January 1 or the day the patient reaches day 91 of ESRD.

Patients who have a functioning transplant on the starting day are included only in the “All ESRD” category. Patients who are on dialysis on the starting date are also included in the “All Dialysis” category. A patient in the “All Dialysis” category may also be reported in one of two sub-groups (Hemodialysis or CAPD/CCPD) based on the dialysis modality on the starting date and whether the patient has been on that modality for at least 60 days. Dialysis patients who are not on hemodialysis or CAPD/CCPD or who have been on that modality for less than 60 days are included only in the “All ESRD” and “All Dialysis” categories. Note that a given patient may be included in one (“All ESRD” only), two (“All ESRD” and “All Dialysis”) or three (“All ESRD”, “All Dialysis”, and hemo or CAPD) categories.

For patients alive on January 1 to be classified as a hemodialysis or CAPD/CCPD patient, the patient must have been on the indicated treatment modality for 60 days before the start of the year. For incident patients, the 60 day test is applied on day 91. For patients returning from a transplant, the 60 day test is applied when they re-enter the analysis on the next January 1.

In determining whether the patient has been on the same modality for at least 60 days, a short (less than 60 days) spell on a different modality is not considered a change in modality if it is preceded and followed by periods of at least 60 days on the primary modality. See Chapter XIII of the 1992 ADR or the *Researcher’s Guide to the USRDS Database* for a description of the application of this “60 day rule.”

The denominator for the mortality rate is patient years at risk. For the "All Dialysis" category, the period at risk is from the starting date through the death date or December 31, whichever is earliest. For patients on dialysis at the beginning of the year, the period at risk is from the starting date through the death date, the date of the next transplant (if any during the year), or December 31 (again, whichever is earliest). Note that a dialysis patient who is transplanted during the year will have different periods of risks when he is used in the "All ESRD" and "All Dialysis" categories.

Note that the period at risk is not censored at the start of a lost to follow-up period during the year. Note also that if a patient is on dialysis and receives a transplant on January 1, then the days at risk are censored at the transplant date, and the patient has 0 days at risk.

Each patient contributes his/her years at risk to the denominator. The numerator is the number of deaths which occurred during the at risk periods. If death occurs during the at risk period for that patient, then the patient contributes a death to the numerator. Days at risk are divided by 365.25 to yield years at risk. The ratio is multiplied by 1,000 to yield deaths per 1,000 years at risk.

Years may be pooled in this method to increase the cell sizes, as is done in Section D of the Reference Tables. The analysis is still done in one year increments. A patient who is alive at the beginning of more than one of the pooled years will be used more than once in this method. A patient who dies during the second of the three years would contribute a full year at risk for the first year and a partial year at risk

and one death for the second year. For each year, the patient's age and treatment modality will be determined as of the beginning of that year. Thus, a patient could be classified as a 45 year old dialysis patient one year and as a 46 year old functioning transplant patient the next.

It should be noted that this method excludes about 14 percent of the deaths which occur in a given year. About 10 percent are excluded because the patient died before day 91 of ESRD, and about 4 percent are excluded because the patient is classified as lost to follow-up on January 1.

### **Annual Death Rates for Transplant Patients**

In previous ADRs annual mortality rates were computed for patients who had a functioning kidney transplant on January 1. These patients were treated in the same way as the all-ESRD category patients. Specifically, this means that the denominator for the mortality rate was the patient years at risk. For each patient the time in days from January 1 through the earliest of the death date and December 31 translated to years was the contribution to the denominator. One criticism of this methodology is that it misses the deaths of patients transplanted in the previous year who died in that year. Given that the first year post transplant has the highest mortality rate, it understates the mortality rate for transplant patients. In addition, it was felt that transplant facilities are more likely to prefer the transplant date as the starting point for mortality analyses.

In this ADR, annual death rates for the first year post transplant and for the

next two years post transplant are reported for patients transplanted in the previous three years, i.e. 1986-1988. The last transplant year is 1988 so that it is possible for all of the patients to experience three years post transplant, if they survive. The denominator for both rates is still the patient years at risk. For the first year rate, patient years are measured in days from the date of transplant to the death date or the end of the year for the first year at risk, i.e., 365 days exactly. For the rate for the second two years, the time at risk is measured from the anniversary date, i.e., exactly one year (365 days) following the transplant date, to the death date or the end of the period, i.e., exactly two years at risk.

Mortality rates for transplant patients with functioning grafts older than three years are not reported. In the earlier ADRs, such patients rates were mixed with survival rates for more recent transplants; this practice somewhat distorted the mortality rates.

---

## **Patient Survival Analysis Methodology**

All new Medicare eligible ESRD patients who passed the USRDS QC filters and who had a first service (dialysis or transplant) between January 1, 1977 and December 31, 1991, were included in the survival analysis. They were followed until December 31, 1992, giving a maximum follow-up of 15 years and a minimum follow-up of one year.

In each table in Section E, the survival rates for the most recent year are preliminary and must be used with caution. The analysis periods for the most recent year in each table extends

into 1992. The period for which data are considered complete for the current database update ends in February 1992. Thus, the death data for the most recent year in each table may be incomplete.

Patient survival probabilities and standard errors are estimated using the Kaplan-Meier method (KAPLAN) as implemented in the SAS LIFETEST (SAS) procedure and Greenwood's formula. The probabilities are expressed as percentages varying from 0 to 100 (rather than probabilities varying from 0 to 1).

Survival probabilities are computed for annual cohorts for the groups shown below. For the four groups which represent subsequent rather than initial modalities, the median times from ESRD to these modalities are also presented.

In the tables for all patients and for dialysis patients, the cohorts are defined as patients starting ESRD therapy in a calendar year and surviving at least 90 days after starting ESRD therapy. Thus, the one year survival probability is the probability of surviving from day 91 to one year plus 90 days (days 91 to 455, or 3 to 15 months). The two year survival probability is the probability of surviving from day 91 to two years plus 90 days. This delay is necessary because many patients under age 65 do not become eligible for Medicare for up to 90 days, and the database does not have data until the patient becomes eligible. Additional tables for patients 65 and over are presented, showing survival from date of ESRD to day 90 and from date of ESRD to day 365. The latter results can be compared with the earlier tables for survival from day 91 to day 455.

Given the likely selection bias involved in categorizing patients into modalities, patient characteristics may vary substantially between modality groups. Therefore, differences in survival across modality groups may not necessarily be ascribed to differences in the efficacy of the treatment modalities, even after adjustments for age, race, sex, and primary disease.

One, two, five, and ten-year Kaplan-Meier product limit estimates of survival probabilities were calculated for each age, race, sex, and primary disease group by the year of incidence.

Some survival results in Section E of the Reference Tables are adjusted for age, race, sex, and primary disease. Starting with this ADR, the standard population for these adjustments is the incident ESRD population for the most recent year; for this ADR, the most recent year is 1991. Previous ADRs used the 1985 ESRD incident population as the standard population.

The procedure used is the direct method of adjustment described earlier in this chapter and used for incidence and prevalence. The number of cells was limited to four age, two sex, three race, and four diagnosis groups. These limitations were imposed because more cells would have led to undesirable smaller sample sizes and the Kaplan-Meier survival estimates are biased high with small cell sizes.

### **All Patients**

The cohort consists of all ESRD patients starting renal replacement therapy in a calendar year and surviving beyond day 90. Patients are censored only at the end of follow-up (December 31, 1992).

### **Dialysis Patients (Censored at First Transplant)**

The cohort consists of all ESRD patients starting renal replacement therapy in a calendar year, surviving beyond day 90, and not receiving a transplant by day 91. Patients are censored at transplantation or at end of follow-up (December 31, 1992).

### **First Renal Transplant (Cadaveric)**

Patients in this cohort satisfy two conditions: 1) they received their first transplant in the designated year, and 2) the transplant donor is cadaveric. Transplants where the donor type is recorded as “other” or “unknown” are excluded from both the cadaveric and living related groups. The cohort is defined based on the year of first transplant regardless of the year of first ESRD service. Patients are followed from date of transplantation, and age is computed at the time of transplantation. Patients are censored only at end of follow-up (December 31, 1992). These patients appear in the “All ESRD” group above, and most also appear in the dialysis modality group. The median time in days from first ESRD service to first transplantation is also presented for these patients.

### **First Renal Transplant (Living Related)**

Patients in this cohort satisfy two conditions: 1) they received their first transplant in the designated year, and 2) the transplant donor is living related. Transplants where the donor type is recorded as “other” or “unknown” are excluded from both the cadaveric and living related groups. The cohort is defined based on the year of first

transplant regardless of the year of first ESRD service. Patients are followed from date of transplantation, and age is computed at the time of transplantation. Patients are censored only at end of follow-up (December 31, 1992). These patients appear in the "All ESRD" group above, and most also appear in the dialysis modality group. The median time in days from first ESRD service to first transplantation is presented for these patients.

---

### **Expected Remaining Lifetime Methodology**

Reported values for the expected remaining lifetime of a patient cohort are estimates of the life expectancy for the typical patient in a particular cohort. Without making assumptions about the parametric form of the distribution of lifetimes within a patient cohort, the expected remaining lifetime for a patient cohort cannot be estimated until all of the patients have died (Gross). In order to give timely (and thus practically useful) estimates of expected remaining lifetime, we truncated lifetimes at age 100.

---

### **Graft Survival Analysis Methodology**

The methodology for graft survival analysis is the same Kaplan-Meier methodology used for patient survival. For persons who have received multiple transplants, only the first transplant is analyzed. For the patient survival analysis, once a patient receives a transplant, that patient is thereafter classified as a transplant patient even if the graft fails. Graft survival analyses are presented separately for cadaveric and living related first transplants.

Transplants where the donor type is recorded as "other" or "unknown" are excluded.

The starting date for the graft survival analysis is the date of the first transplant. The ending date is the earlier of death or graft failure. The derivation of the graft failure date is discussed in Chapter XIV and in the *Researcher's Guide to the USRDS Database*.

Standard errors (after 90 days, 1, 2, 3, 5, and 10 years) are presented for both patient survival and graft survival analyses. The calculation of the standard error of the difference between two graft survival curves (Chapter VII) is the square root of the sum of the squared individual standard errors, since the covariance is zero. The p values reported in Chapter VII for the differences between survival curves are based on a t test of the difference in survival.

Some adjusted survival probabilities are computed using the same methods as for patient survival. Starting with this ADR, the standard population for these adjustments is patients who received a transplant in the past three years, 1989-1991 for this ADR. Prior ADRs used the 1985 ESRD incident population as the standard population.

---

### **Treatment Modality**

The treatment modality categories and the procedures for determining treatment modality are described in Chapter XIV of this ADR, in Chapters XI and XIII of the 1992 ADR, and in the *Researcher's Guide to the USRDS Database*. In Section C of the Reference Tables, the "60 day rule" discussed in Chapter XII is applied. This rule requires that the patient be on a new modality for

at least 60 days before it is considered a change in modality. Patients who are lost to follow-up are not included in the year end modality counts.

Section C of the Reference Tables also includes tabulations of patients by modality at two years. Lost to follow-up is included as a category in these tables.

---

## EPO Data

Three of the figures included in Chapter V (Treatment Modalities) show data about the use of erythropoietin (EPO) in conjunction with dialysis treatment in 1990 through 1992. This section describes the data and methodology used for these figures.

These data come from the data in the Quarterly Dialysis Records (QDR) in the USRDS database. The QDRs report on outpatient claims filed by institutions and do not report on claims filed by physicians and other Medicare Part B providers. Each QDR summarizes treatments received by one patient in one calendar quarter from one provider. A patient who uses more than one provider in a quarter will have more than one record for that quarter.

The treatment modality shown on the QDR is the last treatment modality in the quarter, and this is used to classify a patient by treatment modality for that quarter. Patients are classified as hemodialysis, CAPD/CCPD, or other. Patients with more than one QDR in a quarter are classified as "other" if the records do not all show the same modality.

**Figure V-15: Percent of Medicare Dialysis Patients Receiving Epo by Quarter, 1989-1992.** Estimates are

computed as  $epo\_patients / total\_patients$  for each modality, PD and HD. *Epo\_patients* is the sum of Medicare patients where the EPO administration field or the EPO dollars field in the QDR is greater than zero in that quarter. *Total\_patients* is the sum of any patients with a record in the QDR. Each patient is counted only once per quarter.

**Figure V-20: Hemodialysis patients receiving outpatient blood transfusions by Quarter, 1989-1992.** Estimates are computed as  $blood\_patients / total\_patients$ . *Blood\_patients* is the sum of patients where blood units received, reported in the QDR, is greater than 0 in that quarter. Total patients is the sum of any patients with a record in the QDR.

**Figure V-21: Average patient hematocrit and Epo Dose, by Quarter, 1989-1992. Center hemodialysis patients only.** Estimates are computed as mean hematocrit among patients for whom a hematocrit is reported and mean and median EPO dose among patients for who EPO dose is reported. Patients were classified as center hemodialysis excluding any HD patient where the dialysis setting field in the QDR was 'home'.

The EPO dose field on the QDRs is somewhat problematic. Before January 1991, the EPO dose field in the QDR was defined as the last EPO dose in the period. Since January 1991, the repeated dose has been defined as the average over the billing period. This average dose is calculated by HCFA using an input of the total EPO units from the claim.

Because patients in the QDR have one record per patient provider in that quarter, in the case of more than one record per patient, the latest of the EPO dose field and the hematocrit field were taken.

---

## Hospitalization

The USRDS database contains data on more than 2.7 million inpatient hospital stays. Section H of the Reference Tables reports on hospital days and hospital admissions per year at risk for hospitalization using a method which closely parallels the mortality rate method described above. A three year period is used in order to dampen the effect of short term variations.

In previous ADRs, the algorithm used for calculating hospital length of stay counted the day of admission as well as the day of discharge. In the 1994 ADR, this algorithm has been changed to count the day of admission but not the day of discharge. Patients discharged on the day of admission and patients discharged on the next day are counted as staying one day. This change was made to bring the ADR calculation into agreement with the Commission on Professional and Hospital Activities (CPHA) algorithm, which has been used extensively in hospitals and in hospital utilization research since 1955.

The cohort for this analysis is the same as for the mortality rate methodology, with the additional restriction that the date recorded in the HIM ID POST DATE variable must have been reached by the starting date (January 1 or day 91 of ESRD for this patient). This date defines the beginning of the period for which the hospitalization data are “complete.”

This period is constrained by Medicare eligibility and by the transmission of hospitalization data from the Medicare billing system to the HCFA ESRD PMMIS data system and then to the USRDS. From 10 percent to 15 percent of patients alive on January 1 are removed from the cohort by this requirement.

The term “complete” is placed in quotes because there is uncertainty about the quality and completeness of the hospitalization data. Known constraints on the completeness of the data are described below. For patients under 65 who are not already eligible for Medicare due to disability, Medicare eligibility may not start until up to 90 days after the start of therapy. During this period hospitalization data are not expected to be complete. Similarly, some transplant patients lose Medicare eligibility after three years of a successful transplant.

Hospital records are passed from the Medicare billing system to the ESRD PMMIS system if the record shows dialysis or a kidney transplant, or if both systems recognize the patient as an ESRD patient. The point at which this recognition occurs is most reliably indicated by the HIM posting date in the PMMIS and USRDS databases. Before this date, at least some of the hospitalizations will be missing. In most cases this date is shortly after the start of Medicare eligibility, or at the change from non-ESRD to ESRD eligibility. Both of these problems constrain the starting date for the period of complete data, and affect most dramatically patients who are in the program for shorter periods. There is likely to be some downward bias in the data because

hospitalization immediately following renal failure may not be counted.

In addition to the above limitations, the data from the HCFA PMMIS excluded one night stays before 1992. Also, Medicare rules require that for the first 21 months Medicare is the secondary payer after any other employer paid group health insurance which the patient may have. This may result in less complete hospitalization data in the first 21 months if the other insurer covers the complete cost of the hospitalization. The completeness of the hospitalization data is, and will continue to be, another quality control issue for USRDS. The limited comparisons that have been made to date between USRDS data and actual hospital records have produced mixed and inconclusive results.

hispanic origin: 1989. U.S. Government Printing Office, Washington, DC, 1990.

U.S. Bureau of the Census, Current Population Reports, Series P25, No. 1086, Population Estimates and Projections, U.S. Government Printing Office, Washington, DC, 1992.

U.S. Bureau of the Census, 1980 Census of Population, General Population Characteristics, United States Summary, PC801B1, U.S. Government Printing Office, Washington, DC, 1983.

U.S. Bureau of the Census, 1990 Census of Population and Housing: Modified Age/Race, Sex, and Hispanic Origin (MARS) State and County File, Data Users Services Division, The Bureau of the Census, Washington, DC, 1992.

Wolfe RA, Gaylin DS, Port FK, Held PJ, Wood CL. Using USRDS generated mortality tables to compare local ESRD mortality rates to national rates. *Kidney Int* 1992; 42:991-996.

---

## References

Fleiss JL. *Statistical Methods for Rates and Proportions*, second edition. New York: John Wiley & Sons, 1981.

Gross AJ, Clark VA. *Survival Distributions: Reliability Applications in the Biomedical Sciences*. New York: John Wiley & Sons. 1975.

Kaplan EL, Meier P. Nonparametric Estimation from Incomplete Observations. *J Am Statist Assoc* 1958; 53:457-481.

SAS Institute Inc. *SAS User's Guide: Statistics*, version 5 edition. Cary, NC: SAS Institute Inc. 1985.

U.S. Bureau of the Census, Current Population Reports, Series P25, No. 1045, United States population estimates by age, sex, race, and Hispanic origin: 1980 to 1988, U.S. Government Printing Office, Washington, DC, 1990.

U.S. Bureau of the Census, Current Population Reports, Series P25, No. 1057, U.S. population estimates by age, sex, race, and

